

Gwendolen A Jull

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8459424/gwendolen-a-jull-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

311
papers

19,484
citations

73
h-index

130
g-index

416
ext. papers

21,515
ext. citations

3.2
avg, IF

6.78
L-index

#	Paper	IF	Citations
311	The temporal behaviour of migraine related neck pain does not inform on the origin of neck pain: An observational study.. <i>Musculoskeletal Science and Practice</i> , 2022 , 58, 102522	2.4	1
310	Classification criteria for cervical radiculopathy: An international e-Delphi study. <i>Musculoskeletal Science and Practice</i> , 2022 , 61, 102596	2.4	0
309	Deep and superficial cervical muscles respond differently to unstable motor skill tasks. <i>Human Movement Science</i> , 2021 , 80, 102893	2.4	0
308	Altered neuromuscular activity and postural stability during standing balance tasks in persons with non-specific neck pain. <i>Journal of Electromyography and Kinesiology</i> , 2021 , 61, 102608	2.5	
307	Neck pain associated with migraine does not necessarily reflect cervical musculoskeletal dysfunction. <i>Headache</i> , 2021 , 61, 882-894	4.2	6
306	Die Hands-on/Hands-off-Debatte. <i>Muskuloskelettale Physiotherapie</i> , 2021 , 25, 117-124		
305	Cervical musculoskeletal impairments in migraine. <i>Archives of Physiotherapy</i> , 2021 , 11, 27	2.5	0
304	Implementation of a novel stratified PATHway of CarE for common musculoskeletal (MSK) conditions in primary care: protocol for a multicentre pragmatic randomised controlled trial (the PACE MSK trial). <i>BMJ Open</i> , 2021 , 11, e057705	3	
303	Response rate and comparison of clinical features associated with positive or negative responses to a scapular positioning test in patients with neck pain and altered scapular alignment: a cross-sectional study. <i>BMJ Open</i> , 2021 , 11, e057459	3	1
302	Validation of a clinical examination to differentiate a cervicogenic source of headache: a diagnostic prediction model using controlled diagnostic blocks. <i>BMJ Open</i> , 2020 , 10, e035245	3	11
301	Low back pain websites do not meet the needs of consumers: A study of online resources at three time points. <i>Health Information Management Journal</i> , 2020 , 49, 137-149	2.6	4
300	Outcome measures for assessing the effectiveness of non-pharmacological interventions in frequent episodic or chronic migraine: a Delphi study. <i>BMJ Open</i> , 2020 , 10, e029855	3	20
299	Cervical musculoskeletal impairments in migraine and tension type headache: A systematic review and meta-analysis. <i>Musculoskeletal Science and Practice</i> , 2019 , 42, 67-83	2.4	38
298	Motor cortex representation of deep and superficial neck flexor muscles in individuals with and without neck pain. <i>Human Brain Mapping</i> , 2019 , 40, 2759-2770	5.9	8
297	Self-reported sinus headaches are associated with neck pain and cervical musculoskeletal dysfunction: a preliminary observational case control study. <i>Journal of Manual and Manipulative Therapy</i> , 2019 , 27, 245-252	1.6	
296	A Lawyer With Whiplash 2019 , 241-258		
295	Recommendations For Core Outcome Domain Set For Whiplash-Associated Disorders (CATWAD). <i>Clinical Journal of Pain</i> , 2019 , 35, 727-736	3.5	10

294	Essential key messages about diagnosis, imaging, and self-care for people with low back pain: a modified Delphi study of consumer and expert opinions. <i>Pain</i> , 2019 , 160, 2787-2797	8	11
293	Neck muscle vibration produces diverse responses in balance and gait speed between individuals with and without neck pain. <i>Musculoskeletal Science and Practice</i> , 2018 , 35, 25-29	2.4	10
292	Benefits and Threats to Using Social Media for Presenting and Implementing Evidence. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2018 , 48, 3-7	4.2	4
291	Effects of local treatment with and without sensorimotor and balance exercise in individuals with neck pain: protocol for a randomized controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2018 , 19, 48	2.8	10
290	Psychological Features and Their Relationship to Movement-Based Subgroups in People Living With Low Back Pain. <i>Archives of Physical Medicine and Rehabilitation</i> , 2018 , 99, 121-128	2.8	7
289	Effects of deep cervical flexor training on impaired physiological functions associated with chronic neck pain: a systematic review. <i>BMC Musculoskeletal Disorders</i> , 2018 , 19, 415	2.8	35
288	Cervical musculoskeletal dysfunction in headache: How should it be defined?. <i>Musculoskeletal Science and Practice</i> , 2018 , 38, 148-150	2.4	12
287	Effectiveness of physiotherapy for seniors with recurrent headaches associated with neck pain and dysfunction: a randomized controlled trial. <i>Spine Journal</i> , 2017 , 17, 46-55	4	13
286	Low Pain Catastrophization and Disability Predict Successful Outcome to Radiofrequency Neurotomy in Individuals with Chronic Whiplash. <i>Pain Practice</i> , 2016 , 16, 311-9	3	11
285	Response to Letter to the Editor re: 'Movement-based subgrouping in low back pain: synergy and divergence in approaches'. <i>Physiotherapy</i> , 2016 , 102, e3	3	
284	Neck muscle function in violinists/violists with and without neck pain. <i>Clinical Rheumatology</i> , 2016 , 35, 1045-51	3.9	16
283	Discord Between Approaches to Spinal and Extremity Disorders: Is It Logical?. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2016 , 46, 938-941	4.2	8
282	Physical Therapist-Delivered Pain Coping Skills Training and Exercise for Knee Osteoarthritis: Randomized Controlled Trial. <i>Arthritis Care and Research</i> , 2016 , 68, 590-602	4.7	87
281	Does increased superficial neck flexor activity in the craniocervical flexion test reflect reduced deep flexor activity in people with neck pain?. <i>Manual Therapy</i> , 2016 , 25, 43-7		43
280	Movement-based subgrouping in low back pain: synergy and divergence in approaches. <i>Physiotherapy</i> , 2016 , 102, 159-69	3	25
279	International consensus on the most useful physical examination tests used by physiotherapists for patients with headache: A Delphi study. <i>Manual Therapy</i> , 2016 , 23, 17-24		45
278	Mechanical properties of the trapezius during scapular elevation in people with chronic whiplash associated disorders--A case-control ultrasound speckle tracking analysis. <i>Manual Therapy</i> , 2016 , 21, 177-82		9
277	Designing an online resource for people with low back pain: health-care provider perspectives. <i>Australian Journal of Primary Health</i> , 2016 , 22, 159-166	1.4	7

276	Can a professional development workshop with follow-up alter practitioner behaviour and outcomes for neck pain patients? A randomised controlled trial. <i>Manual Therapy</i> , 2016 , 25, 87-93		8
275	Is tailored treatment superior to non-tailored treatment for pain and disability in women with non-specific neck pain? A randomized controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2016 , 17, 408	2.8	10
274	Implementing a self-management intervention for people with a chronic compensable musculoskeletal injury in a workers compensation context: a process evaluation. <i>Journal of Occupational Rehabilitation</i> , 2015 , 25, 412-22	3.6	
273	Morphological changes in the cervical muscles of women with chronic whiplash can be modified with exercise-A pilot study. <i>Muscle and Nerve</i> , 2015 , 52, 772-9	3.4	29
272	A profession-wide collaboration to embed role-play simulation into Australian entry-level physiotherapy clinical training. <i>Physiotherapy</i> , 2015 , 101, e1047	3	5
271	External validation of a clinical prediction rule to predict full recovery and ongoing moderate/severe disability following acute whiplash injury. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2015 , 45, 242-50	4.2	54
270	Modulation of Cervical Facet Joint Nociception and Pain Attenuates Physical and Psychological Features of Chronic Whiplash: A Prospective Study. <i>PM and R</i> , 2015 , 7, 913-921	2.2	8
269	Comprehensive physiotherapy exercise programme or advice for chronic whiplash (PROMISE): a pragmatic randomised controlled trial. <i>Physiotherapy</i> , 2015 , 101, e1001-e1002	3	2
268	Preliminary evaluation of dorsal muscle activity during resisted cervical extension in patients with longstanding pain and disability following anterior cervical decompression and fusion surgery. <i>Physiotherapy</i> , 2015 , 101, 69-74	3	10
267	Inter-and intra-tester reliability of a battery of cervical movement control dysfunction tests. <i>Manual Therapy</i> , 2015 , 20, 570-9		8
266	Does posture of the cervical spine influence dorsal neck muscle activity when lifting?. <i>Manual Therapy</i> , 2014 , 19, 32-6		19
265	Physical therapist-delivered cognitive-behavioral therapy: a qualitative study of physical therapists' perceptions and experiences. <i>Physical Therapy</i> , 2014 , 94, 197-209	3.3	62
264	Commentary to: Surgery for cervical radiculopathy followed by physiotherapy may resolve symptoms faster than physiotherapy alone, but with few differences at two years. <i>Journal of Physiotherapy</i> , 2014 , 60, 109	2.9	
263	Enhancing the vocational outcomes of people with chronic disabilities caused by a musculoskeletal condition: development and evaluation of content of self-management training modules. <i>Work</i> , 2014 , 49, 455-64	1.6	4
262	A speckle tracking application of ultrasound to evaluate activity of multilayered cervical muscles. <i>Journal of Rehabilitation Medicine</i> , 2014 , 46, 662-7	3.4	4
261	Information needs of people with low back pain for an online resource: a qualitative study of consumer views. <i>Disability and Rehabilitation</i> , 2014 , 36, 1085-91	2.4	18
260	Can physical therapists deliver a pain coping skills program? An examination of training processes and outcomes. <i>Physical Therapy</i> , 2014 , 94, 1443-54	3.3	28
259	Differential changes in muscle composition exist in traumatic and nontraumatic neck pain. <i>Spine</i> , 2014 , 39, 39-47	3.3	70

258	Cervical radiofrequency neurotomy reduces central hyperexcitability and improves neck movement in individuals with chronic whiplash. <i>Pain Medicine</i> , 2014 , 15, 128-41	2.8	25
257	A description of neck motor performance, neck pain, fatigue, and mental effort while driving in a sample with chronic whiplash-associated disorders. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2014 , 93, 665-74	2.6	10
256	Derivation of a clinical decision guide in the diagnosis of cervical facet joint pain. <i>Archives of Physical Medicine and Rehabilitation</i> , 2014 , 95, 1695-701	2.8	31
255	Comprehensive physiotherapy exercise programme or advice for chronic whiplash (PROMISE): a pragmatic randomised controlled trial. <i>Lancet, The</i> , 2014 , 384, 133-41	4.0	106
254	Cervical radiofrequency neurotomy reduces psychological features in individuals with chronic whiplash symptoms. <i>Pain Physician</i> , 2014 , 17, 265-74	1.8	9
253	Immediate and short-term effects of Mulligan's mobilization with movement on knee pain and disability associated with knee osteoarthritis--a prospective case series. <i>Physiotherapy Theory and Practice</i> , 2013 , 29, 87-95	1.5	28
252	Applying principles of self-management to facilitate workers to return to or remain at work with a chronic musculoskeletal condition. <i>Manual Therapy</i> , 2013 , 18, 274-80		14
251	Management of acute whiplash: a randomized controlled trial of multidisciplinary stratified treatments. <i>Pain</i> , 2013 , 154, 1798-1806	8	62
250	Cervical muscle activity during loaded arm lifts in patients 10 years postsurgery for cervical disc disease. <i>Journal of Manipulative and Physiological Therapeutics</i> , 2013 , 36, 292-9	1.3	9
249	Patients with non-specific neck disorders commonly report upper limb disability. <i>Manual Therapy</i> , 2013 , 18, 492-7		23
248	Assessment of driving-related performance in chronic whiplash using an advanced driving simulator. <i>Accident Analysis and Prevention</i> , 2013 , 60, 5-14	6.1	8
247	People seeking treatment for a new episode of neck pain typically have rapid improvement in symptoms: an observational study. <i>Journal of Physiotherapy</i> , 2013 , 59, 31-7	2.9	26
246	Characteristics of a new episode of neck pain. <i>Manual Therapy</i> , 2013 , 18, 254-7		14
245	Intrarater and interrater reliability of select clinical tests in patients referred for diagnostic facet joint blocks in the cervical spine. <i>Archives of Physical Medicine and Rehabilitation</i> , 2013 , 94, 1628-34	2.8	33
244	A comparison of physical and psychological features of responders and non-responders to cervical facet blocks in chronic whiplash. <i>BMC Musculoskeletal Disorders</i> , 2013 , 14, 313	2.8	29
243	Response to Commentary on Our Article Titled Intrarater and Interrater Reliability of Select Clinical Tests in Patients Referred for Diagnostic Facet Joint Blocks in the Cervical Spine— <i>Archives of Physical Medicine and Rehabilitation</i> , 2013 , 94, 1638-1640	2.8	
242	Sensory and sensorimotor features in violinists and violists with neck pain. <i>Archives of Physical Medicine and Rehabilitation</i> , 2013 , 94, 2523-2528	2.8	16
241	Editors' response. <i>Manual Therapy</i> , 2013 , 18, e12		

240	Neck pain driving index: appropriateness of the rating scale and unidimensionality of the strategic, tactical, and operational levels. <i>Archives of Physical Medicine and Rehabilitation</i> , 2013 , 94, 1842-6	2.8	6
239	Baseline characteristics of patients with nerve-related neck and arm pain predict the likely response to neural tissue management. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2013 , 43, 379-91	4.2	14
238	Registrierung Klinischer Studien. <i>Physioscience</i> , 2013 , 9, 47-49	0.3	
237	Contributions of physical and cognitive impairments to self-reported driving difficulty in chronic whiplash-associated disorders. <i>Spine</i> , 2013 , 38, 1554-60	3.3	17
236	Simulation can contribute a part of cardiorespiratory physiotherapy clinical education: two randomized trials. <i>Simulation in Healthcare</i> , 2013 , 8, 32-42	2.8	61
235	Clinical trial registration in physiotherapy journals: recommendations from the international society of physiotherapy journal editors. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2013 , 65, 109-15	0.8	2
234	L'enregistrement des essais cliniques dans les revues de physiothérapie: recommandations de l'International Society of Physiotherapy Journal Editors. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2013 , 65, 109-15	0.8	
233	Is there a difference in the pattern of muscle activity when performing neck exercises with a guild board versus a pulley?. <i>Journal of Rehabilitation Medicine</i> , 2013 , 45, 900-5	3.4	6
232	Recommendations From the International Society of Physiotherapy Journal Editors: Clinical Trial Registration in Physiotherapy Journals. <i>Journal, Physical Therapy Education</i> , 2013 , 27, 7-9	0.6	
231	Neural tissue management provides immediate clinically relevant benefits without harmful effects for patients with nerve-related neck and arm pain: a randomised trial. <i>Journal of Physiotherapy</i> , 2012 , 58, 23-31	2.9	70
230	Screening of patients suitable for diagnostic cervical facet joint blocks--a role for physiotherapists. <i>Manual Therapy</i> , 2012 , 17, 180-3		8
229	As we move into 2012. <i>Manual Therapy</i> , 2012 , 17, 1		1
228	Can a functional postural exercise improve performance in the cranio-cervical flexion test?--a preliminary study. <i>Manual Therapy</i> , 2012 , 17, 219-24		29
227	Can simulation replace part of clinical time? Two parallel randomised controlled trials. <i>Medical Education</i> , 2012 , 46, 657-67	3.7	115
226	Physiotherapy movement based classification approaches to low back pain: comparison of subgroups through review and developer/expert survey. <i>BMC Musculoskeletal Disorders</i> , 2012 , 13, 24	2.8	72
225	The influence of neck pain on sensorimotor function in the elderly. <i>Archives of Gerontology and Geriatrics</i> , 2012 , 55, 667-72	4	32
224	The validity of upper-limb neurodynamic tests for detecting peripheral neuropathic pain. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2012 , 42, 413-24	4.2	73
223	Minimum repetitions for stable measures of visual dependency using the dot version of the computer-based Rod-Frame test. <i>Manual Therapy</i> , 2012 , 17, 466-9		14

222 Response to Jones. *Manual Therapy*, **2012**, 17, e15

221 Assessment and validation of prognostic models for poor functional recovery 12 months after whiplash injury: a multicentre inception cohort study. *Pain*, **2012**, 153, 1727-1734 8 43

220 Training mode-dependent changes in motor performance in neck pain. *Archives of Physical Medicine and Rehabilitation*, **2012**, 93, 1225-33 2.8 28

219 A physiotherapist-delivered integrated exercise and pain coping skills training intervention for individuals with knee osteoarthritis: a randomised controlled trial protocol. *BMC Musculoskeletal Disorders*, **2012**, 13, 129 2.8 23

218 Validity and reliability of the perceived deficit questionnaire to assess cognitive symptoms in people with chronic whiplash-associated disorders. *Archives of Physical Medicine and Rehabilitation*, **2012**, 93, 1774-81 2.8 20

217 The Neck Pain Driving Index (NPDI) for chronic whiplash-associated disorders: development, reliability, and validity assessment. *Spine Journal*, **2012**, 12, 912-20.e1 4 13

216 Altered trapezius muscle behavior in individuals with neck pain and clinical signs of scapular dysfunction. *Journal of Manipulative and Physiological Therapeutics*, **2012**, 35, 346-53 1.3 48

215 Comparing outcomes of medial parapatellar and subvastus approaches in total knee arthroplasty: a randomized controlled trial. *Journal of Arthroplasty*, **2012**, 27, 347-353.e1 4.4 19

214 A comparison of patellar vascularity between the medial parapatellar and subvastus approaches in total knee arthroplasty. *Journal of Arthroplasty*, **2012**, 27, 1123-7.e1 4.4 10

213 Magnetic resonance imaging changes in the size and shape of the oropharynx following acute whiplash injury. *Journal of Orthopaedic and Sports Physical Therapy*, **2012**, 42, 912-8 4.2 11

212 The change in deep cervical flexor activity after training is associated with the degree of pain reduction in patients with chronic neck pain. *Clinical Journal of Pain*, **2012**, 28, 628-34 3.5 80

211 Driving with a chronic whiplash-associated disorder: a review of patients' perspectives. *Archives of Physical Medicine and Rehabilitation*, **2011**, 92, 106-10 2.8 15

210 Is there altered activity of the extensor muscles in chronic mechanical neck pain? A functional magnetic resonance imaging study. *Archives of Physical Medicine and Rehabilitation*, **2011**, 92, 929-34 2.8 66

209 A novel protocol to develop a prediction model that identifies patients with nerve-related neck and arm pain who benefit from the early introduction of neural tissue management. *Contemporary Clinical Trials*, **2011**, 32, 760-70 2.3 13

208 Considerations in the physical rehabilitation of patients with whiplash-associated disorders. *Spine*, **2011**, 36, S286-91 3.3 8

207 Toward optimal early management after whiplash injury to lessen the rate of transition to chronicity: discussion paper 5. *Spine*, **2011**, 36, S335-42 3.3 24

206 Association between intensity of pain and impairment in onset and activation of the deep cervical flexors in patients with persistent neck pain. *Clinical Journal of Pain*, **2011**, 27, 309-14 3.5 61

205 A radiographic analysis of the influence of initial neck posture on cervical segmental movement at end-range extension in asymptomatic subjects. *Manual Therapy*, **2011**, 16, 74-9 10

204	Normal kinematics of the upper cervical spine during the Flexion-Rotation Test - In vivo measurements using magnetic resonance imaging. <i>Manual Therapy</i> , 2011 , 16, 167-71	56	
203	Head eye co-ordination and gaze stability in subjects with persistent whiplash associated disorders. <i>Manual Therapy</i> , 2011 , 16, 252-7	52	
202	The relationship between superficial muscle activity during the craniocervical flexion test and clinical features in patients with chronic neck pain. <i>Manual Therapy</i> , 2011 , 16, 452-5	47	
201	Absence of the inferior portion of the trapezius muscle in three family members. <i>Manual Therapy</i> , 2011 , 16, 629-35	2	
200	Does the region of pain influence the presence of sensorimotor disturbances in neck pain disorders?. <i>Manual Therapy</i> , 2011 , 16, 636-40	26	
199	The case of the missing lower trapezius muscle. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2011 , 41, 614	4.2	
198	Internet-based outpatient telerehabilitation for patients following total knee arthroplasty: a randomized controlled trial. <i>Journal of Bone and Joint Surgery - Series A</i> , 2011 , 93, 113-20	5.6	212
197	The temporal development of fatty infiltrates in the neck muscles following whiplash injury: an association with pain and posttraumatic stress. <i>PLoS ONE</i> , 2011 , 6, e21194	3.7	81
196	Interactive effects from self-reported physical and psychosocial factors in the workplace on neck pain and disability in female office workers. <i>Ergonomics</i> , 2010 , 53, 502-13	2.9	26
195	Craniocervical orientation affects muscle activation when exercising the cervical extensors in healthy subjects. <i>Archives of Physical Medicine and Rehabilitation</i> , 2010 , 91, 1418-22	2.8	38
194	A randomized controlled trial comparing manipulation with mobilization for recent onset neck pain. <i>Archives of Physical Medicine and Rehabilitation</i> , 2010 , 91, 1313-8	2.8	86
193	Magnetic resonance imaging findings of fatty infiltrate in the cervical flexors in chronic whiplash. <i>Spine</i> , 2010 , 35, 948-54	3.3	89
192	A comparison of craniocervical and cervicothoracic muscle strength in healthy individuals. <i>Journal of Applied Biomechanics</i> , 2010 , 26, 400-6	1.2	22
191	The effect of a scapular postural correction strategy on trapezius activity in patients with neck pain. <i>Manual Therapy</i> , 2010 , 15, 562-6	73	
190	Does self-management for return to work increase the effectiveness of vocational rehabilitation for chronic compensated musculoskeletal disorders? Protocol for a randomised controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2010 , 11, 115	2.8	7
189	Systematic review of medial parapatellar and subvastus approaches in total knee arthroplasty. <i>Journal of Arthroplasty</i> , 2010 , 25, 728-34	4.4	17
188	Invited commentary. <i>Physical Therapy</i> , 2009 , 89, 48-50	3.3	1
187	Psychological, cognitive and quality of life features in the elderly with chronic headache. <i>Gerontology</i> , 2009 , 55, 683-93	5.5	14

186	Bungsansatz bei HWS-Störungen: Wie fließen die Ergebnisse der Forschung in die Praxis ein? <i>Manuelletherapie</i> , 2009, 13, 110-116	0.1	0
185	Head repositioning accuracy to neutral: a comparative study of error calculation. <i>Manual Therapy</i> , 2009, 14, 110-4		31
184	Performance in the cranio-cervical flexion test is altered in elderly subjects. <i>Manual Therapy</i> , 2009, 14, 475-9		20
183	Cervical musculoskeletal impairment is common in elders with headache. <i>Manual Therapy</i> , 2009, 14, 636-41		24
182	The effect of therapeutic exercise on activation of the deep cervical flexor muscles in people with chronic neck pain. <i>Manual Therapy</i> , 2009, 14, 696-701		196
181	A randomised clinical trial of a comprehensive exercise program for chronic whiplash: trial protocol. <i>BMC Musculoskeletal Disorders</i> , 2009, 10, 149	2.8	16
180	The clinical presentation of chronic whiplash and the relationship to findings of MRI fatty infiltrates in the cervical extensor musculature: a preliminary investigation. <i>European Spine Journal</i> , 2009, 18, 1371-87	2.7	38
179	Head eye co-ordination using simultaneous measurement of eye in head and head in space movements: potential for use in subjects with a whiplash injury. <i>Journal of Clinical Monitoring and Computing</i> , 2009, 23, 31-40	2	19
178	Contribution of individual, workplace, psychosocial and physiological factors to neck pain in female office workers. <i>European Journal of Pain</i> , 2009, 13, 985-91	3.7	19
177	Muscle dysfunction in cervical spine pain: implications for assessment and management. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2009, 39, 324-33	4.2	132
176	Physiotherapy management of cervicogenic headache: Part 1 2009, 169-180		
175	Physiotherapy management of cervicogenic headachePart 2 2009, 181-194		
174	Widespread sensory hypersensitivity is not a feature of chronic headache in elders. <i>Clinical Journal of Pain</i> , 2009, 25, 699-704	3.5	20
173	The influence of cervical traction, compression, and spurling test on cervical intervertebral foramen size. <i>Spine</i> , 2009, 34, 1658-62	3.3	22
172	Standing balance: a comparison between idiopathic and whiplash-induced neck pain. <i>Manual Therapy</i> , 2008, 13, 183-91		87
171	MRI study of the cross-sectional area for the cervical extensor musculature in patients with persistent whiplash associated disorders (WAD). <i>Manual Therapy</i> , 2008, 13, 258-65		83
170	The influence of neck pain on balance and gait parameters in community-dwelling elders. <i>Manual Therapy</i> , 2008, 13, 317-24		45
169	Training the cervical muscles with prescribed motor tasks does not change muscle activation during a functional activity. <i>Manual Therapy</i> , 2008, 13, 507-12		37

168	Alterations in Cervical Muscle Function in Neck Pain 2008 , 41-58		2
167	Sensory Manifestations of Neck Pain 2008 , 5-19		
166	Structure and Function of the Cervical Region 2008 , 21-39		
165	The Cervical Spine and Sensorimotor Control 2008 , 59-71		
164	Whiplash-associated Disorders 2008 , 101-115		2
163	Cervicogenic Headache 2008 , 117-130		1
162	Differential Diagnosis of Cervicobrachial Pain 2008 , 131-143		1
161	Clinical Assessment 2008 , 145-153		
160	Clinical Assessment: Physical Examination of the Cervical Region 2008 , 155-187		2
159	Principles of Management of Cervical Disorders 2008 , 189-206		1
158	Clinical assessment of the deep cervical flexor muscles: the craniocervical flexion test. <i>Journal of Manipulative and Physiological Therapeutics</i> , 2008 , 31, 525-33	1.3	257
157	Self-reported driving habits in subjects with persistent whiplash-associated disorder: relationship to sensorimotor and psychologic features. <i>Archives of Physical Medicine and Rehabilitation</i> , 2008 , 89, 1097-102	2.8	28
156	Comparison of sensorimotor disturbance between subjects with persistent whiplash-associated disorder and subjects with vestibular pathology associated with acoustic neuroma. <i>Archives of Physical Medicine and Rehabilitation</i> , 2008 , 89, 522-30	2.8	59
155	MRI analysis of the size and shape of the oropharynx in chronic whiplash. <i>Otolaryngology - Head and Neck Surgery</i> , 2008 , 138, 747-51	5.5	8
154	Quantitative sensory measures distinguish office workers with varying levels of neck pain and disability. <i>Pain</i> , 2008 , 137, 257-265	8	69
153	Therapeutic Exercise for Cervical Disorders 2008 , 207-229		4
152	Fatty infiltrate in the cervical extensor muscles is not a feature of chronic, insidious-onset neck pain. <i>Clinical Radiology</i> , 2008 , 63, 681-7	2.9	98
151	Neck movement and muscle activity characteristics in female office workers with neck pain. <i>Spine</i> , 2008 , 33, 555-63	3.3	117

150	Disturbances in Postural Stability, Head and Eye Movement Control in Cervical Disorders 2008 , 73-86		
149	Psychological and Psychosocial Factors in Neck Pain 2008 , 87-99		
148	Alterations in cervical muscle activity in functional and stressful tasks in female office workers with neck pain. <i>European Journal of Applied Physiology</i> , 2008 , 103, 253-64	3.4	70
147	Associations between individual and workplace risk factors for self-reported neck pain and disability among female office workers. <i>Applied Ergonomics</i> , 2008 , 39, 171-82	4.2	50
146	Effect of neck exercise on sitting posture in patients with chronic neck pain. <i>Physical Therapy</i> , 2007 , 87, 408-17	3.3	240
145	Specific therapeutic exercise of the neck induces immediate local hypoalgesia. <i>Journal of Pain</i> , 2007 , 8, 832-9	5.2	97
144	Magnetic resonance imaging study of cross-sectional area of the cervical extensor musculature in an asymptomatic cohort. <i>Clinical Anatomy</i> , 2007 , 20, 35-40	2.5	51
143	Retraining cervical joint position sense: the effect of two exercise regimes. <i>Journal of Orthopaedic Research</i> , 2007 , 25, 404-12	3.8	167
142	Cervical musculoskeletal impairment in frequent intermittent headache. Part 1: Subjects with single headaches. <i>Cephalalgia</i> , 2007 , 27, 793-802	6.1	160
141	Cervical musculoskeletal impairment in frequent intermittent headache. Part 2: subjects with concurrent headache types. <i>Cephalalgia</i> , 2007 , 27, 891-8	6.1	57
140	Efficacy of manipulation for non-specific neck pain of recent onset: design of a randomised controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2007 , 8, 18	2.8	14
139	Cranio-cervical flexor muscle impairment at maximal, moderate, and low loads is a feature of neck pain. <i>Manual Therapy</i> , 2007 , 12, 34-9		87
138	Recruitment of the deep cervical flexor muscles during a postural-correction exercise performed in sitting. <i>Manual Therapy</i> , 2007 , 12, 139-43		98
137	Wireless orientation sensors: their suitability to measure head movement for neck pain assessment. <i>Manual Therapy</i> , 2007 , 12, 380-5		43
136	Equine physiotherapy: a comparative view of the science underlying the profession. <i>Equine Veterinary Journal</i> , 2007 , 39, 90-4	2.4	26
135	Muscle specificity in tests of cervical flexor muscle performance. <i>Journal of Electromyography and Kinesiology</i> , 2007 , 17, 35-40	2.5	70
134	Interventions to prevent back pain and back injury in nurses: a systematic review. <i>Occupational and Environmental Medicine</i> , 2007 , 64, 642-50	2.1	133
133	Specificity in retraining craniocervical flexor muscle performance. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2007 , 37, 3-9	4.2	55

132	Conservative treatments for tennis elbow do subgroups of patients respond differently?. <i>Rheumatology</i> , 2007 , 46, 1601-5	3.9	46
131	Does the presence of sensory hypersensitivity influence outcomes of physical rehabilitation for chronic whiplash?--A preliminary RCT. <i>Pain</i> , 2007 , 129, 28-34	8	183
130	Interaction of psychosocial risk factors explain increased neck problems among female office workers. <i>Pain</i> , 2007 , 129, 311-320	8	26
129	Cyclic pneumatic soft-tissue compression enhances recovery following fracture of the distal radius: a randomised controlled trial. <i>Australian Journal of Physiotherapy</i> , 2007 , 53, 247-52		20
128	No model of clinical education for physiotherapy students is superior to another: a systematic review. <i>Australian Journal of Physiotherapy</i> , 2007 , 53, 19-28		111
127	The relationship of cervical joint position error to balance and eye movement disturbances in persistent whiplash. <i>Manual Therapy</i> , 2006 , 11, 99-106		138
126	Clinical tests of musculoskeletal dysfunction in the diagnosis of cervicogenic headache. <i>Manual Therapy</i> , 2006 , 11, 118-29		186
125	Diagnosis of Cervicogenic Headache. <i>Journal of Manual and Manipulative Therapy</i> , 2006 , 14, 136-138	1.6	3
124	Cyclic pneumatic soft-tissue compression accelerates the union of distal radial osteotomies in an ovine model. <i>Journal of Bone and Joint Surgery: British Volume</i> , 2006 , 88, 411-5		12
123	Balance, mobility and gaze stability deficits remain following surgical removal of vestibular schwannoma (acoustic neuroma): an observational study. <i>Australian Journal of Physiotherapy</i> , 2006 , 52, 211-6		21
122	An endurance-strength training regime is effective in reducing myoelectric manifestations of cervical flexor muscle fatigue in females with chronic neck pain. <i>Clinical Neurophysiology</i> , 2006 , 117, 828-37	4.3	114
121	Mobilisation with movement and exercise, corticosteroid injection, or wait and see for tennis elbow: randomised trial. <i>BMJ, The</i> , 2006 , 333, 939	5.9	314
120	Further evaluation of an EMG technique for assessment of the deep cervical flexor muscles. <i>Journal of Electromyography and Kinesiology</i> , 2006 , 16, 621-8	2.5	54
119	Physical and psychological factors maintain long-term predictive capacity post-whiplash injury. <i>Pain</i> , 2006 , 122, 102-8	8	212
118	Fatty infiltration in the cervical extensor muscles in persistent whiplash-associated disorders: a magnetic resonance imaging analysis. <i>Spine</i> , 2006 , 31, E847-55	3.3	192
117	Applications in Rehabilitation Medicine and Related Fields 2005 , 403-433		
116	Is the method of signal analysis and test selection important for measuring standing balance in subjects with persistent whiplash?. <i>Gait and Posture</i> , 2005 , 21, 395-402	2.6	26
115	Magnetic resonance imaging analysis of the upper cervical spine extensor musculature in an asymptomatic cohort: an index of fat within muscle. <i>Clinical Radiology</i> , 2005 , 60, 355-63	2.9	49

114	Physical and psychological factors predict outcome following whiplash injury. <i>Pain</i> , 2005 , 114, 141-8	8	279
113	Do dorsal head contact forces have the potential to identify impairment during graded craniocervical flexor muscle contractions?. <i>Archives of Physical Medicine and Rehabilitation</i> , 2005 , 86, 1763-6	2.8	7
112	Smooth pursuit neck torsion test in whiplash-associated disorders: relationship to self-reports of neck pain and disability, dizziness and anxiety. <i>Journal of Rehabilitation Medicine</i> , 2005 , 37, 219-23	3.4	77
111	A New Method of Isometric Dynamometry for the Craniocervical Flexor Muscles. <i>Physical Therapy</i> , 2005 , 85, 556-564	3.3	38
110	Effect of cyclic pneumatic soft tissue compression on simulated distal radius fractures. <i>Clinical Orthopaedics and Related Research</i> , 2005 , 183-8	2.2	8
109	Widespread sensory hypersensitivity is a feature of chronic whiplash-associated disorder but not chronic idiopathic neck pain. <i>Clinical Journal of Pain</i> , 2005 , 21, 175-81	3.5	216
108	Predictors of responsiveness to physiotherapy management of cervicogenic headache. <i>Cephalalgia</i> , 2005 , 25, 101-8	6.1	46
107	Standing balance in persistent whiplash: a comparison between subjects with and without dizziness. <i>Journal of Rehabilitation Medicine</i> , 2005 , 37, 224-9	3.4	131
106	A new method of isometric dynamometry for the craniocervical flexor muscles. <i>Physical Therapy</i> , 2005 , 85, 556-64	3.3	8
105	Headache in a mature athlete 2004 , 135-148		
104	Impairment in the cervical flexors: a comparison of whiplash and insidious onset neck pain patients. <i>Manual Therapy</i> , 2004 , 9, 89-94		305
103	Feedforward activity of the cervical flexor muscles during voluntary arm movements is delayed in chronic neck pain. <i>Experimental Brain Research</i> , 2004 , 157, 43-8	2.3	218
102	Neck flexor muscle fatigue is side specific in patients with unilateral neck pain. <i>European Journal of Pain</i> , 2004 , 8, 71-7	3.7	71
101	Rehabilitation after total knee replacement via low-bandwidth telemedicine: the patient and therapist experience. <i>Journal of Telemedicine and Telecare</i> , 2004 , 10 Suppl 1, 85-7	6.8	43
100	Neuromuscular efficiency of the sternocleidomastoid and anterior scalene muscles in patients with chronic neck pain. <i>Disability and Rehabilitation</i> , 2004 , 26, 712-7	2.4	73
99	Response to Dr Ferrari's Letter to the Editor 'The clinical relevance of symptom amplification' <i>Pain</i> , 2004 , 107, 277-278	8	2
98	The clinical relevance of symptom amplification. <i>Pain</i> , 2004 , 107, 276	8	14
97	More on the clinical and scientific relevance of 'symptom amplification' and psychological factors in pain. <i>Pain</i> , 2004 , 110, 499-500; author reply 500-2	8	4

96	Response to Drs Greve and Bianchini. <i>Pain</i> , 2004 , 110, 501-502	8	1
95	Lack of correlation between sternocleidomastoid and scalene muscle fatigability and duration of symptoms in chronic neck pain patients. <i>Neurophysiologie Clinique</i> , 2004 , 34, 159-65	2.7	17
94	Spatio-temporal evaluation of neck muscle activation during postural perturbations in healthy subjects. <i>Journal of Electromyography and Kinesiology</i> , 2004 , 14, 463-74	2.5	34
93	Diagnosis of Cervical Disorders: Exploring a Mechanistic Approach. <i>Hong Kong Physiotherapy Journal</i> , 2004 , 22, 2-6	1	1
92	Characterization of acute whiplash-associated disorders. <i>Spine</i> , 2004 , 29, 182-8	3.3	129
91	Patients with neck pain demonstrate reduced electromyographic activity of the deep cervical flexor muscles during performance of the craniocervical flexion test. <i>Spine</i> , 2004 , 29, 2108-14	3.3	359
90	Patients with chronic neck pain demonstrate altered patterns of muscle activation during performance of a functional upper limb task. <i>Spine</i> , 2004 , 29, 1436-40	3.3	279
89	Measurement of Upper Cervical Flexion and Extension with the 3-Space Fastrak Measurement System: A Repeatability Study. <i>Journal of Manual and Manipulative Therapy</i> , 2003 , 11, 198-203	1.6	10
88	An Electromyographic Analysis of the Deep Cervical Flexor Muscles in Performance of Craniocervical Flexion. <i>Physical Therapy</i> , 2003 , 83, 899-906	3.3	147
87	Dizziness and unsteadiness following whiplash injury: characteristic features and relationship with cervical joint position error. <i>Journal of Rehabilitation Medicine</i> , 2003 , 35, 36-43	3.4	267
86	Relationship between cranio-cervical flexion range of motion and pressure change during the cranio-cervical flexion test. <i>Manual Therapy</i> , 2003 , 8, 92-6		71
85	Measuring range of active cervical rotation in a position of full head flexion using the 3D Fastrak measurement system: an intra-tester reliability study. <i>Manual Therapy</i> , 2003 , 8, 176-9		53
84	Can the Internet be used as a medium to evaluate knee angle?. <i>Manual Therapy</i> , 2003 , 8, 242-6		68
83	Justifying the on-going physiotherapy management of long-term patients. <i>Manual Therapy</i> , 2003 , 8, 254-6		13
82	Control subjects in whiplash studies. <i>Manual Therapy</i> , 2003 , 8, 52-3; author reply 54-5		1
81	Response to Drs Kwan and Friel. <i>Manual Therapy</i> , 2003 , 8, 54-55		
80	Cervicogenic headache: locus of control and success of treatment. <i>Headache</i> , 2003 , 43, 956-61	4.2	12
79	Recent advances in therapeutic exercise for the neck: implications for patients with head and neck pain. <i>Australian Endodontic Journal</i> , 2003 , 29, 138-42	1.7	29

78	Myoelectric manifestations of sternocleidomastoid and anterior scalene muscle fatigue in chronic neck pain patients. <i>Clinical Neurophysiology</i> , 2003 , 114, 488-95	4.3	131
77	The development of psychological changes following whiplash injury. <i>Pain</i> , 2003 , 106, 481-489	8	167
76	Development of motor system dysfunction following whiplash injury. <i>Pain</i> , 2003 , 103, 65-73	8	241
75	Sensory hypersensitivity occurs soon after whiplash injury and is associated with poor recovery. <i>Pain</i> , 2003 , 104, 509-517	8	365
74	Comment on Sterling, M., et al., Motor system dysfunction following whiplash injury, <i>Pain</i> 103 (2003) 65-73. <i>Pain</i> , 2003 , 105, 507	8	
73	Response to comment by Kwan and Friel. <i>Pain</i> , 2003 , 105, 508	8	
72	Low-bandwidth telerehabilitation for patients who have undergone total knee replacement: preliminary results. <i>Journal of Telemedicine and Telecare</i> , 2003 , 9 Suppl 2, S44-7	6.8	69
71	A study of five cervicocephalic relocation tests in three different subject groups. <i>Clinical Rehabilitation</i> , 2003 , 17, 768-74	3.3	125
70	The diagnostic reliability of Internet-based observational kinematic gait analysis. <i>Journal of Telemedicine and Telecare</i> , 2003 , 9 Suppl 2, S48-51	6.8	25
69	An electromyographic analysis of the deep cervical flexor muscles in performance of craniocervical flexion. <i>Physical Therapy</i> , 2003 , 83, 899-906	3.3	26
68	Responses to a clinical test of mechanical provocation of nerve tissue in whiplash associated disorder. <i>Manual Therapy</i> , 2002 , 7, 89-94		93
67	Repeatability of surface EMG variables in the sternocleidomastoid and anterior scalene muscles. <i>European Journal of Applied Physiology</i> , 2002 , 87, 542-9	3.4	75
66	Are cervical physical outcome measures influenced by the presence of symptomatology?. <i>Physiotherapy Research International</i> , 2002 , 7, 113-21	1.8	33
65	Reliability of detecting 'onset of pain' and 'submaximal pain' during neural provocation testing of the upper quadrant. <i>Physiotherapy Research International</i> , 2002 , 7, 146-56	1.8	116
64	Management of Cervicogenic Headache 2002 , 239-272		4
63	A randomized controlled trial of exercise and manipulative therapy for cervicogenic headache. <i>Spine</i> , 2002 , 27, 1835-43; discussion 1843	3.3	485
62	Pressure Pain Thresholds in Chronic Whiplash Associated Disorder: Further Evidence of Altered Central Pain Processing. <i>Journal of Musculoskeletal Pain</i> , 2002 , 10, 69-81		94
61	Physical outcome measurements via the Internet: reliability at two Internet speeds. <i>Journal of Telemedicine and Telecare</i> , 2002 , 8 Suppl 3, 50-52	6.8	33

60	Location of innervation zones of sternocleidomastoid and scalene muscles--a basis for clinical and research electromyography applications. <i>Clinical Neurophysiology</i> , 2002 , 113, 57-63	4.3	230
59	Manual therapy produces greater relief of neck pain than physiotherapy or general practitioner care. <i>Australian Journal of Physiotherapy</i> , 2002 , 48, 240-1		1
58	Use of high and low velocity cervical manipulative therapy procedures by Australian manipulative physiotherapists. <i>Australian Journal of Physiotherapy</i> , 2002 , 48, 189-93		45
57	Physiotherapy, a responsible profession to use cervical manipulation. Response to Refshauge et al. <i>Australian Journal of Physiotherapy</i> , 2002 , 48, 180-183		12
56	Cervicocephalic kinaesthesia: reliability of a new test approach. <i>Physiotherapy Research International</i> , 2001 , 6, 224-35	1.8	53
55	Co-activation of the abdominal and pelvic floor muscles during voluntary exercises. <i>Neurourology and Urodynamics</i> , 2001 , 20, 31-42	2.3	338
54	Mid-thoracic tenderness: a comparison of pressure pain threshold between spinal regions, in asymptomatic subjects. <i>Manual Therapy</i> , 2001 , 6, 34-9		40
53	Cervical mobilisation: concurrent effects on pain, sympathetic nervous system activity and motor activity. <i>Manual Therapy</i> , 2001 , 6, 72-81		325
52	The effect of musculoskeletal pain on motor activity and control. <i>Journal of Pain</i> , 2001 , 2, 135-45	5.2	138
51	Long-term effects of specific stabilizing exercises for first-episode low back pain. <i>Spine</i> , 2001 , 26, E243-8,3		567
50	Cervical range of motion discriminates between asymptomatic persons and those with whiplash. <i>Spine</i> , 2001 , 26, 2090-4	3.3	186
49	Pressure pain thresholds of upper limb peripheral nerve trunks in asymptomatic subjects. <i>Physiotherapy Research International</i> , 2000 , 5, 220-9	1.8	23
48	Deep Cervical Flexor Muscle Dysfunction in Whiplash. <i>Journal of Musculoskeletal Pain</i> , 2000 , 8, 143-154		171
47	Motor control problems in patients with spinal pain: a new direction for therapeutic exercise. <i>Journal of Manipulative and Physiological Therapeutics</i> , 2000 , 23, 115-117	1.3	88
46	Does strengthening the abdominal muscles prevent low back pain?. <i>Journal of Rheumatology</i> , 2000 , 27, 2286-8	4.1	4
45	Lumbar spine kinesthesia in patients with low back pain. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 1999 , 29, 294-9	4.2	61
44	Further clinical clarification of the muscle dysfunction in cervical headache. <i>Cephalalgia</i> , 1999 , 19, 179-85,1		218
43	Use of real-time ultrasound imaging for feedback in rehabilitation. <i>Manual Therapy</i> , 1998 , 3, 125-131		69

42	Breast surgery and neural tissue mechanosensitivity. <i>Australian Journal of Physiotherapy</i> , 1998 , 44, 31-37		10
41	Characterization of cervicogenic headache. <i>Physical Therapy Reviews</i> , 1998 , 3, 95-105	0.7	2
40	Electromyographic amplitude and frequency changes in the iliocostalis lumborum and multifidus muscles during a trunk holding test. <i>Physical Therapy</i> , 1997 , 77, 954-61	3.3	84
39	Inter-examiner reliability to detect painful upper cervical joint dysfunction. <i>Australian Journal of Physiotherapy</i> , 1997 , 43, 125-129		104
38	Active stabilisation training for screen based keyboard operators - a single case study. <i>Australian Journal of Physiotherapy</i> , 1997 , 43, 235-242		10
37	Upper trapezius muscle activity during the brachial plexus tension test in asymptomatic subjects. <i>Manual Therapy</i> , 1997 , 2, 144-149		53
36	Management of cervical headache. <i>Manual Therapy</i> , 1997 , 2, 182-190		68
35	Retrodiscal tissue of the temporomandibular joint: clinical anatomy and its role in diagnosis and treatment of arthropathies. <i>Manual Therapy</i> , 1997 , 2, 191-198		8
34	Toward a clinical test of lumbar spine kinesthesia. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 1996 , 24, 354-8	4.2	61
33	An in vivo study of the primary and coupled rotations of the thoracic spine. <i>Clinical Biomechanics</i> , 1996 , 11, 311-316	2.2	116
32	Multifidus muscle recovery is not automatic after resolution of acute, first-episode low back pain. <i>Spine</i> , 1996 , 21, 2763-9	3.3	733
31	Morphology of the cervical intervertebral disc: implications for McKenzie's model of the disc derangement syndrome. <i>Manual Therapy</i> , 1996 , 1, 76-81		15
30	Evaluation of the relationship between laboratory and clinical tests of transversus abdominis function. <i>Physiotherapy Research International</i> , 1996 , 1, 30-40	1.8	111
29	An Initial Analysis of Thoracic Spine Movement During Unilateral Arm Elevation. <i>Journal of Manual and Manipulative Therapy</i> , 1995 , 3, 15-20	1.6	27
28	Kinematic Analysis of the Lumbar Lateral Flexion and Lumbar Lateral Shift Movement Techniques. <i>Journal of Manual and Manipulative Therapy</i> , 1995 , 3, 104-109	1.6	7
27	Ultrasound imaging in rehabilitation. <i>Australian Journal of Physiotherapy</i> , 1995 , 41, 187-93		46
26	Magnetic resonance imaging and ultrasonography of the lumbar multifidus muscle. Comparison of two different modalities. <i>Spine</i> , 1995 , 20, 54-8	3.3	209
25	Muscle control-pain control. What exercises would you prescribe?. <i>Manual Therapy</i> , 1995 , 1, 2-10		438

24	Cervical musculoskeletal dysfunction in post-concussional headache. <i>Cephalalgia</i> , 1994 , 14, 273-9; discussion 257	6.1	101
23	Manual examination: is pain provocation a major cue for spinal dysfunction?. <i>Australian Journal of Physiotherapy</i> , 1994 , 40, 159-65		50
22	The relationship between upper trapezius muscle length and upper quadrant neural tissue extensibility. <i>Australian Journal of Physiotherapy</i> , 1994 , 40, 99-103		29
21	Effectiveness of Manipulative Physiotherapy in the Management of Cervicogenic Headache: A Single Case Study. <i>Physiotherapy</i> , 1994 , 80, 417-423	3	37
20	Evidence of lumbar multifidus muscle wasting ipsilateral to symptoms in patients with acute/subacute low back pain. <i>Spine</i> , 1994 , 19, 165-72	3.3	645
19	The influence of thoracic posture and movement on range of arm elevation. <i>Physiotherapy Theory and Practice</i> , 1993 , 9, 143-148	1.5	61
18	Towards a measurement of active muscle control for lumbar stabilisation. <i>Australian Journal of Physiotherapy</i> , 1993 , 39, 187-93		70
17	The relationship between the dynamic and static function of abdominal muscles. <i>Australian Journal of Physiotherapy</i> , 1993 , 39, 9-13		40
16	Adverse tension in the neural system. A preliminary study of tennis elbow. <i>Australian Journal of Physiotherapy</i> , 1993 , 39, 15-22		61
15	Techniques for active lumbar stabilisation for spinal protection: A pilot study. <i>Australian Journal of Physiotherapy</i> , 1992 , 38, 105-12		100
14	A modified upper limb tension test: An investigation of responses in normal subjects. <i>Australian Journal of Physiotherapy</i> , 1991 , 37, 143-52		42
13	An initial evaluation of eight abdominal exercises for their ability to provide stabilisation for the lumbar spine. <i>Australian Journal of Physiotherapy</i> , 1990 , 36, 6-11		64
12	A survey of work profiles and histories of physiotherapy graduates from the university of queensland. <i>Australian Journal of Physiotherapy</i> , 1989 , 35, 246-53		3
11	The distribution and characteristics of neck-arm pain in patients with and without a neurological deficit. <i>Australian Journal of Physiotherapy</i> , 1989 , 35, 3-8		9
10	Evaluation of a temporary prosthetic insert in the rehabilitation of elderly ischaemic below-knee amputees: a pilot study. <i>Australian Journal of Physiotherapy</i> , 1988 , 34, 133-8		3
9	The accuracy of manual diagnosis for cervical zygapophysial joint pain syndromes. <i>Medical Journal of Australia</i> , 1988 , 148, 233-6	4	285
8	Thoracic kyphosis and mobility: the effect of age. <i>Physiotherapy Practice</i> , 1987 , 3, 154-162		27
7	A Motion Profile of the Lumbar Spine in an Ageing Population Assessed by Manual Examination. <i>Physiotherapy Practice</i> , 1987 , 3, 70-81		34

6	The relationship of low back pain to postural changes during pregnancy. <i>Australian Journal of Physiotherapy</i> , 1987 , 33, 10-7	56
5	The influence of segmental level and direction of movement on age changes in lumbar motion as assessed by manual examination. <i>Physiotherapy Practice</i> , 1987 , 3, 107-116	3
4	Physiotherapy at the XII Commonwealth Games Part I: Organization and Utilization of Services. <i>Australian Journal of Physiotherapy</i> , 1984 , 30, 3-9	3
3	Physiotherapy at the XII Commonwealth Games Part II: Injuries and Management. <i>Australian Journal of Physiotherapy</i> , 1984 , 30, 10-4	3
2	The role of passive mobilization in the immediate management of the fractured neck of humerus. <i>Australian Journal of Physiotherapy</i> , 1979 , 25, 107-14	2
1	Manual Therapy164-176	4