Thierry Lefevre

List of Publications by Citations

Source: https://exaly.com/author-pdf/8471990/thierry-lefevre-publications-by-citations.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

2,513 25 49 g-index

72 2,734 4.4 5.01 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
66	Protein secondary structure and orientation in silk as revealed by Raman spectromicroscopy. <i>Biophysical Journal</i> , 2007 , 92, 2885-95	2.9	317
65	Study of protein conformation and orientation in silkworm and spider silk fibers using Raman microspectroscopy. <i>Biomacromolecules</i> , 2004 , 5, 2247-57	6.9	256
64	Molecular differences in the formation and structure of fine-stranded and particulate beta-lactoglobulin gels. <i>Biopolymers</i> , 2000 , 54, 578-86	2.2	199
63	Effects of ultra-high pressure homogenization on the properties and structure of interfacial protein layer in whey protein-stabilized emulsion. <i>Food Chemistry</i> , 2009 , 113, 191-195	8.5	114
62	Formation of intermolecular beta-sheet structures: a phenomenon relevant to protein film structure at oil-water interfaces of emulsions. <i>Journal of Colloid and Interface Science</i> , 2003 , 263, 59-67	9.3	101
61	Structural and interaction properties of Elactoglobulin as studied by FTIR spectroscopy. <i>International Journal of Food Science and Technology</i> , 1999 , 34, 419-428	3.8	98
60	Review structure of silk by raman spectromicroscopy: from the spinning glands to the fibers. <i>Biopolymers</i> , 2012 , 97, 322-36	2.2	75
59	Native spider silk as a biological optical fiber. <i>Applied Physics Letters</i> , 2013 , 102, 123702	3.4	75
58	Characterization by Raman microspectroscopy of the strain-induced conformational transition in fibroin fibers from the silkworm Samia cynthia ricini. <i>Biomacromolecules</i> , 2006 , 7, 2512-21	6.9	74
57	Evidence by infrared spectroscopy of the presence of two types of Bheets in major ampullate spider silk and silkworm silk. <i>Soft Matter</i> , 2013 , 9, 208-215	3.6	73
56	Diversity of molecular transformations involved in the formation of spider silks. <i>Journal of Molecular Biology</i> , 2011 , 405, 238-53	6.5	68
55	Conformation and orientation of proteins in various types of silk fibers produced by Nephila clavipes spiders. <i>Biomacromolecules</i> , 2009 , 10, 2945-53	6.9	67
54	Changes and roles of secondary structures of whey protein for the formation of protein membrane at soy oil/water interface under high-pressure homogenization. <i>Journal of Agricultural and Food Chemistry</i> , 2007 , 55, 10924-31	5.7	67
53	Conformational and orientational transformation of silk proteins in the major ampullate gland of Nephila clavipes spiders. <i>Biomacromolecules</i> , 2008 , 9, 2399-407	6.9	64
52	Interaction of beta-lactoglobulin with phospholipid bilayers: a molecular level elucidation as revealed by infrared spectroscopy. <i>International Journal of Biological Macromolecules</i> , 2000 , 28, 59-67	7.9	62
51	In situ conformation of spider silk proteins in the intact major ampullate gland and in solution. <i>Biomacromolecules</i> , 2007 , 8, 2342-4	6.9	57
50	Molecular description of the formation and structure of plasticized globular protein films. <i>Biomacromolecules</i> , 2005 , 6, 3209-19	6.9	54

(2013-2001)

49	Molecular structure and interaction of biopolymers as viewed by Fourier transform infrared spectroscopy: model studies on Elactoglobulin. <i>Food Hydrocolloids</i> , 2001 , 15, 365-376	10.6	52	
48	Study of protein aggregation using two-dimensional correlation infrared spectroscopy and spectral simulations. <i>Biopolymers</i> , 2004 , 73, 705-15	2.2	45	
47	Conformational rearrangement of beta-lactoglobulin upon interaction with an anionic membrane. <i>BBA - Proteins and Proteomics</i> , 2001 , 1549, 37-50		39	
46	Attenuated total reflection infrared spectroscopy: an efficient technique to quantitatively determine the orientation and conformation of proteins in single silk fibers. <i>Applied Spectroscopy</i> , 2008 , 62, 956-62	3.1	38	
45	Orientation-insensitive spectra for Raman microspectroscopy. <i>Applied Spectroscopy</i> , 2006 , 60, 841-6	3.1	36	
44	Effect of pH on the structure of the recombinant C-terminal domain of Nephila clavipes dragline silk protein. <i>Biomacromolecules</i> , 2014 , 15, 4447-54	6.9	34	
43	Spider silk as a blueprint for greener materials: a review. <i>International Materials Reviews</i> , 2016 , 61, 127-	1 56 .1	33	
42	Spider wrapping silk fibre architecture arising from its modular soluble protein precursor. <i>Scientific Reports</i> , 2015 , 5, 11502	4.9	25	
41	Mimicking and Understanding the Agglutination Effect of the Antimicrobial Peptide Thanatin Using Model Phospholipid Vesicles. <i>Biochemistry</i> , 2015 , 54, 3932-41	3.2	24	
40	A comparative study between human skin substitutes and normal human skin using Raman microspectroscopy. <i>Acta Biomaterialia</i> , 2014 , 10, 2703-11	10.8	24	
39	Study by Raman spectromicroscopy of the effect of tensile deformation on the molecular structure of Bombyx mori silk. <i>Vibrational Spectroscopy</i> , 2009 , 51, 136-141	2.1	23	
38	Unexpected Esheets and molecular orientation in flagelliform spider silk as revealed by Raman spectromicroscopy. <i>Soft Matter</i> , 2012 , 8, 6350	3.6	18	
37	Vibrational Circular Dichroism Reveals Supramolecular Chirality Inversion of Esynuclein Peptide Assemblies upon Interactions with Anionic Membranes. <i>ACS Nano</i> , 2019 , 13, 3232-3242	16.7	18	
36	Hydrodynamical properties of recombinant spider silk proteins: Effects of pH, salts and shear, and implications for the spinning process. <i>Biopolymers</i> , 2013 , 99, 582-93	2.2	17	
35	Understanding amyloid fibril formation using protein fragments: structural investigations via vibrational spectroscopy and solid-state NMR. <i>Biophysical Reviews</i> , 2018 , 10, 1133-1149	3.7	16	
34	Solvation of Lauric Acid Studied by Vibrational Spectroscopies. <i>Applied Spectroscopy</i> , 1995 , 49, 1268-127	7 4 .1	15	
33	Major Ampullate Spider Silk with Indistinguishable Spidroin Dope Conformations Leads to Different Fiber Molecular Structures. <i>International Journal of Molecular Sciences</i> , 2016 , 17,	6.3	15	
32	Structure and membrane interactions of the Eamyloid fragment 25-35 as viewed using spectroscopic approaches. <i>Physical Chemistry Chemical Physics</i> , 2013 , 15, 7228-39	3.6	14	

31	Vitamin E-phospholipid interactions in model multilayer membranes: A spectroscopic study. <i>Biospectroscopy</i> , 1998 , 2, 391-403		13
30	Mutant bovine odorant-binding protein: Temperature affects the protein stability and dynamics as revealed by infrared spectroscopy and molecular dynamics simulations. <i>Proteins: Structure, Function and Bioinformatics</i> , 2008 , 72, 769-78	4.2	13
29	Structure and pH-induced alterations of recombinant and natural spider silk proteins in solution. <i>Biopolymers</i> , 2012 , 97, 337-46	2.2	12
28	Structure and mechanical properties of spider silk films at the air-water interface. <i>Langmuir</i> , 2013 , 29, 7931-8	4	11
27	Surface properties and conformation of Nephila clavipes spider recombinant silk proteins at the air-water interface. <i>Langmuir</i> , 2009 , 25, 8170-80	4	11
26	A quantitative analysis of the supercontraction-induced molecular disorientation of major ampullate spider silk. <i>Physical Chemistry Chemical Physics</i> , 2017 , 19, 31487-31498	3.6	10
25	Besides fibrillization: putative role of the peptide fragment 71-82 on the structural and assembly behavior of Esynuclein. <i>Biochemistry</i> , 2014 , 53, 6463-72	3.2	10
24	Raman and Fourier transform infrared study of phytol effects on saturated and unsaturated lipid multibilayers. <i>Journal of Raman Spectroscopy</i> , 2003 , 34, 4-12	2.3	10
23	Crystallization of water in multilamellar vesicles. European Biophysics Journal, 2002, 31, 126-35	1.9	9
22	Binding of pediocin PA-1 with anionic lipid induces model membrane destabilization. <i>Applied and Environmental Microbiology</i> , 2003 , 69, 6777-84	4.8	9
21	Discriminating Lipid- from Protein-Calcium Binding To Understand the Interaction between Recoverin and Phosphatidylglycerol Model Membranes. <i>Biochemistry</i> , 2016 , 55, 3481-91	3.2	8
20	Structural and Mechanical Roles for the C-Terminal Nonrepetitive Domain Become Apparent in Recombinant Spider Aciniform Silk. <i>Biomacromolecules</i> , 2017 , 18, 3678-3686	6.9	8
19	Using infrared and Raman microspectroscopies to compare ex vivo involved psoriatic skin with normal human skin. <i>Journal of Biomedical Optics</i> , 2015 , 20, 067004	3.5	8
18	Importance of the Reference Spectrum on Generalized Two-Dimensional Correlation Spectroscopy: Relation between Intensity Variations and Synchronism. <i>Journal of Physical Chemistry A</i> , 2003 , 107, 6366-6372	2.8	8
17	Interaction of oil bodies proteins with phospholipid bilayers: A molecular level elucidation as revealed by infrared spectroscopy. <i>International Journal of Biological Macromolecules</i> , 2019 , 122, 873-8	887 ^{.9}	8
16	Spider silk inspired materials and sustainability: perspective. <i>Materials Technology</i> , 2016 , 1-16	2.1	7
15	The thermal stability of recoverin depends on calcium binding and its myristoyl moiety as revealed by infrared spectroscopy. <i>Biochemistry</i> , 2014 , 53, 48-56	3.2	6
14	Membrane fluidity is a driving force for recoverin myristoyl immobilization in zwitterionic lipids. Biochemical and Biophysical Research Communications, 2017, 490, 1268-1273	3.4	6

LIST OF PUBLICATIONS

13	Mapping molecular orientation in dry and wetNephila clavipesdragline spider silk. <i>Journal of Physics: Conference Series</i> , 2009 , 186, 012089	0.3	6
12	Chapter 8 Characterization of Molecular Orientation. Comprehensive Analytical Chemistry, 2008, 295-33	5 1.9	6
11	Structure of a Parkinson& Disease-Involved Esynuclein Peptide Is Modulated by Membrane Composition and Physical State. <i>Journal of Physical Chemistry B</i> , 2020 , 124, 3469-3481	3.4	5
10	Effect of mechanical deformation on the structure of regenerated Bombyx mori silk fibroin films as revealed using Raman and infrared spectroscopy. <i>Applied Spectroscopy</i> , 2015 , 69, 689-98	3.1	5
9	15. Protein- and peptide-based materials: a source of inspiration for innovation 2014 , 415-442		4
8	Solid-state nuclear magnetic resonance (NMR) spectroscopy reveals distinctive protein dynamics in closely related spider silks. <i>Canadian Journal of Chemistry</i> , 2011 , 89, 1047-1054	0.9	4
7	Thermotropic aspects of multilamellar organisation of mono-unsaturated phospholipid OPPC. <i>Chemistry and Physics of Lipids</i> , 1998 , 92, 79-89	3.7	4
6	Novel approaches to probe the binding of recoverin to membranes. <i>European Biophysics Journal</i> , 2018 , 47, 679-691	1.9	3
5	A surface spectroscopy study of a Pseudomonas fluorescens biofilm in the presence of an immobilized air bubble. <i>Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy</i> , 2019 , 222, 117163	4.4	1
4	Lipid membrane interactions of a fluorinated peptide with potential ion channel-forming ability. <i>Peptide Science</i> , 2019 , 111, e24051	3	1
3	Raman Microspectroscopy: An Ideal Technique to Study the Conformation and Orientation of Proteins in Silkworm and Spider Silk Fibers. <i>Microscopy and Microanalysis</i> , 2004 , 10, 1314-1315	0.5	
2	Troxerutine and Vitamin E Interactions in Model Membranes. a Spectroscopic Study 1995 , 407-408		
1	Transdermal diffusion, spatial distribution and physical state of a potential anticancer drug in mouse skin as studied by diffusion and spectroscopic techniques. <i>Biomedical Spectroscopy and Imaging</i> , 2018 , 7, 47-61	1.3	