

# Jun Hata

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8472997/publications.pdf>

Version: 2024-02-01

141  
papers

6,264  
citations

81900

39  
h-index

82547

72  
g-index

145  
all docs

145  
docs citations

145  
times ranked

10381  
citing authors

#	ARTICLE	IF	CITATIONS
1	Multiancestry genome-wide association study of 520,000 subjects identifies 32 loci associated with stroke and stroke subtypes. <i>Nature Genetics</i> , 2018, 50, 524-537.	21.4	1,124
2	Secular Trends in Cardiovascular Disease and Its Risk Factors in Japanese. <i>Circulation</i> , 2013, 128, 1198-1205.	1.6	250
3	Midlife and Late-Life Blood Pressure and Dementia in Japanese Elderly. <i>Hypertension</i> , 2011, 58, 22-28.	2.7	214
4	LDL Cholesterol and the Development of Stroke Subtypes and Coronary Heart Disease in a General Japanese Population. <i>Stroke</i> , 2009, 40, 382-388.	2.0	189
5	Effects of Visit-to-Visit Variability in Systolic Blood Pressure on Macrovascular and Microvascular Complications in Patients With Type 2 Diabetes Mellitus. <i>Circulation</i> , 2013, 128, 1325-1334.	1.6	189
6	High-Sensitivity C-Reactive Protein and Coronary Heart Disease in a General Population of Japanese. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2008, 28, 1385-1391.	2.4	180
7	Trends in dementia prevalence, incidence, and survival rate in a Japanese community. <i>Neurology</i> , 2017, 88, 1925-1932.	1.1	154
8	Epidemiology of Stroke and Coronary Artery Disease in Asia. <i>Circulation Journal</i> , 2013, 77, 1923-1932.	1.6	151
9	Prestroke Glycemic Control Is Associated With the Functional Outcome in Acute Ischemic Stroke. <i>Stroke</i> , 2011, 42, 2788-2794.	2.0	134
10	Secular Trends in the Incidence of and Risk Factors for Ischemic Stroke and Its Subtypes in Japanese Population. <i>Circulation</i> , 2008, 118, 2672-2678.	1.6	119
11	Day-to-Day Blood Pressure Variability and Risk of Dementia in a General Japanese Elderly Population. <i>Circulation</i> , 2017, 136, 516-525.	1.6	113
12	Trends in the prevalence of chronic kidney disease and its risk factors in a general Japanese population: The Hisayama Study. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 2557-2564.	0.7	111
13	Functional SNP in an Sp1-binding site of AGR1 gene is associated with susceptibility to brain infarction. <i>Human Molecular Genetics</i> , 2007, 16, 630-639.	2.9	105
14	Tooth Loss and Risk of Dementia in the Community: the Hisayama Study. <i>Journal of the American Geriatrics Society</i> , 2017, 65, e95-e100.	2.6	103
15	Association between ratio of serum eicosapentaenoic acid to arachidonic acid and risk of cardiovascular disease: The Hisayama Study. <i>Atherosclerosis</i> , 2013, 231, 261-267.	0.8	101
16	Intracerebral hemorrhage location and outcome among INTERACT2 participants. <i>Neurology</i> , 2017, 88, 1408-1414.	1.1	101
17	High Blood Pressure After Acute Ischemic Stroke Is Associated With Poor Clinical Outcomes. <i>Hypertension</i> , 2014, 63, 54-60.	2.7	99
18	Self-Reported Dietary Intake of Potassium, Calcium, and Magnesium and Risk of Dementia in the Japanese: The Hisayama Study. <i>Journal of the American Geriatrics Society</i> , 2012, 60, 1515-1520.	2.6	93

#	ARTICLE	IF	CITATIONS
19	Milk and Dairy Consumption and Risk of Dementia in an Elderly Japanese Population: The Hisayama Study. <i>Journal of the American Geriatrics Society</i> , 2014, 62, 1224-1230.	2.6	92
20	Tongue Microbiota and Oral Health Status in Community-Dwelling Elderly Adults. <i>MSphere</i> , 2018, 3, .	2.9	73
21	Insulin resistance and clinical outcomes after acute ischemic stroke. <i>Neurology</i> , 2018, 90, e1470-e1477.	1.1	72
22	Prevalence and Causes of Functional Disability in an Elderly General Population of Japanese: The Hisayama Study. <i>Journal of Epidemiology</i> , 2012, 22, 222-229.	2.4	71
23	Association Between Diabetes and Hippocampal Atrophy in Elderly Japanese: The Hisayama Study. <i>Diabetes Care</i> , 2016, 39, 1543-1549.	8.6	71
24	Day-by-Day Blood Pressure Variability and Functional Outcome After Acute Ischemic Stroke. <i>Stroke</i> , 2015, 46, 1832-1839.	2.0	67
25	The long-term association between physical activity and risk of dementia in the community: the Hisayama Study. <i>European Journal of Epidemiology</i> , 2016, 31, 267-274.	5.7	67
26	Association Between Daily Sleep Duration and Risk of Dementia and Mortality in a Japanese Community. <i>Journal of the American Geriatrics Society</i> , 2018, 66, 1911-1918.	2.6	64
27	Moyamoya Disease Susceptibility Variant <i>RNF213</i> p.R4810K Increases the Risk of Ischemic Stroke Attributable to Large-Artery Atherosclerosis. <i>Circulation</i> , 2019, 139, 295-298.	1.6	64
28	Plasma C-Reactive Protein and Clinical Outcomes after Acute Ischemic Stroke: A Prospective Observational Study. <i>PLoS ONE</i> , 2016, 11, e0156790.	2.5	59
29	Midlife and Late-Life Smoking and Risk of Dementia in the Community: The Hisayama Study. <i>Journal of the American Geriatrics Society</i> , 2015, 63, 2332-2339.	2.6	56
30	Sex Differences in Short-Term Outcomes After Acute Ischemic Stroke. <i>Stroke</i> , 2015, 46, 471-476.	2.0	55
31	Hematocrit and the risk of cardiovascular disease in a Japanese community: The Hisayama Study. <i>Atherosclerosis</i> , 2015, 242, 199-204.	0.8	54
32	Small Dense Low-Density Lipoprotein Cholesterol and the Risk of Coronary Heart Disease in a Japanese Community. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 669-682.	2.0	52
33	Development and validation of a cardiovascular risk prediction model for Japanese: the Hisayama study. <i>Hypertension Research</i> , 2009, 32, 1119-1122.	2.7	51
34	Midlife and late-life handgrip strength and risk of cause-specific death in a general Japanese population: the Hisayama Study. <i>Journal of Epidemiology and Community Health</i> , 2014, 68, 663-668.	3.7	48
35	Study design and baseline characteristics of a population-based prospective cohort study of dementia in Japan: the Japan Prospective Studies Collaboration for Aging and Dementia (JPSC-AD). <i>Environmental Health and Preventive Medicine</i> , 2020, 25, 64.	3.4	47
36	Serum Soluble Triggering Receptor Expressed on Myeloid Cells 2 as a Biomarker for Incident Dementia: The Hisayama Study. <i>Annals of Neurology</i> , 2019, 85, 47-58.	5.3	45

#	ARTICLE	IF	CITATIONS
37	N-Terminal Pro-Brain Natriuretic Peptide and Risk of Cardiovascular Events in a Japanese Community. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011, 31, 2997-3003.	2.4	44
38	Serum Uric Acid as a Risk Factor for Chronic Kidney Disease in a Japanese Community—“ The Hisayama Study ”. <i>Circulation Journal</i> , 2016, 80, 1857-1862.	1.6	44
39	Association Between Onset-to-Door Time and Clinical Outcomes After Ischemic Stroke. <i>Stroke</i> , 2017, 48, 3049-3056.	2.0	44
40	Significance of plasma adiponectin for diagnosis, neurological severity and functional outcome in ischemic stroke “ Research for Biomarkers in Ischemic Stroke (REBIOS). <i>Metabolism: Clinical and Experimental</i> , 2014, 63, 1093-1103.	3.4	42
41	Left Atrial Size and Long-Term Risk of Recurrent Stroke After Acute Ischemic Stroke in Patients With Nonvalvular Atrial Fibrillation. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	40
42	Albuminuria Increases the Risks for Both Alzheimer Disease and Vascular Dementia in Community-Dwelling Japanese Elderly: The Hisayama Study. <i>Journal of the American Heart Association</i> , 2018, 7, .	3.7	40
43	Combined Effects of Smoking and Hypercholesterolemia on the Risk of Stroke and Coronary Heart Disease in Japanese: The Hisayama Study. <i>Cerebrovascular Diseases</i> , 2011, 31, 477-484.	1.7	38
44	Trends in the Prevalence of Myopia and Myopic Maculopathy in a Japanese Population: The Hisayama Study. , 2019, 60, 2781.		38
45	Non-high-density lipoprotein cholesterol and the development of coronary heart disease and stroke subtypes in a general Japanese population: The Hisayama Study. <i>Atherosclerosis</i> , 2014, 233, 343-348.	0.8	37
46	Gastrointestinal Bleeding in Acute Ischemic Stroke: Recent Trends from the Fukuoka Stroke Registry. <i>Cerebrovascular Diseases Extra</i> , 2014, 4, 156-164.	1.5	37
47	Reduced Estimated GFR and Cardiac Remodeling: A Population-Based Autopsy Study. <i>American Journal of Kidney Diseases</i> , 2019, 74, 373-381.	1.9	34
48	Association of hemoglobin A1c and glycated albumin with carotid atherosclerosis in community-dwelling Japanese subjects: the Hisayama Study. <i>Cardiovascular Diabetology</i> , 2015, 14, 84.	6.8	33
49	Prevalence and Mortality of Sarcopenia in a Community-dwelling Older Japanese Population: The Hisayama Study. <i>Journal of Epidemiology</i> , 2021, 31, 320-327.	2.4	33
50	Serum Angiopoietin-Like Protein 2 Is a Novel Risk Factor for Cardiovascular Disease in the Community. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016, 36, 1686-1691.	2.4	31
51	Alternative Measures of Hyperglycemia and Risk of Alzheimer’s Disease in the Community: The Hisayama Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, 3002-3010.	3.6	31
52	Association between Axial Length and Myopic Maculopathy. <i>Ophthalmology Retina</i> , 2019, 3, 867-873.	2.4	30
53	Patterns and Levels of Sedentary Behavior and Physical Activity in a General Japanese Population: The Hisayama Study. <i>Journal of Epidemiology</i> , 2018, 28, 260-265.	2.4	29
54	Plasma S100A12 is associated with functional outcome after ischemic stroke: Research for Biomarkers in Ischemic Stroke. <i>Journal of the Neurological Sciences</i> , 2014, 340, 75-79.	0.6	28

#	ARTICLE	IF	CITATIONS
55	Decline in Handgrip Strength From Midlife to Late-Life is Associated With Dementia in a Japanese Community: The Hisayama Study. <i>Journal of Epidemiology</i> , 2020, 30, 15-23.	2.4	26
56	Insulin Resistance and the Development of Cardiovascular Disease in a Japanese Community: the Hisayama Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012, 19, 977-985.	2.0	26
57	Genome-Wide Polygenic Score and the Risk of Ischemic Stroke in a Prospective Cohort. <i>Stroke</i> , 2020, 51, 759-765.	2.0	25
58	Dietary fiber intake and risk of type 2 diabetes in a general Japanese population: The Hisayama Study. <i>Journal of Diabetes Investigation</i> , 2021, 12, 527-536.	2.4	24
59	Development and Validation of a Risk Prediction Model for Atherosclerotic Cardiovascular Disease in Japanese Adults: The Hisayama Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 345-361.	2.0	23
60	Disrupted tongue microbiota and detection of nonindigenous bacteria on the day of allogeneic hematopoietic stem cell transplantation. <i>PLoS Pathogens</i> , 2020, 16, e1008348.	4.7	22
61	Adjustment of Cell-Type Composition Minimizes Systematic Bias in Blood DNA Methylation Profiles Derived by DNA Collection Protocols. <i>PLoS ONE</i> , 2016, 11, e0147519.	2.5	21
62	Associations with health-related quality of life after intracerebral haemorrhage: pooled analysis of INTERACT studies. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2017, 88, 70-75.	1.9	21
63	Dietary Protein Intake and Stroke Risk in a General Japanese Population. <i>Stroke</i> , 2017, 48, 1478-1486.	2.0	21
64	Development and validation of a risk assessment tool for gastric cancer in a general Japanese population. <i>Gastric Cancer</i> , 2018, 21, 383-390.	5.3	21
65	The effect of metabolic syndrome defined by various criteria on the development of ischemic stroke subtypes in a general Japanese population. <i>Atherosclerosis</i> , 2010, 210, 249-255.	0.8	20
66	Prevalence of and risk factors for cerebral microbleeds in a general Japanese elderly community. <i>Neurology: Clinical Practice</i> , 2018, 8, 223-231.	1.6	20
67	Development and validation of modified risk prediction models for cardiovascular disease and its subtypes: The Hisayama Study. <i>Atherosclerosis</i> , 2018, 279, 38-44.	0.8	19
68	Association between the ratio of serum arachidonic acid to eicosapentaenoic acid and the presence of depressive symptoms in a general Japanese population: the Hisayama Study. <i>Journal of Affective Disorders</i> , 2018, 237, 73-79.	4.1	19
69	Intravenous Thrombolysis with Recombinant Tissue Plasminogen Activator for Ischemic Stroke Patients over 80 Years Old: The Fukuoka Stroke Registry. <i>PLoS ONE</i> , 2014, 9, e110444.	2.5	18
70	The ratio of serum eicosapentaenoic acid to arachidonic acid and risk of cancer death in a Japanese community: The Hisayama Study. <i>Journal of Epidemiology</i> , 2017, 27, 578-583.	2.4	18
71	Serum Non-High-Density Lipoprotein Cholesterol and Risk of Cardiovascular Disease in Community Dwellers with Chronic Kidney Disease: the Hisayama Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017, 24, 706-715.	2.0	18
72	Prevalence and Risk Factors for Polypoidal Choroidal Vasculopathy in a General Japanese Population: The Hisayama Study. <i>Seminars in Ophthalmology</i> , 2018, 33, 813-819.	1.6	18

#	ARTICLE	IF	CITATIONS
73	Association Between Serum Î²-Alanine and Risk of Dementia. American Journal of Epidemiology, 2019, 188, 1637-1645.	3.4	18
74	Serum homocysteine and risk of dementia in Japan. Journal of Neurology, Neurosurgery and Psychiatry, 2020, 91, 540-546.	1.9	18
75	Trends in the prevalence of type 2 diabetes and prediabetes in a Japanese community, 1988â€“2012: the Hisayama Study. Diabetology International, 2019, 10, 198-205.	1.4	17
76	White-coat and masked hypertension are associated with albuminuria in a general population: the Hisayama Study. Hypertension Research, 2017, 40, 937-943.	2.7	16
77	Influence of the Accumulation of Unhealthy Eating Habits on Obesity in a General Japanese Population: The Hisayama Study. Nutrients, 2020, 12, 3160.	4.1	16
78	Association Between Serum Vitamin D and All-Cause and Cause-Specific Death in a General Japanese Populationâ€”The Hisayama Study â€”. Circulation Journal, 2017, 81, 1315-1321.	1.6	15
79	Association of Embolic Sources With Cause-Specific Functional Outcomes Among Adults With Cryptogenic Stroke. JAMA Network Open, 2018, 1, e182953.	5.9	15
80	Association between serum glycated albumin and risk of cardiovascular disease in a Japanese community: The Hisayama Study. Atherosclerosis, 2020, 311, 52-59.	0.8	15
81	Association of Albuminuria With White Matter Hyperintensities Volume on Brain Magnetic Resonance Imaging in Elderly Japaneseâ€”The Hisayama Study â€”. Circulation Journal, 2020, 84, 935-942.	1.6	15
82	Periodontal status and lung function decline in the community: the Hisayama study. Scientific Reports, 2018, 8, 13354.	3.3	14
83	Insulin Resistance Is a Risk Factor for Increased Intraocular Pressure: The Hisayama Study. , 2015, 56, 7983.		13
84	Emotional Loneliness Is Associated With a Risk of Dementia in a General Japanese Older Population: The Hisayama Study. Journals of Gerontology - Series B Psychological Sciences and Social Sciences, 2020, 76, 1756-1766.	3.9	13
85	Five-Year Incidence of Myopic Maculopathy in a General Japanese Population. JAMA Ophthalmology, 2020, 138, 887.	2.5	13
86	Serum uric acid levels and cardiovascular mortality in a general Japanese population: the Hisayama Study. Hypertension Research, 2020, 43, 560-568.	2.7	13
87	The Association of Small Dense Low-Density Lipoprotein Cholesterol and Coronary Heart Disease in Subjects at High Cardiovascular Risk. Journal of Atherosclerosis and Thrombosis, 2021, 28, 79-89.	2.0	13
88	Long-term association of vegetable and fruit intake with risk of dementia in Japanese older adults: the Hisayama study. BMC Geriatrics, 2022, 22, 257.	2.7	13
89	Association Between Genetic Risk and Development of Type 2 Diabetes in a General Japanese Population: The Hisayama Study. Journal of Clinical Endocrinology and Metabolism, 2019, 104, 3213-3222.	3.6	12
90	Risk prediction models for mortality in patients with cardiovascular disease: The BioBank Japan project. Journal of Epidemiology, 2017, 27, S71-S76.	2.4	11

#	ARTICLE	IF	CITATIONS
91	Dairy consumption and risk of functional disability in an elderly Japanese population: the Hisayama Study. <i>American Journal of Clinical Nutrition</i> , 2019, 109, 1664-1671.	4.7	11
92	Serum elaidic acid concentration and risk of dementia. <i>Neurology</i> , 2019, 93, e2053-e2064.	1.1	11
93	Association of glucose tolerance status with pancreatic $\beta$ -cell mass in community-based autopsy samples of Japanese individuals: The Hisayama Study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1197-1206.	2.4	11
94	Multiple-region grey matter atrophy as a predictor for the development of dementia in a community: the Hisayama Study. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2022, 93, 263-271.	1.9	11
95	A potential novel pathological implication of serum soluble triggering receptor expressed on myeloid cell 2 in insulin resistance in a general Japanese population: The Hisayama study. <i>Diabetes Research and Clinical Practice</i> , 2018, 146, 225-232.	2.8	10
96	Lifetime cumulative incidence of dementia in a community-dwelling elderly population in Japan. <i>Neurology</i> , 2020, 95, e508-e518.	1.1	10
97	Comparison of the contributions of impaired beta cell function and insulin resistance to the development of type 2 diabetes in a Japanese community: the Hisayama Study. <i>Diabetologia</i> , 2021, 64, 1775-1784.	6.3	10
98	Serum Ethylamine Levels as an Indicator of Theanine Consumption and the Risk of Type 2 Diabetes in a General Japanese Population: The Hisayama Study. <i>Diabetes Care</i> , 2019, 42, 1234-1240.	8.6	9
99	Glucose Tolerance Levels and Circumpapillary Retinal Nerve Fiber Layer Thickness in a General Japanese Population: The Hisayama Study. <i>American Journal of Ophthalmology</i> , 2019, 205, 140-146.	3.3	9
100	Poor glycemic control and posterior circulation ischemic stroke. <i>Neurology: Clinical Practice</i> , 2019, 9, 129-139.	1.6	9
101	Elevated serum glycosylated albumin and glycosylated albumin-hemoglobin A <sub>1c</sub> ratio were associated with hippocampal atrophy in a general elderly population of Japanese: The Hisayama Study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 971-979.	2.4	9
102	Pathologic Diabetic Nephropathy in Autopsied Diabetic Cases With Normoalbuminuria From a Japanese Community-Based Study. <i>Kidney International Reports</i> , 2021, 6, 3035-3044.	0.8	9
103	Objectively measured sedentary time and diabetes mellitus in a general Japanese population: The Hisayama Study. <i>Journal of Diabetes Investigation</i> , 2019, 10, 809-816.	2.4	8
104	$\beta$ -Cell Function and Clinical Outcome in Nondiabetic Patients With Acute Ischemic Stroke. <i>Stroke</i> , 2021, 52, 2621-2628.	2.0	8
105	Association between chronic low back pain and regional brain atrophy in a Japanese older population: the Hisayama Study. <i>Pain</i> , 2022, 163, 2185-2193.	4.2	8
106	Morning and Evening Blood Pressures Are Associated With Intima-Media Thickness in a General Population: The Hisayama Study. <i>Circulation Journal</i> , 2017, 81, 1647-1653.	1.6	7
107	Long-term regular exercise and intraocular pressure: the Hisayama Study. <i>Graefe's Archive for Clinical and Experimental Ophthalmology</i> , 2019, 257, 2461-2469.	1.9	7
108	Risk Prediction Model for Incident Atrial Fibrillation in a General Japanese Population: The Hisayama Study. <i>Circulation Journal</i> , 2021, 85, 1373-1382.	1.6	7

#	ARTICLE	IF	CITATIONS
109	10-year trend of tooth loss and associated factors in a Japanese population-based longitudinal study. <i>BMJ Open</i> , 2021, 11, e048114.	1.9	7
110	Diabetes Mellitus, Elevated Hemoglobin A1c, and Glycated Albumin Are Associated with the Presence of All-Cause Dementia and Alzheimer's Disease: The JPSC-AD Study. <i>Journal of Alzheimer's Disease</i> , 2022, 85, 235-247.	2.6	7
111	Association Between Diabetes and Gray Matter Atrophy Patterns in a General Older Japanese Population: The Hisayama Study. <i>Diabetes Care</i> , 2022, 45, 1364-1371.	8.6	7
112	Changes in Body Weight and Concurrent Changes in Cardiovascular Risk Profiles in Community Residents in Japan: the Hisayama Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, . .	2.0	6
113	Current status of the certification of long-term care insurance among individuals with dementia in a Japanese community: The Hisayama Study. <i>Psychiatry and Clinical Neurosciences</i> , 2021, 75, 182-184.	1.8	6
114	Midlife and late-life diabetes and sarcopenia in a general older Japanese population: The Hisayama Study. <i>Journal of Diabetes Investigation</i> , 2021, 12, 1899-1907.	2.4	6
115	Day-by-Day Blood Pressure Variability in the Subacute Stage of Ischemic Stroke and Long-Term Recurrence. <i>Stroke</i> , 2022, 53, 70-78.	2.0	6
116	Association of Airflow Limitation With Carotid Atherosclerosis in a Japanese Community The Hisayama Study. <i>Circulation Journal</i> , 2017, 81, 1846-1853.	1.6	6
117	Yogurt product intake and reduction of tooth loss risk in a Japanese community. <i>Journal of Clinical Periodontology</i> , 2022, 49, 345-352.	4.9	6
118	PKC $\delta$ deficiency improves lipid metabolism and atherosclerosis in apolipoprotein E-deficient mice. <i>Genes To Cells</i> , 2016, 21, 1030-1048.	1.2	5
119	Trends in the prevalence of airflow limitation in a general Japanese population: two serial cross-sectional surveys from the Hisayama Study. <i>BMJ Open</i> , 2019, 9, e023673.	1.9	5
120	30-minute postload plasma glucose levels during an oral glucose tolerance test predict the risk of future type 2 diabetes: the Hisayama Study. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001156.	2.8	5
121	Serum N-terminal pro-B-type natriuretic peptide as a predictor for future development of atrial fibrillation in a general population: the Hisayama Study. <i>International Journal of Cardiology</i> , 2020, 320, 90-96.	1.7	5
122	High Serum Folate Concentrations Are Associated with Decreased Risk of Mortality among Japanese Adults. <i>Journal of Nutrition</i> , 2021, 151, 657-665.	2.9	5
123	Serum High-Sensitivity C-Reactive Protein Levels and the Development of Atrial Fibrillation in a General Japanese Population The Hisayama Study. <i>Circulation Journal</i> , 2021, 85, 1365-1372.	1.6	5
124	Higher-resolution quantification of white matter hypointensities by large-scale transfer learning from 2D images on the JPSC-AD cohort. <i>Human Brain Mapping</i> , 2022, 43, 3998-4012.	3.6	5
125	Ratios of serum eicosapentaenoic acid to arachidonic acid and docosahexaenoic acid to arachidonic acid were inversely associated with serum resistin levels: The Hisayama Study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 482-489.	2.4	4
126	Recent status of self-measured home blood pressure in the Japanese general population: a modern database on self-measured home blood pressure (MDAS). <i>Hypertension Research</i> , 2020, 43, 1403-1412.	2.7	4



#	ARTICLE	IF	CITATIONS
127	N-Terminal Pro-B-Type Natriuretic Peptide and Incident CKD. <i>Kidney International Reports</i> , 2021, 6, 976-985.	0.8	4
128	Urinary N-terminal pro-B-type natriuretic peptide as a biomarker for cardiovascular events in a general Japanese population: the Hisayama Study. <i>Environmental Health and Preventive Medicine</i> , 2021, 26, 47.	3.4	4
129	Baseline periodontal status and modifiable risk factors are associated with tooth loss over a 10-year period: Estimates of population attributable risk in a Japanese community. <i>Journal of Periodontology</i> , 2022, 93, 526-536.	3.4	4
130	Long-Term Trends in The 5-Year Risk of Recurrent Stroke over A Half Century in A Japanese Community: The Hisayama Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 1759-1773.	2.0	4
131	Impact of the 1425G/A Polymorphism of PRKCH on the Recurrence of Ischemic Stroke: Fukuoka Stroke Registry. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2014, 23, 1356-1361.	1.6	3
132	Secular trends in the incidence, risk factors, and prognosis of transient ischemic attack in Japan: The Hisayama Study. <i>Atherosclerosis</i> , 2018, 273, 84-90.	0.8	3
133	Anticoagulation and Risk of Stroke Recurrence in Patients with Embolic Stroke of Undetermined Source Having No Potential Source of Embolism. <i>Cerebrovascular Diseases</i> , 2020, 49, 601-608.	1.7	3
134	Development of a risk prediction model for incident hypertension in Japanese individuals: the Hisayama Study. <i>Hypertension Research</i> , 2021, 44, 1221-1229.	2.7	2
135	Biomarkers for stroke: the Hisayama Study. <i>Nosotchu</i> , 2015, 37, 352-357.	0.1	2
136	Serum Uric Acid Levels and Nephrosclerosis in a Population-Based Autopsy Study: The Hisayama Study. <i>American Journal of Nephrology</i> , 2022, 53, 69-77.	3.1	2
137	Serum NT-proBNP levels and histopathological myocardial fibrosis in autopsied cases from a Japanese community: The Hisayama Study. <i>Journal of Cardiology</i> , 2021, 78, 237-243.	1.9	1
138	Association of daily sleep duration with the incident dementia by serum soluble <sc>TREM2</sc> in a community. <i>Journal of the American Geriatrics Society</i> , 2022, 70, 1147-1156.	2.6	1
139	A Comparative Study of Site-Specific Distribution of Aging-Related Tau Astroglipathy and Its Risk Factors Between Alzheimer Disease and Cognitive Healthy Brains: The Hisayama Study. <i>Journal of Neuropathology and Experimental Neurology</i> , 2022, 81, 106-116.	1.7	1
140	Epidemiology of glucose intolerance, dyslipidemia, and stroke: the Hisayama Study. <i>Nosotchu</i> , 2016, 38, 442-448.	0.1	0
141	Airflow limitation and tongue microbiota in community-dwelling elderly individuals. <i>ERJ Open Research</i> , 2021, 7, 00616-2020.	2.6	0