

Eline H Van Den Berg

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8475355/publications.pdf>

Version: 2024-02-01

18
papers

433
citations

758635

12
h-index

940134

16
g-index

18
all docs

18
docs citations

18
times ranked

770
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence and determinants of non-alcoholic fatty liver disease in lifelines: A large Dutch population cohort. <i>PLoS ONE</i> , 2017, 12, e0171502.	1.1	74
2	Higher free triiodothyronine is associated with non-alcoholic fatty liver disease in euthyroid subjects: the Lifelines Cohort Study. <i>Metabolism: Clinical and Experimental</i> , 2017, 67, 62-71.	1.5	71
3	Non-Alcoholic Fatty Liver Disease and Risk of Incident Type 2 Diabetes: Role of Circulating Branched-Chain Amino Acids. <i>Nutrients</i> , 2019, 11, 705.	1.7	67
4	High prevalence of apolipoprotein B dyslipoproteinemias in non-alcoholic fatty liver disease: The lifelines cohort study. <i>Metabolism: Clinical and Experimental</i> , 2017, 72, 37-46.	1.5	36
5	Liver transplantation for NASH cirrhosis is not performed at the expense of major post-operative morbidity. <i>Digestive and Liver Disease</i> , 2018, 50, 68-75.	0.4	28
6	Oxidative stress is associated with suspected non-alcoholic fatty liver disease and all-cause mortality in the general population. <i>Liver International</i> , 2020, 40, 2148-2159.	1.9	28
7	Cholesterol efflux capacity is impaired in subjects with an elevated Fatty Liver Index, a proxy of non-alcoholic fatty liver disease. <i>Atherosclerosis</i> , 2018, 277, 21-27.	0.4	20
8	Nonalcoholic fatty liver disease, circulating ketone bodies and all-cause mortality in a general population-based cohort. <i>European Journal of Clinical Investigation</i> , 2021, 51, e13627.	1.7	20
9	Prescription of statins in suspected non-alcoholic fatty liver disease and high cardiovascular risk, a population-based study. <i>Liver International</i> , 2019, 39, 1343-1354.	1.9	18
10	Plasma lecithin:cholesterol acyltransferase and phospholipid transfer protein activity independently associate with nonalcoholic fatty liver disease. <i>European Journal of Clinical Investigation</i> , 2018, 48, e12988.	1.7	17
11	Serum paraoxonase 1 activity is paradoxically maintained in nonalcoholic fatty liver disease despite low HDL cholesterol. <i>Journal of Lipid Research</i> , 2019, 60, 168-175.	2.0	17
12	Higher Sodium Intake Assessed by 24 Hour Urinary Sodium Excretion Is Associated with Non-Alcoholic Fatty Liver Disease: The PREVEND Cohort Study. <i>Journal of Clinical Medicine</i> , 2019, 8, 2157.	1.0	16
13	Plasma ApoE elevations are associated with NAFLD: The PREVEND Study. <i>PLoS ONE</i> , 2019, 14, e0220659.	1.1	11
14	Branched chain amino acids are associated with metabolic complications in liver transplant recipients. <i>Clinical Biochemistry</i> , 2022, 102, 26-33.	0.8	5
15	Profoundly Disturbed Lipoproteins in Cirrhotic Patients: Role of Lipoprotein-Z, a Hepatotoxic LDL-like Lipoprotein. <i>Journal of Clinical Medicine</i> , 2022, 11, 1223.	1.0	3
16	High plasma levels of betaine, a trimethylamine N-oxide-related metabolite, are associated with the severity of cirrhosis. <i>Liver International</i> , 2022, , .	1.9	2
17	Author response to Letter to the Editor: "Statins and non-alcoholic fatty liver disease". <i>Liver International</i> , 2019, 39, 1788-1789.	1.9	0
18	Lipoprotein Z, an abnormal LDL-like lipoprotein, independently predicts mortality in cirrhosis. <i>European Journal of Internal Medicine</i> , 2022, , .	1.0	0