

Eline H Van Den Berg

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8475355/publications.pdf>

Version: 2024-02-01

18
papers

433
citations

759233

12
h-index

940533

16
g-index

18
all docs

18
docs citations

18
times ranked

770
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence and determinants of non-alcoholic fatty liver disease in lifelines: A large Dutch population cohort. PLoS ONE, 2017, 12, e0171502.	2.5	74
2	Higher free triiodothyronine is associated with non-alcoholic fatty liver disease in euthyroid subjects: the Lifelines Cohort Study. Metabolism: Clinical and Experimental, 2017, 67, 62-71.	3.4	71
3	Non-Alcoholic Fatty Liver Disease and Risk of Incident Type 2 Diabetes: Role of Circulating Branched-Chain Amino Acids. Nutrients, 2019, 11, 705.	4.1	67
4	High prevalence of apolipoprotein B dyslipoproteinemias in non-alcoholic fatty liver disease: The lifelines cohort study. Metabolism: Clinical and Experimental, 2017, 72, 37-46.	3.4	36
5	Liver transplantation for NASH cirrhosis is not performed at the expense of major post-operative morbidity. Digestive and Liver Disease, 2018, 50, 68-75.	0.9	28
6	Oxidative stress is associated with suspected non-alcoholic fatty liver disease and all-cause mortality in the general population. Liver International, 2020, 40, 2148-2159.	3.9	28
7	Cholesterol efflux capacity is impaired in subjects with an elevated Fatty Liver Index, a proxy of non-alcoholic fatty liver disease. Atherosclerosis, 2018, 277, 21-27.	0.8	20
8	Nonalcoholic fatty liver disease, circulating ketone bodies and all-cause mortality in a general population-based cohort. European Journal of Clinical Investigation, 2021, 51, e13627.	3.4	20
9	Prescription of statins in suspected non-alcoholic fatty liver disease and high cardiovascular risk, a population-based study. Liver International, 2019, 39, 1343-1354.	3.9	18
10	Plasma lecithin:cholesterol acyltransferase and phospholipid transfer protein activity independently associate with nonalcoholic fatty liver disease. European Journal of Clinical Investigation, 2018, 48, e12988.	3.4	17
11	Serum paraoxonase 1 activity is paradoxically maintained in nonalcoholic fatty liver disease despite low HDL cholesterol. Journal of Lipid Research, 2019, 60, 168-175.	4.2	17
12	Higher Sodium Intake Assessed by 24 Hour Urinary Sodium Excretion Is Associated with Non-Alcoholic Fatty Liver Disease: The PREVEND Cohort Study. Journal of Clinical Medicine, 2019, 8, 2157.	2.4	16
13	Plasma ApoE elevations are associated with NAFLD: The PREVEND Study. PLoS ONE, 2019, 14, e0220659.	2.5	11
14	Branched chain amino acids are associated with metabolic complications in liver transplant recipients. Clinical Biochemistry, 2022, 102, 26-33.	1.9	5
15	Profoundly Disturbed Lipoproteins in Cirrhotic Patients: Role of Lipoprotein-Z, a Hepatotoxic LDL-like Lipoprotein. Journal of Clinical Medicine, 2022, 11, 1223.	2.4	3
16	High plasma levels of betaine, a trimethylamine N -oxide-related metabolite, are associated with the severity of cirrhosis. Liver International, 2022, , .	3.9	2
17	Author response to Letter to the Editor: "Statins and non-alcoholic fatty liver disease". Liver International, 2019, 39, 1788-1789.	3.9	0
18	Lipoprotein Z, an abnormal LDL-like lipoprotein, independently predicts mortality in cirrhosis. European Journal of Internal Medicine, 2022, , .	2.2	0