

Daisuke Koya

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8479150/daisuke-koya-publications-by-citations.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

155
papers

9,808
citations

42
h-index

97
g-index

173
ext. papers

11,656
ext. citations

4.9
avg, IF

6.15
L-index

#	Paper	IF	Citations
155	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016 , 12, 1-222	10.2	3838
154	Calorie restriction enhances cell adaptation to hypoxia through Sirt1-dependent mitochondrial autophagy in mouse aged kidney. <i>Journal of Clinical Investigation</i> , 2010 , 120, 1043-55	15.9	451
153	Amelioration of accelerated diabetic mesangial expansion by treatment with a PKC beta inhibitor in diabetic db/db mice, a rodent model for type 2 diabetes. <i>FASEB Journal</i> , 2000 , 14, 439-47	0.9	374
152	Resveratrol improves oxidative stress and protects against diabetic nephropathy through normalization of Mn-SOD dysfunction in AMPK/SIRT1-independent pathway. <i>Diabetes</i> , 2011 , 60, 634-43	0.9	254
151	Linagliptin-mediated DPP-4 inhibition ameliorates kidney fibrosis in streptozotocin-induced diabetic mice by inhibiting endothelial-to-mesenchymal transition in a therapeutic regimen. <i>Diabetes</i> , 2014 , 63, 2120-31	0.9	245
150	Effects of antioxidants in diabetes-induced oxidative stress in the glomeruli of diabetic rats. <i>Journal of the American Society of Nephrology: JASN</i> , 2003 , 14, S250-3	12.7	212
149	Impaired Podocyte Autophagy Exacerbates Proteinuria in Diabetic Nephropathy. <i>Diabetes</i> , 2016 , 65, 755-67	0.9	174
148	Dietary restriction ameliorates diabetic nephropathy through anti-inflammatory effects and regulation of the autophagy via restoration of Sirt1 in diabetic Wistar fatty (fa/fa) rats: a model of type 2 diabetes. <i>Experimental Diabetes Research</i> , 2011 , 2011, 908185		159
147	Obesity-mediated autophagy insufficiency exacerbates proteinuria-induced tubulointerstitial lesions. <i>Journal of the American Society of Nephrology: JASN</i> , 2013 , 24, 1769-81	12.7	156
146	The protective role of Sirt1 in vascular tissue: its relationship to vascular aging and atherosclerosis. <i>Aging</i> , 2016 , 8, 2290-2307	5.6	147
145	Inhibition of mTOR signaling with rapamycin attenuates renal hypertrophy in the early diabetic mice. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 340, 296-301	3.4	130
144	Rodent models of diabetic nephropathy: their utility and limitations. <i>International Journal of Nephrology and Renovascular Disease</i> , 2016 , 9, 279-290	2.5	129
143	Nutrient sensing, autophagy, and diabetic nephropathy. <i>Diabetes</i> , 2012 , 61, 23-9	0.9	125
142	Role of the endothelial-to-mesenchymal transition in renal fibrosis of chronic kidney disease. <i>Clinical and Experimental Nephrology</i> , 2013 , 17, 488-97	2.5	120
141	Interactions of DPP-4 and integrin β 1 influences endothelial-to-mesenchymal transition. <i>Kidney International</i> , 2015 , 88, 479-89	9.9	104
140	Renal protective effects of resveratrol. <i>Oxidative Medicine and Cellular Longevity</i> , 2013 , 2013, 568093	6.7	97
139	Sirtuins and Type 2 Diabetes: Role in Inflammation, Oxidative Stress, and Mitochondrial Function. <i>Frontiers in Endocrinology</i> , 2019 , 10, 187	5.7	92

138	MicroRNAs in kidney fibrosis and diabetic nephropathy: roles on EMT and EndMT. <i>BioMed Research International</i> , 2013 , 2013, 125469	3	88
137	Fatty acids are novel nutrient factors to regulate mTORC1 lysosomal localization and apoptosis in podocytes. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2014 , 1842, 1097-108	6.9	82
136	N-Acetyl-seryl-aspartyl-lysyl-proline inhibits TGF-beta-mediated plasminogen activator inhibitor-1 expression via inhibition of Smad pathway in human mesangial cells. <i>Journal of the American Society of Nephrology: JASN</i> , 2003 , 14, 863-72	12.7	71
135	SIRT1 inactivation induces inflammation through the dysregulation of autophagy in human THP-1 cells. <i>Biochemical and Biophysical Research Communications</i> , 2012 , 427, 191-6	3.4	67
134	Sirtuins as possible drug targets in type 2 diabetes. <i>Current Drug Targets</i> , 2013 , 14, 622-36	3	67
133	Autophagy: A Novel Therapeutic Target for Diabetic Nephropathy. <i>Diabetes and Metabolism Journal</i> , 2015 , 39, 451-60	5	63
132	Genetic variations in the gene encoding TFAP2B are associated with type 2 diabetes mellitus. <i>Journal of Human Genetics</i> , 2005 , 50, 283-292	4.3	63
131	Regulating Autophagy as a Therapeutic Target for Diabetic Nephropathy. <i>Current Diabetes Reports</i> , 2017 , 17, 53	5.6	60
130	N-acetyl-seryl-aspartyl-lysyl-proline prevents renal insufficiency and mesangial matrix expansion in diabetic db/db mice. <i>Diabetes</i> , 2005 , 54, 838-45	0.9	60
129	N-acetyl-seryl-aspartyl-lysyl-proline inhibits diabetes-associated kidney fibrosis and endothelial-mesenchymal transition. <i>BioMed Research International</i> , 2014 , 2014, 696475	3	59
128	SIRT3 deficiency leads to induction of abnormal glycolysis in diabetic kidney with fibrosis. <i>Cell Death and Disease</i> , 2018 , 9, 997	9.8	59
127	Ipragliflozin improves mitochondrial abnormalities in renal tubules induced by a high-fat diet. <i>Journal of Diabetes Investigation</i> , 2018 , 9, 1025-1032	3.9	55
126	Association between single nucleotide polymorphisms within genes encoding sirtuin families and diabetic nephropathy in Japanese subjects with type 2 diabetes. <i>Clinical and Experimental Nephrology</i> , 2011 , 15, 381-390	2.5	54
125	A very-low-protein diet ameliorates advanced diabetic nephropathy through autophagy induction by suppression of the mTORC1 pathway in Wistar fatty rats, an animal model of type 2 diabetes and obesity. <i>Diabetologia</i> , 2016 , 59, 1307-17	10.3	54
124	The impact of dietary protein intake on longevity and metabolic health. <i>EBioMedicine</i> , 2019 , 43, 632-6408.8		53
123	Urinary Potassium Excretion and Renal and Cardiovascular Complications in Patients with Type 2 Diabetes and Normal Renal Function. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2015 , 10, 2152-8	6.9	52
122	Endothelin-1 induces cyclooxygenase-2 expression via nuclear factor of activated T-cell transcription factor in glomerular mesangial cells. <i>Journal of the American Society of Nephrology: JASN</i> , 2001 , 12, 1359-1368	12.7	52
121	Inhibition of Dipeptidyl Peptidase-4 Accelerates Epithelial-Mesenchymal Transition and Breast Cancer Metastasis via the CXCL12/CXCR4/mTOR Axis. <i>Cancer Research</i> , 2019 , 79, 735-746	10.1	52

120	PDGFR β Regulates Adipose Tissue Expansion and Glucose Metabolism via Vascular Remodeling in Diet-Induced Obesity. <i>Diabetes</i> , 2017 , 66, 1008-1021	0.9	51
119	Anti-aging molecule, Sirt1: a novel therapeutic target for diabetic nephropathy. <i>Archives of Pharmacal Research</i> , 2013 , 36, 230-6	6.1	50
118	The role of autophagy in the pathogenesis of diabetic nephropathy. <i>Journal of Diabetes Research</i> , 2013 , 2013, 193757	3.9	49
117	N-acetyl-seryl-aspartyl-lysyl-proline ameliorates the progression of renal dysfunction and fibrosis in WKY rats with established anti-glomerular basement membrane nephritis. <i>Journal of the American Society of Nephrology: JASN</i> , 2006 , 17, 674-85	12.7	47
116	Lipid mediators in diabetic nephropathy. <i>Fibrogenesis and Tissue Repair</i> , 2014 , 7, 12		45
115	Renal protective effects of empagliflozin via inhibition of EMT and aberrant glycolysis in proximal tubules. <i>JCI Insight</i> , 2020 , 5,	9.9	44
114	Effect of Antifibrotic MicroRNAs Crosstalk on the Action of N-acetyl-seryl-aspartyl-lysyl-proline in Diabetes-related Kidney Fibrosis. <i>Scientific Reports</i> , 2016 , 6, 29884	4.9	44
113	Role of nutrient-sensing signals in the pathogenesis of diabetic nephropathy. <i>BioMed Research International</i> , 2014 , 2014, 315494	3	42
112	Mammalian autophagy is essential for hepatic and renal ketogenesis during starvation. <i>Scientific Reports</i> , 2016 , 6, 18944	4.9	41
111	Dipeptidyl peptidase-4 and kidney fibrosis in diabetes. <i>Fibrogenesis and Tissue Repair</i> , 2016 , 9, 1		40
110	Calorie restriction in overweight males ameliorates obesity-related metabolic alterations and cellular adaptations through anti-aging effects, possibly including AMPK and SIRT1 activation. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2013 , 1830, 4820-7	4	39
109	d-alpha-tocopherol treatment prevents glomerular dysfunctions in diabetic rats through inhibition of protein kinase C-diacylglycerol pathway. <i>BioFactors</i> , 1998 , 7, 69-76	6.1	37
108	Anti-albuminuric effects of spironolactone in patients with type 2 diabetic nephropathy: a multicenter, randomized clinical trial. <i>Clinical and Experimental Nephrology</i> , 2015 , 19, 1098-106	2.5	36
107	FGFR1 is critical for the anti-endothelial mesenchymal transition effect of N-acetyl-seryl-aspartyl-lysyl-proline via induction of the MAP4K4 pathway. <i>Cell Death and Disease</i> , 2017 , 8, e2965	9.8	36
106	Eplerenone prevented obesity-induced inflammasome activation and glucose intolerance. <i>Journal of Endocrinology</i> , 2017 , 235, 179-191	4.7	34
105	Predictive properties of plasma amino acid profile for cardiovascular disease in patients with type 2 diabetes. <i>PLoS ONE</i> , 2014 , 9, e101219	3.7	34
104	Renal mitochondrial oxidative stress is enhanced by the reduction of Sirt3 activity, in Zucker diabetic fatty rats. <i>Redox Report</i> , 2018 , 23, 153-159	5.9	33
103	Combinational effect of genes for the renin-angiotensin system in conferring susceptibility to diabetic nephropathy. <i>Journal of Human Genetics</i> , 2007 , 52, 143-151	4.3	33

102	Linagliptin but not Sitagliptin inhibited transforming growth factor- β -induced endothelial DPP-4 activity and the endothelial-mesenchymal transition. <i>Biochemical and Biophysical Research Communications</i> , 2016 , 471, 184-90	3.4	32
101	Clinical therapeutic strategies for early stage of diabetic kidney disease. <i>World Journal of Diabetes</i> , 2014 , 5, 342-56	4.7	31
100	Inhibition of Angiotensin-Converting Enzyme Ameliorates Renal Fibrosis by Mitigating DPP-4 Level and Restoring Antifibrotic MicroRNAs. <i>Genes</i> , 2020 , 11,	4.2	30
99	Role of angiotensin II-mediated AMPK inactivation on obesity-related salt-sensitive hypertension. <i>Biochemical and Biophysical Research Communications</i> , 2012 , 418, 559-64	3.4	30
98	Loss of endothelial glucocorticoid receptor accelerates diabetic nephropathy. <i>Nature Communications</i> , 2021 , 12, 2368	17.4	30
97	Oral Administration of N-Acetyl-seryl-aspartyl-lysyl-proline Ameliorates Kidney Disease in Both Type 1 and Type 2 Diabetic Mice via a Therapeutic Regimen. <i>BioMed Research International</i> , 2016 , 2016, 9172157	3	30
96	Endothelial autophagy deficiency induces IL6 - dependent endothelial mesenchymal transition and organ fibrosis. <i>Autophagy</i> , 2020 , 16, 1905-1914	10.2	28
95	The Japanese clinical practice guideline for acute kidney injury 2016. <i>Clinical and Experimental Nephrology</i> , 2018 , 22, 985-1045	2.5	28
94	Deficiency in catechol-o-methyltransferase is linked to a disruption of glucose homeostasis in mice. <i>Scientific Reports</i> , 2017 , 7, 7927	4.9	26
93	Effects of high sodium intake and diuretics on the circadian rhythm of blood pressure in type 2 diabetic patients treated with an angiotensin II receptor blocker. <i>Clinical and Experimental Nephrology</i> , 2009 , 13, 300-306	2.5	26
92	Role of dietary amino acid balance in diet restriction-mediated lifespan extension, renoprotection, and muscle weakness in aged mice. <i>Aging Cell</i> , 2018 , 17, e12796	9.9	26
91	Autophagy in metabolic disease and ageing. <i>Nature Reviews Endocrinology</i> , 2021 , 17, 647-661	15.2	26
90	The Effect of Piceatannol from Passion Fruit (<i>Passiflora edulis</i>) Seeds on Metabolic Health in Humans. <i>Nutrients</i> , 2017 , 9,	6.7	25
89	Catechol-O-Methyltransferase Deficiency Leads to Hypersensitivity of the Pressor Response Against Angiotensin II. <i>Hypertension</i> , 2017 , 69, 1156-1164	8.5	24
88	1-Methylnicotinamide ameliorates lipotoxicity-induced oxidative stress and cell death in kidney proximal tubular cells. <i>Free Radical Biology and Medicine</i> , 2015 , 89, 831-41	7.8	24
87	Successful recovery of infective endocarditis-induced rapidly progressive glomerulonephritis by steroid therapy combined with antibiotics: a case report. <i>BMC Nephrology</i> , 2004 , 5, 18	2.7	24
86	Recent Insights Into SREBP as a Direct Mediator of Kidney Fibrosis via Lipid-Independent Pathways. <i>Frontiers in Pharmacology</i> , 2020 , 11, 265	5.6	23
85	N-acetyl-seryl-aspartyl-lysyl-proline: a valuable endogenous anti-fibrotic peptide for combating kidney fibrosis in diabetes. <i>Frontiers in Pharmacology</i> , 2014 , 5, 70	5.6	23

84	Secular changes in clinical manifestations of kidney disease among Japanese adults with type 2 diabetes from 1996 to 2014. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1032-1040	3.9	23
83	Elevation of the antifibrotic peptide N-acetyl-seryl-aspartyl-lysyl-proline: a blood pressure-independent beneficial effect of angiotensin I-converting enzyme inhibitors. <i>Fibrogenesis and Tissue Repair</i> , 2011 , 4, 25		22
82	Endothelial FGFR1 (Fibroblast Growth Factor Receptor 1) Deficiency Contributes Differential Fibrogenic Effects in Kidney and Heart of Diabetic Mice. <i>Hypertension</i> , 2020 , 76, 1935-1944	8.5	21
81	Metabolic reprogramming by N-acetyl-seryl-aspartyl-lysyl-proline protects against diabetic kidney disease. <i>British Journal of Pharmacology</i> , 2020 , 177, 3691-3711	8.6	21
80	A Low-Protein Diet for Diabetic Kidney Disease: Its Effect and Molecular Mechanism, an Approach from Animal Studies. <i>Nutrients</i> , 2018 , 10,	6.7	21
79	CD38 inhibition by apigenin ameliorates mitochondrial oxidative stress through restoration of the intracellular NAD/NADH ratio and Sirt3 activity in renal tubular cells in diabetic rats. <i>Aging</i> , 2020 , 12, 11325-11336	5.6	21
78	Cancer biology in diabetes. <i>Journal of Diabetes Investigation</i> , 2014 , 5, 251-64	3.9	20
77	Pituitary apoplexy following gonadotropin-releasing hormone agonist administration with gonadotropin-secreting pituitary adenoma. <i>Journal of Clinical Neuroscience</i> , 2015 , 22, 601-3	2.2	20
76	The Japanese Clinical Practice Guideline for acute kidney injury 2016. <i>Journal of Intensive Care</i> , 2018 , 6, 48	7	18
75	MicroRNA148b-3p inhibits mTORC1-dependent apoptosis in diabetes by repressing TNFR2 in proximal tubular cells. <i>Kidney International</i> , 2016 , 90, 1211-1225	9.9	18
74	Dipeptidyl peptidase-4 plays a pathogenic role in BSA-induced kidney injury in diabetic mice. <i>Scientific Reports</i> , 2019 , 9, 7519	4.9	17
73	The relevance of the Renin-Angiotensin system in the development of drugs to combat preeclampsia. <i>International Journal of Endocrinology</i> , 2015 , 2015, 572713	2.7	17
72	Manganese Superoxide Dismutase Dysfunction and the Pathogenesis of Kidney Disease. <i>Frontiers in Physiology</i> , 2020 , 11, 755	4.6	17
71	AMP-Activated Protein (AMPK) in Pathophysiology of Pregnancy Complications. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	17
70	Lamp-2 deficiency prevents high-fat diet-induced obese diabetes via enhancing energy expenditure. <i>Biochemical and Biophysical Research Communications</i> , 2015 , 465, 249-55	3.4	16
69	A low-protein diet exerts a beneficial effect on diabetic status and prevents diabetic nephropathy in Wistar fatty rats, an animal model of type 2 diabetes and obesity. <i>Nutrition and Metabolism</i> , 2018 , 15, 20	4.6	14
68	Endothelial SIRT3 regulates myofibroblast metabolic shifts in diabetic kidneys. <i>IScience</i> , 2021 , 24, 102396	6.1	14
67	Conditions, pathogenesis, and progression of diabetic kidney disease and early decliner in Japan. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	13

66	Dipeptidyl peptidase-4 inhibition and renoprotection: the role of antifibrotic effects. <i>Current Opinion in Nephrology and Hypertension</i> , 2017 , 26, 56-66	3.5	13
65	Statin use in patients with diabetes and kidney disease: the Japanese experience. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013 , 20, 407-24	4	13
64	The PKM2 activator TEPP-46 suppresses kidney fibrosis via inhibition of the EMT program and aberrant glycolysis associated with suppression of HIF-1 α accumulation. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 697-709	3.9	13
63	Identification of subgroups of patients with type 2 diabetes with differences in renal function preservation, comparing patients receiving sodium-glucose co-transporter-2 inhibitors with those receiving dipeptidyl peptidase-4 inhibitors, using a supervised machine-learning algorithm	6.7	12
62	A ketogenic amino acid rich diet benefits mitochondrial homeostasis by altering the AKT/4EBP1 and autophagy signaling pathways in the gastrocnemius and soleus. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2018 , 1862, 1547-1555	4	12
61	Hypothalamic AMP-Activated Protein Kinase Regulates Biphasic Insulin Secretion from Pancreatic β Cells during Fasting and in Type 2 Diabetes. <i>EBioMedicine</i> , 2016 , 13, 168-180	8.8	12
60	Decline in estimated glomerular filtration rate is associated with risk of end-stage renal disease in type 2 diabetes with macroalbuminuria: an observational study from JDNCS. <i>Clinical and Experimental Nephrology</i> , 2018 , 22, 377-387	2.5	11
59	Therapeutic management of diabetic kidney disease. <i>Journal of Diabetes Investigation</i> , 2011 , 2, 248-54	3.9	11
58	Stromal cell-derived factor 1 (SDF1) attenuates platelet-derived growth factor-B (PDGF-B)-induced vascular remodeling for adipose tissue expansion in obesity. <i>Angiogenesis</i> , 2020 , 23, 667-684	10.6	11
57	Prevalence of albuminuria and renal dysfunction, and related clinical factors in Japanese patients with diabetes: The Japan Diabetes Complication and its Prevention prospective study \square . <i>Journal of Diabetes Investigation</i> , 2020 , 11, 325-332	3.9	11
56	FGFR1 is essential for N-acetyl-seryl-aspartyl-lysyl-proline regulation of mitochondrial dynamics by upregulating microRNA let-7b-5p. <i>Biochemical and Biophysical Research Communications</i> , 2018 , 495, 2214-2220 ¹⁰	2.4	10
55	Efficacy and tolerability of vildagliptin in type 2 diabetic patients on hemodialysis. <i>Journal of Diabetes Investigation</i> , 2012 , 3, 298-301	3.9	10
54	Methionine abrogates the renoprotective effect of a low-protein diet against diabetic kidney disease in obese rats with type 2 diabetes. <i>Aging</i> , 2020 , 12, 4489-4505	5.6	10
53	Relationship Between Autophagy and Metabolic Syndrome Characteristics in the Pathogenesis of Atherosclerosis. <i>Frontiers in Cell and Developmental Biology</i> , 2021 , 9, 641852	5.7	10
52	The impact of mitochondrial quality control by Sirtuins on the treatment of type 2 diabetes and diabetic kidney disease. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2020 , 1866, 165756	6.9	9
51	Deficiency in Dipeptidyl Peptidase-4 Promotes Chemoresistance through the CXCL12/CXCR4/mTOR/TGF β Signaling Pathway in Breast Cancer Cells. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	9
50	Metformin Mitigates DPP-4 Inhibitor-Induced Breast Cancer Metastasis via Suppression of mTOR Signaling. <i>Molecular Cancer Research</i> , 2021 , 19, 61-73	6.6	9
49	Mechanism of Activation of Mechanistic Target of Rapamycin Complex 1 by Methionine. <i>Frontiers in Cell and Developmental Biology</i> , 2020 , 8, 715	5.7	8

48	Comparative Effects of Direct Renin Inhibitor and Angiotensin Receptor Blocker on Albuminuria in Hypertensive Patients with Type 2 Diabetes. A Randomized Controlled Trial. <i>PLoS ONE</i> , 2016 , 11, e0164936	3.7	8
47	Effect of Methionine Restriction on Aging: Its Relationship to Oxidative Stress. <i>Biomedicines</i> , 2021 , 9,	4.8	8
46	Significance of SGLT2 inhibitors: lessons from renal clinical outcomes in patients with type 2 diabetes and basic researches. <i>Diabetology International</i> , 2020 , 11, 245-251	2.3	7
45	Evaluation of a new care system provided to diabetic patients in the outpatient clinic. <i>Internal Medicine</i> , 2000 , 39, 783-7	1.1	7
44	Dapagliflozin Restores Impaired Autophagy and Suppresses Inflammation in High Glucose-Treated HK-2 Cells. <i>Cells</i> , 2021 , 10,	7.9	7
43	Randomized trial of an intensified, multifactorial intervention in patients with advanced-stage diabetic kidney disease: Diabetic Nephropathy Remission and Regression Team Trial in Japan (DNETT-Japan). <i>Journal of Diabetes Investigation</i> , 2021 , 12, 207-216	3.9	7
42	Pro-inflammatory macrophages coupled with glycolysis remodel adipose vasculature by producing platelet-derived growth factor-B in obesity. <i>Scientific Reports</i> , 2020 , 10, 670	4.9	6
41	Effect of switching to teneligliptin from other dipeptidyl peptidase-4 inhibitors on glucose control and renoprotection in type 2 diabetes patients with diabetic kidney disease. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 706-713	3.9	6
40	Three ileus cases associated with the use of dipeptidyl peptidase-4 inhibitors in diabetic patients. <i>Journal of Diabetes Investigation</i> , 2013 , 4, 673-5	3.9	6
39	Anagliptin ameliorates albuminuria and urinary liver-type fatty acid-binding protein excretion in patients with type 2 diabetes with nephropathy in a glucose-lowering-independent manner. <i>BMJ Open Diabetes Research and Care</i> , 2017 , 5, e000391	4.5	6
38	A new classification of Diabetic Nephropathy 2014: a report from Joint Committee on Diabetic Nephropathy. <i>Diabetology International</i> , 2014 , 5, 207-211	2.3	6
37	CD26/DPP-4: Type 2 Diabetes Drug Target with Potential Influence on Cancer Biology. <i>Cancers</i> , 2021 , 13,	6.6	6
36	Interventions against nutrient-sensing pathways represent an emerging new therapeutic approach for diabetic nephropathy. <i>Clinical and Experimental Nephrology</i> , 2014 , 18, 210-3	2.5	5
35	Diabetic Nephropathy Remission and Regression Team Trial in Japan (DNETT-Japan): Rationale and study design. <i>Diabetes Research and Clinical Practice</i> , 2010 , 87, 228-32	7.4	5
34	Interactions among Long Non-Coding RNAs and microRNAs Influence Disease Phenotype in Diabetes and Diabetic Kidney Disease. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	5
33	Sirtuins and Renal Oxidative Stress. <i>Antioxidants</i> , 2021 , 10,	7.1	5
32	Resveratrol is essential for the anti-endothelial mesenchymal transition effects of N-acetyl-seryl-aspartyl-lysyl-proline. <i>FEBS Open Bio</i> , 2019 , 9, 1029-1038	2.7	4
31	Supplementation with Red Wine Extract Increases Insulin Sensitivity and Peripheral Blood Mononuclear Sirt1 Expression in Nondiabetic Humans. <i>Nutrients</i> , 2020 , 12,	6.7	4

30	Role of Sirt1 as a Regulator of Autophagy 2016 , 89-100		4
29	Relevance of Autophagy Induction by Gastrointestinal Hormones: Focus on the Incretin-Based Drug Target and Glucagon. <i>Frontiers in Pharmacology</i> , 2019 , 10, 476	5.6	3
28	N-Acetyl-seryl-aspartyl-lysyl-proline is a potential biomarker of renal function in normoalbuminuric diabetic patients with eGFR ≥ 30 ml/min/1.73 m ² . <i>Clinical and Experimental Nephrology</i> , 2019 , 23, 1004-1012	2.5	3
27	Cyclic and intermittent very low-protein diet can have beneficial effects against advanced diabetic nephropathy in Wistar Fatty (fa/fa) rats, an animal model of type 2 diabetes and obesity. <i>Nephrology</i> , 2017 , 22, 1030-1034	2.2	3
26	Epidermal growth factor receptor signaling and the progression of diabetic nephropathy. <i>Journal of Diabetes Investigation</i> , 2015 , 6, 519-21	3.9	3
25	Medical nutrition therapy and dietary counseling for patients with diabetes-energy, carbohydrates, protein intake and dietary counseling. <i>Diabetology International</i> , 2020 , 11, 224-239	2.3	3
24	Sodium-glucose cotransporter 2 inhibitors in type 2 diabetes patients with renal function impairment slow the annual renal function decline, in a real clinical practice. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1577-1585	3.9	3
23	The Japanese Clinical Practice Guideline for acute kidney injury 2016. <i>Renal Replacement Therapy</i> , 2018 , 4,	2.3	3
22	Levofloxacin-induced Achilles tendon rupture in a patient with systemic microscopic polyangiitis. <i>Modern Rheumatology</i> , 2005 , 15, 217-219	3.3	2
21	Anterior pituitary function in Rathke's cleft cysts versus nonfunctioning pituitary adenomas. <i>Endocrine Journal</i> , 2021 , 68, 943-952	2.9	2
20	CD-1 mice: A novel type 2 diabetic mouse model with progressive kidney fibrosis. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1470-1481	3.9	1
19	Case report of superior mesenteric artery syndrome that developed in a lean type 2 diabetes patient and was associated with rapid body weight loss after sodium-glucose cotransporter 2 inhibitor administration. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1359-1362	3.9	1
18	Diabetic kidney disease: Its current trends and future therapeutic perspectives. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1174-1176	3.9	1
17	Exercise Ameliorates Diabetic Kidney Disease in Type 2 Diabetic Fatty Rats. <i>Antioxidants</i> , 2021 , 10,	7.1	1
16	NAD Homeostasis in Diabetic Kidney Disease. <i>Frontiers in Medicine</i> , 2021 , 8, 703076	4.9	1
15	Rapid enlargement of an intracranial germ cell tumor after gonadotropin hormone therapy. <i>Journal of Clinical Neuroscience</i> , 2016 , 31, 185-8	2.2	1
14	Dietary Magnesium Insufficiency Induces Salt-Sensitive Hypertension in Mice Associated With Reduced Kidney Catechol-O-Methyl Transferase Activity. <i>Hypertension</i> , 2021 , 78, 138-150	8.5	1
13	Effects of SGLT2 Inhibitors on Atherosclerosis: Lessons from Cardiovascular Clinical Outcomes in Type 2 Diabetic Patients and Basic Researches.. <i>Journal of Clinical Medicine</i> , 2021 , 11,	5.1	1

12	Impact of empagliflozin on diabetic kidney disease. <i>Journal of Diabetes Investigation</i> , 2017 , 8, 658-660	3.9	o
11	Restoration of the Hypothalamic-pituitary-adrenal Response to Hypoglycemia in Type 2 Diabetes by Avoiding Chronic Hypoglycemia. <i>Internal Medicine</i> , 2016 , 55, 3471-3473	1.1	o
10	Severe electrolytes disorders with the interstitial kidney alterations in the patient with the history of total thyroidectomy and parathyroidectomy: possible role of vitamin D deficiency. <i>Clinical Case Reports (discontinued)</i> , 2018 , 6, 983-989	0.7	
9	Corrigendum to [Legumain/asparaginyl endopeptidase controls extracellular matrix remodeling through the degradation of fibronectin in mouse renal proximal tubular cells][FEBS Lett. 581 (2007) 1417-1424]. <i>FEBS Letters</i> , 2007 , 581, 3579-3579	3.8	
8	Insulin-like growth factor I stimulates glucose uptake and expression of glucose transporter 1 in cultured mesangial cells. <i>Clinical and Experimental Nephrology</i> , 1999 , 3, 159-162	2.5	
7	Backcross db Gene into CD-1 Background Results in Novel Type 2 Diabetic Mouse Model with Progressive Kidney Fibrosis. <i>Diabetes</i> , 2018 , 67, 500-P	0.9	
6	Renal Mitochondrial Oxidative Stress Induced by NAD ⁺ -Dependent Sirt3 Inactivation via Overexpression of CD38 (NAD ⁺ ase) in Diabetic Kidney Disease. <i>Diabetes</i> , 2018 , 67, 495-P	0.9	
5	513-P: Adenosine Signal Plays an Important Role in Renoprotective Effects of SGLT2 Inhibitor in Proteinuric Diabetic Mice. <i>Diabetes</i> , 2019 , 68, 513-P	0.9	
4	463-P: Linagliptin Ameliorated Heart Damage Associated with the Suppression of Necroptosis in Type 1 Diabetic Mice. <i>Diabetes</i> , 2019 , 68, 463-P	0.9	
3	Patient Assessment and Diagnosis 2017 , 47-56		
2	Rapid effects of pitavastatin on uric acid homeostasis. <i>Gout and Nucleic Acid Metabolism</i> , 2011 , 35, 39-47		
1	Novel Missense Mutation p.Ile424Ser in an Individual with Multiple Hepatic Cysts: A Case Report.. <i>Medicines (Basel, Switzerland)</i> , 2022 , 9,	4.1	