

Daisuke Koya

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8479150/publications.pdf>

Version: 2024-02-01

161
papers

13,090
citations

41258

49
h-index

24179

110
g-index

173
all docs

173
docs citations

173
times ranked

22195
citing authors

#	ARTICLE	IF	CITATIONS
1	Adenosine/A1R signaling pathway did not play dominant roles on the influence of SGLT2 inhibitor in the kidney of BSAâ€œoverloaded STZâ€œinduced diabetic mice. <i>Journal of Diabetes Investigation</i> , 2022, , .	1.1	1
2	Novel PKD2 Missense Mutation p.Ile424Ser in an Individual with Multiple Hepatic Cysts: A Case Report. <i>Medicines (Basel, Switzerland)</i> , 2022, 9, 25.	0.7	0
3	Effects of SGLT2 Inhibitors on Atherosclerosis: Lessons from Cardiovascular Clinical Outcomes in Type 2 Diabetic Patients and Basic Researches. <i>Journal of Clinical Medicine</i> , 2022, 11, 137.	1.0	15
4	Rationale, Design and Baseline Characteristics of the Effect of Canagliflozin in Type 2 Diabetic Patients with Microalbuminuria in Japanese Population (<scp>CANPIONE</scp>) study. <i>Diabetes, Obesity and Metabolism</i> , 2022, , .	2.2	1
5	Randomized trial of an intensified, multifactorial intervention in patients with advancedâ€œstage diabetic kidney disease: Diabetic Nephropathy Remission and Regression Team Trial in Japan (DNETTâ€œJapan). <i>Journal of Diabetes Investigation</i> , 2021, 12, 207-216.	1.1	17
6	Metformin Mitigates DPP-4 Inhibitor-Induced Breast Cancer Metastasis via Suppression of mTOR Signaling. <i>Molecular Cancer Research</i> , 2021, 19, 61-73.	1.5	22
7	The PKM2 activator TEPPâ€œ46 suppresses kidney fibrosis via inhibition of the EMT program and aberrant glycolysis associated with suppression of HIFâ€œ1â€œ accumulation. <i>Journal of Diabetes Investigation</i> , 2021, 12, 697-709.	1.1	44
8	Anterior pituitary function in Rathkeâ€œ's cleft cysts <i>> versus</i> nonfunctioning pituitary adenomas. <i>Endocrine Journal</i> , 2021, 68, 943-952.	0.7	5
9	Effect of Methionine Restriction on Aging: Its Relationship to Oxidative Stress. <i>Biomedicines</i> , 2021, 9, 130.	1.4	39
10	Sodiumâ€œglucose cotransporterâ€œ2 inhibitors in typeâ€œ2 diabetes patients with renal function impairment slow the annual renal function decline, in a real clinical practice. <i>Journal of Diabetes Investigation</i> , 2021, 12, 1577-1585.	1.1	6
11	Relationship Between Autophagy and Metabolic Syndrome Characteristics in the Pathogenesis of Atherosclerosis. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 641852.	1.8	26
12	Loss of endothelial glucocorticoid receptor accelerates diabetic nephropathy. <i>Nature Communications</i> , 2021, 12, 2368.	5.8	79
13	CD26/DPP-4: Type 2 Diabetes Drug Target with Potential Influence on Cancer Biology. <i>Cancers</i> , 2021, 13, 2191.	1.7	20
14	Endothelial SIRT3 regulates myofibroblast metabolic shifts in diabetic kidneys. <i>IScience</i> , 2021, 24, 102390.	1.9	50
15	Interactions among Long Non-Coding RNAs and microRNAs Influence Disease Phenotype in Diabetes and Diabetic Kidney Disease. <i>International Journal of Molecular Sciences</i> , 2021, 22, 6027.	1.8	19
16	Dapagliflozin Restores Impaired Autophagy and Suppresses Inflammation in High Glucose-Treated HK-2 Cells. <i>Cells</i> , 2021, 10, 1457.	1.8	60
17	NAD+ Homeostasis in Diabetic Kidney Disease. <i>Frontiers in Medicine</i> , 2021, 8, 703076.	1.2	10
18	Dietary Magnesium Insufficiency Induces Salt-Sensitive Hypertension in Mice Associated With Reduced Kidney Catechol-O-Methyl Transferase Activity. <i>Hypertension</i> , 2021, 78, 138-150.	1.3	4

#	ARTICLE	IF	CITATIONS
19	Sirtuins and Renal Oxidative Stress. <i>Antioxidants</i> , 2021, 10, 1198.	2.2	27
20	Autophagy in metabolic disease and ageing. <i>Nature Reviews Endocrinology</i> , 2021, 17, 647-661.	4.3	159
21	Exercise Ameliorates Diabetic Kidney Disease in Type 2 Diabetic Fatty Rats. <i>Antioxidants</i> , 2021, 10, 1754.	2.2	8
22	Prevalence of albuminuria and renal dysfunction, and related clinical factors in Japanese patients with diabetes: The Japan Diabetes Complication and its Prevention prospective study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 325-332.	1.1	21
23	Stromal cell-derived factor 1 (SDF1) attenuates platelet-derived growth factor-B (PDGF-B)-induced vascular remodeling for adipose tissue expansion in obesity. <i>Angiogenesis</i> , 2020, 23, 667-684.	3.7	19
24	Manganese Superoxide Dismutase Dysfunction and the Pathogenesis of Kidney Disease. <i>Frontiers in Physiology</i> , 2020, 11, 755.	1.3	52
25	Medical nutrition therapy and dietary counseling for patients with diabetes-energy, carbohydrates, protein intake and dietary counseling. <i>Diabetology International</i> , 2020, 11, 224-239.	0.7	7
26	Supplementation with Red Wine Extract Increases Insulin Sensitivity and Peripheral Blood Mononuclear Sirt1 Expression in Nondiabetic Humans. <i>Nutrients</i> , 2020, 12, 3108.	1.7	8
27	Mechanism of Activation of Mechanistic Target of Rapamycin Complex 1 by Methionine. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 715.	1.8	21
28	Endothelial FGFR1 (Fibroblast Growth Factor Receptor 1) Deficiency Contributes Differential Fibrogenic Effects in Kidney and Heart of Diabetic Mice. <i>Hypertension</i> , 2020, 76, 1935-1944.	1.3	55
29	Metabolic reprogramming by N-acetylcysteine protects against diabetic kidney disease. <i>British Journal of Pharmacology</i> , 2020, 177, 3691-3711.	2.7	42
30	CD44 ^{hi} db/db mice: A novel type 2 diabetic mouse model with progressive kidney fibrosis. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1470-1481.	1.1	5
31	Significance of SGLT2 inhibitors: lessons from renal clinical outcomes in patients with type 2 diabetes and basic researches. <i>Diabetology International</i> , 2020, 11, 245-251.	0.7	13
32	The impact of mitochondrial quality control by Sirtuins on the treatment of type 2 diabetes and diabetic kidney disease. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2020, 1866, 165756.	1.8	15
33	Case report of superior mesenteric artery syndrome that developed in a lean type 2 diabetes patient and was associated with rapid body weight loss after sodium-glucose cotransporter 2 inhibitor administration. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1359-1362.	1.1	3
34	Inhibition of Angiotensin-Converting Enzyme Ameliorates Renal Fibrosis by Mitigating DPP-4 Level and Restoring Antifibrotic MicroRNAs. <i>Genes</i> , 2020, 11, 211.	1.0	54
35	Endothelial autophagy deficiency induces IL6 - dependent endothelial mesenchymal transition and organ fibrosis. <i>Autophagy</i> , 2020, 16, 1905-1914.	4.3	65
36	Pro-inflammatory macrophages coupled with glycolysis remodel adipose vasculature by producing platelet-derived growth factor-B in obesity. <i>Scientific Reports</i> , 2020, 10, 670.	1.6	18

#	ARTICLE	IF	CITATIONS
37	Deficiency in Dipeptidyl Peptidase-4 Promotes Chemoresistance Through the CXCL12/CXCR4/mTOR/TGF β 2 Signaling Pathway in Breast Cancer Cells. <i>International Journal of Molecular Sciences</i> , 2020, 21, 805.	1.8	18
38	Conditions, pathogenesis, and progression of diabetic kidney disease and early decliner in Japan. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e000902.	1.2	31
39	Recent Insights Into SREBP as a Direct Mediator of Kidney Fibrosis via Lipid-Independent Pathways. <i>Frontiers in Pharmacology</i> , 2020, 11, 265.	1.6	53
40	Renal protective effects of empagliflozin via inhibition of EMT and aberrant glycolysis in proximal tubules. <i>JCI Insight</i> , 2020, 5, .	2.3	131
41	Methionine abrogates the renoprotective effect of a low-protein diet against diabetic kidney disease in obese rats with type 2 diabetes. <i>Aging</i> , 2020, 12, 4489-4505.	1.4	18
42	CD38 inhibition by apigenin ameliorates mitochondrial oxidative stress through restoration of the intracellular NAD ⁺ /NADH ratio and Sirt3 activity in renal tubular cells in diabetic rats. <i>Aging</i> , 2020, 12, 11325-11336.	1.4	61
43	Efficacy of SGLT2 inhibitor in type 2 diabetic patients under dietary instructions: A pilot study. <i>Clinical and Medical Investigations</i> , 2020, 5, .	0.3	1
44	Effect of switching to teneligliptin from other dipeptidyl peptidase-4 inhibitors on glucose control and renoprotection in type 2 diabetes patients with diabetic kidney disease. <i>Journal of Diabetes Investigation</i> , 2019, 10, 706-713.	1.1	7
45	Diabetic kidney disease: Its current trends and future therapeutic perspectives. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1174-1176.	1.1	3
46	Relevance of Autophagy Induction by Gastrointestinal Hormones: Focus on the Incretin-Based Drug Target and Glucagon. <i>Frontiers in Pharmacology</i> , 2019, 10, 476.	1.6	11
47	Proposal of classification of "chronic kidney disease (CKD) with diabetes" in clinical setting. <i>Diabetology International</i> , 2019, 10, 180-182.	0.7	1
48	Dipeptidyl peptidase-4 plays a pathogenic role in BSA-induced kidney injury in diabetic mice. <i>Scientific Reports</i> , 2019, 9, 7519.	1.6	25
49	Identification of subgroups of patients with type 2 diabetes with differences in renal function preservation, comparing patients receiving sodium-glucose cotransporter-2 inhibitors with those receiving dipeptidyl peptidase-4 inhibitors, using a supervised machine learning algorithm (PROFILE). <i>Tj ETQq1 1202784314sgBT /O Metabolism</i> , 2019, 21, 1025-1034.		
50	Klotho is essential for the anti-endothelial mesenchymal transition effects of N-acetyl-seryl-aspartyl-lysyl-proline. <i>FEBS Open Bio</i> , 2019, 9, 1029-1038.	1.0	7
51	Sirtuins and Type 2 Diabetes: Role in Inflammation, Oxidative Stress, and Mitochondrial Function. <i>Frontiers in Endocrinology</i> , 2019, 10, 187.	1.5	170
52	N-Acetyl-seryl-aspartyl-lysyl-proline is a potential biomarker of renal function in normoalbuminuric diabetic patients with eGFR ≥ 30 mL/min/1.73 m ² . <i>Clinical and Experimental Nephrology</i> , 2019, 23, 1004-1012.	0.7	5
53	The impact of dietary protein intake on longevity and metabolic health. <i>EBioMedicine</i> , 2019, 43, 632-640.	2.7	97
54	Inhibition of Dipeptidyl Peptidase-4 Accelerates Epithelial-Mesenchymal Transition and Breast Cancer Metastasis via the CXCL12/CXCR4/mTOR Axis. <i>Cancer Research</i> , 2019, 79, 735-746.	0.4	86

#	ARTICLE	IF	CITATIONS
55	Secular changes in clinical manifestations of kidney disease among Japanese adults with type 2 diabetes from 1996 to 2014. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1032-1040.	1.1	39
56	513-P: Adenosine Signal Plays an Important Role in Renoprotective Effects of SGLT2 Inhibitor in Proteinuric Diabetic Mice. <i>Diabetes</i> , 2019, 68, 513-P.	0.3	0
57	463-P: Linagliptin Ameliorated Heart Damage Associated with the Suppression of Necroptosis in Type 1 Diabetic Mice. <i>Diabetes</i> , 2019, 68, 463-P.	0.3	0
58	Ipragliflozin improves mitochondrial abnormalities in renal tubules induced by a high-fat diet. <i>Journal of Diabetes Investigation</i> , 2018, 9, 1025-1032.	1.1	88
59	FGFR1 is essential for N-acetyl-seryl-aspartyl-lysyl-proline regulation of mitochondrial dynamics by upregulating microRNA let-7b-5p. <i>Biochemical and Biophysical Research Communications</i> , 2018, 495, 2214-2220.	1.0	13
60	Severe electrolytes disorders with the interstitial kidney alterations in the patient with the history of total thyroidectomy and parathyroidectomy: possible role of vitamin D deficiency. <i>Clinical Case Reports (discontinued)</i> , 2018, 6, 983-989.	0.2	0
61	A ketogenic amino acid rich diet benefits mitochondrial homeostasis by altering the AKT/4EBP1 and autophagy signaling pathways in the gastrocnemius and soleus. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2018, 1862, 1547-1555.	1.1	17
62	Decline in estimated glomerular filtration rate is associated with risk of end-stage renal disease in type 2 diabetes with macroalbuminuria: an observational study from JDNCS. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 377-387.	0.7	14
63	SIRT3 deficiency leads to induction of abnormal glycolysis in diabetic kidney with fibrosis. <i>Cell Death and Disease</i> , 2018, 9, 997.	2.7	117
64	AMP-Activated Protein (AMPK) in Pathophysiology of Pregnancy Complications. <i>International Journal of Molecular Sciences</i> , 2018, 19, 3076.	1.8	26
65	The Japanese Clinical Practice Guideline for acute kidney injury 2016. <i>Renal Replacement Therapy</i> , 2018, 4, .	0.3	4
66	Role of dietary amino acid balance in diet restriction-mediated lifespan extension, renoprotection, and muscle weakness in aged mice. <i>Aging Cell</i> , 2018, 17, e12796.	3.0	45
67	The Japanese clinical practice guideline for acute kidney injury 2016. <i>Clinical and Experimental Nephrology</i> , 2018, 22, 985-1045.	0.7	40
68	A Low-Protein Diet for Diabetic Kidney Disease: Its Effect and Molecular Mechanism, an Approach from Animal Studies. <i>Nutrients</i> , 2018, 10, 544.	1.7	38
69	A low-protein diet exerts a beneficial effect on diabetic status and prevents diabetic nephropathy in Wistar fatty rats, an animal model of type 2 diabetes and obesity. <i>Nutrition and Metabolism</i> , 2018, 15, 20.	1.3	23
70	The Japanese Clinical Practice Guideline for acute kidney injury 2016. <i>Journal of Intensive Care</i> , 2018, 6, 48.	1.3	35
71	Renal mitochondrial oxidative stress is enhanced by the reduction of Sirt3 activity, in Zucker diabetic fatty rats. <i>Redox Report</i> , 2018, 23, 153-159.	1.4	42
72	Backcross db Gene into CD-1 Background Results in Novel Type 2 Diabetic Mouse Model with Progressive Kidney Fibrosis. <i>Diabetes</i> , 2018, 67, 500-P.	0.3	0

#	ARTICLE	IF	CITATIONS
73	Renal Mitochondrial Oxidative Stress Induced by NAD ⁺ -Dependent Sirt3 Inactivation via Overexpression of CD38 (NAD ⁺ ase) in Diabetic Kidney Disease. <i>Diabetes</i> , 2018, 67, 495-P.	0.3	0
74	PDGFR β Regulates Adipose Tissue Expansion and Glucose Metabolism via Vascular Remodeling in Diet-Induced Obesity. <i>Diabetes</i> , 2017, 66, 1008-1021.	0.3	66
75	Catechol-O-Methyltransferase Deficiency Leads to Hypersensitivity of the Pressor Response Against Angiotensin II. <i>Hypertension</i> , 2017, 69, 1156-1164.	1.3	28
76	Regulating Autophagy as a Therapeutic Target for Diabetic Nephropathy. <i>Current Diabetes Reports</i> , 2017, 17, 53.	1.7	79
77	Impact of empagliflozin on diabetic kidney disease. <i>Journal of Diabetes Investigation</i> , 2017, 8, 658-660.	1.1	1
78	Eplerenone prevented obesity-induced inflammasome activation and glucose intolerance. <i>Journal of Endocrinology</i> , 2017, 235, 179-191.	1.2	51
79	Deficiency in catechol-o-methyltransferase is linked to a disruption of glucose homeostasis in mice. <i>Scientific Reports</i> , 2017, 7, 7927.	1.6	30
80	Anagliptin ameliorates albuminuria and urinary liver-type fatty acid-binding protein excretion in patients with type 2 diabetes with nephropathy in a glucose-lowering-independent manner. <i>BMJ Open Diabetes Research and Care</i> , 2017, 5, e000391.	1.2	7
81	Dipeptidyl peptidase-4 inhibition and renoprotection. <i>Current Opinion in Nephrology and Hypertension</i> , 2017, 26, 56-66.	1.0	16
82	Cyclic and intermittent very low-protein diet can have beneficial effects against advanced diabetic nephropathy in Wistar fatty (<i>fa/fa</i>) rats, an animal model of type 2 diabetes and obesity. <i>Nephrology</i> , 2017, 22, 1030-1034.	0.7	5
83	FGFR1 is critical for the anti-endothelial mesenchymal transition effect of N-acetyl-seryl-aspartyl-lysyl-proline via induction of the MAP4K4 pathway. <i>Cell Death and Disease</i> , 2017, 8, e2965-e2965.	2.7	61
84	The Effect of Piceatannol from Passion Fruit (<i>Passiflora edulis</i>) Seeds on Metabolic Health in Humans. <i>Nutrients</i> , 2017, 9, 1142.	1.7	38
85	Patient Assessment and Diagnosis. , 2017, , 47-56.		0
86	Oral Administration of N-Acetyl-seryl-aspartyl-lysyl-proline Ameliorates Kidney Disease in Both Type 1 and Type 2 Diabetic Mice via a Therapeutic Regimen. <i>BioMed Research International</i> , 2016, 2016, 1-11.	0.9	36
87	Role of Sirt1 as a Regulator of Autophagy. , 2016, , 89-100.		13
88	Comparative Effects of Direct Renin Inhibitor and Angiotensin Receptor Blocker on Albuminuria in Hypertensive Patients with Type 2 Diabetes. A Randomized Controlled Trial. <i>PLoS ONE</i> , 2016, 11, e0164936.	1.1	11
89	Rodent models of diabetic nephropathy: their utility and limitations. <i>International Journal of Nephrology and Renovascular Disease</i> , 2016, Volume 9, 279-290.	0.8	190
90	The protective role of Sirt1 in vascular tissue: its relationship to vascular aging and atherosclerosis. <i>Aging</i> , 2016, 8, 2290-2307.	1.4	201

#	ARTICLE	IF	CITATIONS
91	Effect of Antifibrotic MicroRNAs Crosstalk on the Action of N-acetyl-seryl-aspartyl-lysyl-proline in Diabetes-related Kidney Fibrosis. <i>Scientific Reports</i> , 2016, 6, 29884.	1.6	60
92	Mammalian autophagy is essential for hepatic and renal ketogenesis during starvation. <i>Scientific Reports</i> , 2016, 6, 18944.	1.6	58
93	A very-low-protein diet ameliorates advanced diabetic nephropathy through autophagy induction by suppression of the mTORC1 pathway in Wistar fatty rats, an animal model of type 2 diabetes and obesity. <i>Diabetologia</i> , 2016, 59, 1307-1317.	2.9	75
94	Rapid enlargement of an intracranial germ cell tumor after gonadotropin hormone therapy. <i>Journal of Clinical Neuroscience</i> , 2016, 31, 185-188.	0.8	5
95	MicroRNA148b-3p inhibits mTORC1-dependent apoptosis in diabetes by repressing TNFR2 in proximal tubular cells. <i>Kidney International</i> , 2016, 90, 1211-1225.	2.6	27
96	Hypothalamic AMP-Activated Protein Kinase Regulates Biphasic Insulin Secretion from Pancreatic β Cells during Fasting and in Type 2 Diabetes. <i>EBioMedicine</i> , 2016, 13, 168-180.	2.7	14
97	Dipeptidyl peptidase-4 and kidney fibrosis in diabetes. <i>Fibrogenesis and Tissue Repair</i> , 2016, 9, 1.	3.4	50
98	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016, 12, 1-222.	4.3	4,701
99	Linagliptin but not Sitagliptin inhibited transforming growth factor- β 2-induced endothelial DPP-4 activity and the endothelial-mesenchymal transition. <i>Biochemical and Biophysical Research Communications</i> , 2016, 471, 184-190.	1.0	38
100	Impaired Podocyte Autophagy Exacerbates Proteinuria in Diabetic Nephropathy. <i>Diabetes</i> , 2016, 65, 755-767.	0.3	243
101	Restoration of the Hypothalamic-pituitary-adrenal Response to Hypoglycemia in Type 2 Diabetes by Avoiding Chronic Hypoglycemia. <i>Internal Medicine</i> , 2016, 55, 3471-3473.	0.3	2
102	The Protective Effect Of A Low-protein Diet Against Tubulo-interstitial Damage In Diabetic Kidneys. <i>Cellular & Molecular Medicine: Open Access</i> , 2016, 02, .	0.4	0
103	Epidermal growth factor receptor signaling and the progression of diabetic nephropathy. <i>Journal of Diabetes Investigation</i> , 2015, 6, 519-521.	1.1	3
104	The Relevance of the Renin-Angiotensin System in the Development of Drugs to Combat Preeclampsia. <i>International Journal of Endocrinology</i> , 2015, 2015, 1-12.	0.6	21
105	Autophagy: A Novel Therapeutic Target for Diabetic Nephropathy. <i>Diabetes and Metabolism Journal</i> , 2015, 39, 451.	1.8	84
106	1-Methylnicotinamide ameliorates lipotoxicity-induced oxidative stress and cell death in kidney proximal tubular cells. <i>Free Radical Biology and Medicine</i> , 2015, 89, 831-841.	1.3	41
107	O13. Catechol-O-methyltransferase deficiency leads to hypersensitivity on the pressor response against angiotensin II. <i>Pregnancy Hypertension</i> , 2015, 5, 212-213.	0.6	0
108	Pituitary apoplexy following gonadotropin-releasing hormone agonist administration with gonadotropin-secreting pituitary adenoma. <i>Journal of Clinical Neuroscience</i> , 2015, 22, 601-603.	0.8	23

#	ARTICLE	IF	CITATIONS
109	Anti-albuminuric effects of spironolactone in patients with type 2 diabetic nephropathy: a multicenter, randomized clinical trial. <i>Clinical and Experimental Nephrology</i> , 2015, 19, 1098-1106.	0.7	49
110	Interactions of DPP-4 and integrin $\alpha 1$ influences endothelial-to-mesenchymal transition. <i>Kidney International</i> , 2015, 88, 479-489.	2.6	127
111	Lamp-2 deficiency prevents high-fat diet-induced obese diabetes via enhancing energy expenditure. <i>Biochemical and Biophysical Research Communications</i> , 2015, 465, 249-255.	1.0	18
112	Urinary Potassium Excretion and Renal and Cardiovascular Complications in Patients with Type 2 Diabetes and Normal Renal Function. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2015, 10, 2152-2158.	2.2	68
113	Predictive Properties of Plasma Amino Acid Profile for Cardiovascular Disease in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2014, 9, e101219.	1.1	41
114	N-acetyl-seryl-aspartyl-lysyl-proline: a valuable endogenous anti-fibrotic peptide for combating kidney fibrosis in diabetes. <i>Frontiers in Pharmacology</i> , 2014, 5, 70.	1.6	26
115	Lipid mediators in diabetic nephropathy. <i>Fibrogenesis and Tissue Repair</i> , 2014, 7, 12.	3.4	54
116	Role of Nutrient-Sensing Signals in the Pathogenesis of Diabetic Nephropathy. <i>BioMed Research International</i> , 2014, 2014, 1-9.	0.9	51
117	N-acetyl-seryl-aspartyl-lysyl-proline Inhibits Diabetes-Associated Kidney Fibrosis and Endothelial-Mesenchymal Transition. <i>BioMed Research International</i> , 2014, 2014, 1-12.	0.9	73
118	A new classification of Diabetic Nephropathy 2014: a report from Joint Committee on Diabetic Nephropathy. <i>Diabetology International</i> , 2014, 5, 207-211.	0.7	10
119	Clinical therapeutic strategies for early stage of diabetic kidney disease. <i>World Journal of Diabetes</i> , 2014, 5, 342.	1.3	42
120	Cancer biology in diabetes. <i>Journal of Diabetes Investigation</i> , 2014, 5, 251-264.	1.1	25
121	Linagliptin-Mediated DPP-4 Inhibition Ameliorates Kidney Fibrosis in Streptozotocin-Induced Diabetic Mice by Inhibiting Endothelial-to-Mesenchymal Transition in a Therapeutic Regimen. <i>Diabetes</i> , 2014, 63, 2120-2131.	0.3	298
122	Interventions against nutrient-sensing pathways represent an emerging new therapeutic approach for diabetic nephropathy. <i>Clinical and Experimental Nephrology</i> , 2014, 18, 210-213.	0.7	6
123	Fatty acids are novel nutrient factors to regulate mTORC1 lysosomal localization and apoptosis in podocytes. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2014, 1842, 1097-1108.	1.8	99
124	Three ileus cases associated with the use of dipeptidyl peptidase-4 inhibitors in diabetic patients. <i>Journal of Diabetes Investigation</i> , 2013, 4, 673-675.	1.1	8
125	Role of the endothelial-to-mesenchymal transition in renal fibrosis of chronic kidney disease. <i>Clinical and Experimental Nephrology</i> , 2013, 17, 488-497.	0.7	145
126	Anti-aging molecule, Sirt1: a novel therapeutic target for diabetic nephropathy. <i>Archives of Pharmacal Research</i> , 2013, 36, 230-236.	2.7	60

#	ARTICLE	IF	CITATIONS
127	Calorie restriction in overweight males ameliorates obesity-related metabolic alterations and cellular adaptations through anti-aging effects, possibly including AMPK and SIRT1 activation. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2013, 1830, 4820-4827.	1.1	41
128	The Role of Autophagy in the Pathogenesis of Diabetic Nephropathy. <i>Journal of Diabetes Research</i> , 2013, 2013, 1-9.	1.0	64
129	MicroRNAs in Kidney Fibrosis and Diabetic Nephropathy: Roles on EMT and EndMT. <i>BioMed Research International</i> , 2013, 2013, 1-10.	0.9	104
130	Obesity-Mediated Autophagy Insufficiency Exacerbates Proteinuria-induced Tubulointerstitial Lesions. <i>Journal of the American Society of Nephrology: JASN</i> , 2013, 24, 1769-1781.	3.0	185
131	Renal Protective Effects of Resveratrol. <i>Oxidative Medicine and Cellular Longevity</i> , 2013, 2013, 1-7.	1.9	123
132	Statin Use in Patients with Diabetes and Kidney Disease: The Japanese Experience. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013, 20, 407-424.	0.9	15
133	Sirtuins as Possible Drug Targets in Type 2 Diabetes. <i>Current Drug Targets</i> , 2013, 14, 622-636.	1.0	74
134	Efficacy and tolerability of vildagliptin in type 2 diabetic patients on hemodialysis. <i>Journal of Diabetes Investigation</i> , 2012, 3, 298-301.	1.1	12
135	Role of angiotensin II-mediated AMPK inactivation on obesity-related salt-sensitive hypertension. <i>Biochemical and Biophysical Research Communications</i> , 2012, 418, 559-564.	1.0	37
136	Nutrient Sensing, Autophagy, and Diabetic Nephropathy. <i>Diabetes</i> , 2012, 61, 23-29.	0.3	141
137	SIRT1 inactivation induces inflammation through the dysregulation of autophagy in human THP-1 cells. <i>Biochemical and Biophysical Research Communications</i> , 2012, 427, 191-196.	1.0	90
138	Therapeutic management of diabetic kidney disease. <i>Journal of Diabetes Investigation</i> , 2011, 2, 248-254.	1.1	12
139	Association between single nucleotide polymorphisms within genes encoding sirtuin families and diabetic nephropathy in Japanese subjects with type 2 diabetes. <i>Clinical and Experimental Nephrology</i> , 2011, 15, 381-390.	0.7	63
140	Elevation of the antifibrotic peptide N-acetyl-seryl-aspartyl-lysyl-proline: a blood pressure-independent beneficial effect of angiotensin I-converting enzyme inhibitors. <i>Fibrogenesis and Tissue Repair</i> , 2011, 4, 25.	3.4	23
141	Resveratrol Improves Oxidative Stress and Protects Against Diabetic Nephropathy Through Normalization of Mn-SOD Dysfunction in AMPK/SIRT1-Independent Pathway. <i>Diabetes</i> , 2011, 60, 634-643.	0.3	300
142	Dietary Restriction Ameliorates Diabetic Nephropathy through Anti-Inflammatory Effects and Regulation of the Autophagy via Restoration of Sirt1 in Diabetic Wistar Fatty (<i>fa/fa</i>) Rats: A Model of Type 2 Diabetes. <i>Experimental Diabetes Research</i> , 2011, 2011, 1-11.	3.8	186
143	Rapid effects of pitavastatin on uric acid homeostasis. <i>Gout and Nucleic Acid Metabolism</i> , 2011, 35, 39-47.	0.0	0
144	Diabetic Nephropathy Remission and Regression Team Trial in Japan (DNETT-Japan): Rationale and study design. <i>Diabetes Research and Clinical Practice</i> , 2010, 87, 228-232.	1.1	10

#	ARTICLE	IF	CITATIONS
145	Calorie restriction enhances cell adaptation to hypoxia through Sirt1-dependent mitochondrial autophagy in mouse aged kidney. <i>Journal of Clinical Investigation</i> , 2010, 120, 1043-1055.	3.9	560
146	Effects of high sodium intake and diuretics on the circadian rhythm of blood pressure in type 2 diabetic patients treated with an angiotensin II receptor blocker. <i>Clinical and Experimental Nephrology</i> , 2009, 13, 300-306.	0.7	29
147	Corrigendum to "Legumain/asparaginyl endopeptidase controls extracellular matrix remodeling through the degradation of fibronectin in mouse renal proximal tubular cells" [FEBS Lett. 581 (2007) 1417-1424]. <i>FEBS Letters</i> , 2007, 581, 3579-3579.	1.3	0
148	Combinational effect of genes for the renin-angiotensin system in conferring susceptibility to diabetic nephropathy. <i>Journal of Human Genetics</i> , 2007, 52, 143-151.	1.1	35
149	Inhibition of mTOR signaling with rapamycin attenuates renal hypertrophy in the early diabetic mice. <i>Biochemical and Biophysical Research Communications</i> , 2006, 340, 296-301.	1.0	150
150	N-Acetyl-Seryl-Aspartyl-Lysyl-Proline Ameliorates the Progression of Renal Dysfunction and Fibrosis in WKY Rats with Established Anti-Glomerular Basement Membrane Nephritis. <i>Journal of the American Society of Nephrology: JASN</i> , 2006, 17, 674-685.	3.0	55
151	Genetic variations in the gene encoding TFAP2B are associated with type 2 diabetes mellitus. <i>Journal of Human Genetics</i> , 2005, 50, 283-292.	1.1	68
152	N-Acetyl-Seryl-Aspartyl-Lysyl-Proline Prevents Renal Insufficiency and Mesangial Matrix Expansion in Diabetic db/db Mice. <i>Diabetes</i> , 2005, 54, 838-845.	0.3	66
153	Levofloxacin-induced Achilles tendon rupture in a patient with systemic microscopic polyangiitis. <i>Modern Rheumatology</i> , 2005, 15, 217-219.	0.9	2
154	Successful recovery of infective endocarditis-induced rapidly progressive glomerulonephritis by steroid therapy combined with antibiotics: a case report. <i>BMC Nephrology</i> , 2004, 5, 18.	0.8	29
155	Effects of Antioxidants in Diabetes-Induced Oxidative Stress in the Glomeruli of Diabetic Rats. <i>Journal of the American Society of Nephrology: JASN</i> , 2003, 14, S250-S253.	3.0	240
156	N-Acetyl-Seryl-Aspartyl-Lysyl-Proline Inhibits TGF- β 2-Mediated Plasminogen Activator Inhibitor-1 Expression via Inhibition of Smad Pathway in Human Mesangial Cells. <i>Journal of the American Society of Nephrology: JASN</i> , 2003, 14, 863-872.	3.0	80
157	Endothelin-1 Induces Cyclooxygenase-2 Expression Via Nuclear Factor of Activated T-Cell Transcription Factor in Glomerular Mesangial Cells. <i>Journal of the American Society of Nephrology: JASN</i> , 2001, 12, 1359-1368.	3.0	61
158	Evaluation of a New Care System Provided to Diabetic Patients in the Outpatient Clinic.. <i>Internal Medicine</i> , 2000, 39, 783-787.	0.3	7
159	Amelioration of accelerated diabetic mesangial expansion by treatment with a PKC β 2 inhibitor in diabetic db/db mice, a rodent model for type 2 diabetes. <i>FASEB Journal</i> , 2000, 14, 439-447.	0.2	417
160	Insulin-like growth factor I stimulates glucose uptake and expression of glucose transporter 1 in cultured mesangial cells. <i>Clinical and Experimental Nephrology</i> , 1999, 3, 159-162.	0.7	0
161	α -tocopherol treatment prevents glomerular dysfunctions in diabetic rats through inhibition of protein kinase C β 1 diacylglycerol pathway. <i>BioFactors</i> , 1998, 7, 69-76.	2.6	39