

RÃ-guel Contente

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8482813/publications.pdf>

Version: 2024-02-01

17

papers

205

citations

1163117

8

h-index

1058476

14

g-index

18

all docs

18

docs citations

18

times ranked

228

citing authors

#	ARTICLE	IF	CITATIONS
1	Feeding ecology of the Brazilian silverside< i>Atherinella brasiliensis</i>(Atherinopsidae) in a sub-tropical estuarine ecosystem. Journal of the Marine Biological Association of the United Kingdom, 2011, 91, 1197-1205.	0.8	40
2	Fishes of Paranaguá Estuarine Complex, South West Atlantic. Biota Neotropica, 2012, 12, 226-238.	1.0	29
3	Fish assemblage structure in an estuary of the Atlantic Forest biodiversity hotspot (southern Brazil). Ichthyological Research, 2011, 58, 38-50.	0.8	26
4	Analysis of fish assemblages in sectors along a salinity gradient based on species, families and functional groups. Brazilian Journal of Oceanography, 2013, 61, 251-264.	0.6	23
5	Size-related shifts in dietary composition of< i>Centropomus parallelus</i>(Perciformes:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 Ichthyology, 2009, 25, 335-342.	0.7	20
6	Feeding ecology of the American freshwater goby < i>Ctenogobius shufeldti</i> (Gobiidae,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 542 Tg Ichthyofauna of the inner shelf of Paraná, Brazil: checklist, geographic distribution, economic importance and conservation status. Biota Neotropica, 2018, 18, .	1.6	14
7	Ichthyofauna of the inner shelf of Paraná, Brazil: checklist, geographic distribution, economic importance and conservation status. Biota Neotropica, 2018, 18, .	0.5	9
8	Checklist of the fish fauna of the Araçruz Bay, São Sebastião Channel, northern coast of São Paulo, Brazil. Check List, 2016, 12, 2004.	0.4	9
9	Occurrence of the non-indigenous <i>Omobranchus punctatus</i> (Blenniidae) on the São Paulo coast, South-Eastern Brazil. Marine Biodiversity Records, 2015, 8, .	1.2	8
10	Diet of the Atlantic rainforest killifish <i>Rivulus santensis</i> (Rivulidae, Cyprinodontiformes) in southeastern Brazil. Journal of Applied Ichthyology, 2010, 26, 930-932.	0.7	7
11	Improving the characterization of fish assemblage structure through the use of multiple sampling methods: a case study in a subtropical tidal flat ecosystem. Environmental Monitoring and Assessment, 2017, 189, 251.	2.7	6
12	Resource depletion and mechanisms for food web robustness in a Neotropical estuary. Marine and Freshwater Research, 2018, 69, 1755.	1.3	4
13	A protocol for measuring spatial variables in soft-sediment tide pools. Zoologia, 2016, 33, .	0.5	2
14	A new record of the non-native fish species <i>Butis koilomatodon</i> (Bleeker 1849) (Teleostei: Eleotridae) for southeastern Brazil. Biotemas, 2016, 29, 113.	0.1	2
15	Complementarity among otter trawl, cast net, and encircling gill net improves the characterization of a subtropical subtidal fish assemblage. Journal of Applied Ichthyology, 2019, 35, 1189-1196.	0.7	2
16	Fish assemblages on the southeastern Brazilian Bight, sampled by midwater trawl during spring and summer seasons: species composition, abundance, and environmental drivers. Fishery Bulletin, 2016, 114, 220-236.	0.2	2
17	COMPARANDO O USO DA PICARÃ‰ E TARRAFA NA CARACTERIZAÇÃO DA ESTRUTURA DA ICTIOFAUNA DO ENTREMARÃ‰S DE UMA BAÃA SUBTROPICAL. Boletim Do Instituto De Pesca, 2018, 44, 208-216.	0.5	0