

Runqiu Jiang

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8483450/runqiu-jiang-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

40
papers

2,251
citations

23
h-index

40
g-index

40
ext. papers

2,861
ext. citations

8
avg, IF

4.88
L-index

#	Paper	IF	Citations
40	Interleukin-22 promotes human hepatocellular carcinoma by activation of STAT3. <i>Hepatology</i> , 2011 , 54, 900-9	11.2	220
39	IL-17A plays a critical role in the pathogenesis of liver fibrosis through hepatic stellate cell activation. <i>Journal of Immunology</i> , 2013 , 191, 1835-44	5.3	192
38	The long noncoding RNA lnc-EGFR stimulates T-regulatory cells differentiation thus promoting hepatocellular carcinoma immune evasion. <i>Nature Communications</i> , 2017 , 8, 15129	17.4	182
37	Theabrownin from Pu-erh tea attenuates hypercholesterolemia via modulation of gut microbiota and bile acid metabolism. <i>Nature Communications</i> , 2019 , 10, 4971	17.4	171
36	HULC and Linc00152 Act as Novel Biomarkers in Predicting Diagnosis of Hepatocellular Carcinoma. <i>Cellular Physiology and Biochemistry</i> , 2015 , 37, 687-96	3.9	160
35	IL-22 is related to development of human colon cancer by activation of STAT3. <i>BMC Cancer</i> , 2013 , 13, 59	4.8	128
34	Circular RNA MAT2B Promotes Glycolysis and Malignancy of Hepatocellular Carcinoma Through the miR-338-3p/PKM2 Axis Under Hypoxic Stress. <i>Hepatology</i> , 2019 , 70, 1298-1316	11.2	124
33	Circulation long non-coding RNAs act as biomarkers for predicting tumorigenesis and metastasis in hepatocellular carcinoma. <i>Oncotarget</i> , 2015 , 6, 4505-15	3.3	120
32	LINC00152 promotes proliferation in hepatocellular carcinoma by targeting EpCAM via the mTOR signaling pathway. <i>Oncotarget</i> , 2015 , 6, 42813-24	3.3	117
31	The aberrant expression of MEG3 regulated by UHRF1 predicts the prognosis of hepatocellular carcinoma. <i>Molecular Carcinogenesis</i> , 2016 , 55, 209-19	5	104
30	miR-22 promotes HBV-related hepatocellular carcinoma development in males. <i>Clinical Cancer Research</i> , 2011 , 17, 5593-603	12.9	57
29	CD24 regulates sorafenib resistance via activating autophagy in hepatocellular carcinoma. <i>Cell Death and Disease</i> , 2018 , 9, 646	9.8	56
28	ROR β +IL-17+ neutrophils play a critical role in hepatic ischemia-reperfusion injury. <i>Journal of Molecular Cell Biology</i> , 2013 , 5, 143-6	6.3	53
27	High expression levels of IKK α and IKK β are necessary for the malignant properties of liver cancer. <i>International Journal of Cancer</i> , 2010 , 126, 1263-74	7.5	53
26	Estrogen-sensitive PTPRO expression represses hepatocellular carcinoma progression by control of STAT3. <i>Hepatology</i> , 2013 , 57, 678-88	11.2	51
25	Interleukin-33 drives hepatic fibrosis through activation of hepatic stellate cells. <i>Cellular and Molecular Immunology</i> , 2018 , 15, 388-398	15.4	50
24	Epstein-Barr virus-encoded latent membrane protein 2A promotes the epithelial-mesenchymal transition in nasopharyngeal carcinoma via metastatic tumor antigen 1 and mechanistic target of rapamycin signaling induction. <i>Journal of Virology</i> , 2014 , 88, 11872-85	6.6	35

23	IL-6 promotes PD-L1 expression in monocytes and macrophages by decreasing protein tyrosine phosphatase receptor type O expression in human hepatocellular carcinoma 2020 , 8,		31
22	Inhibition of MTA1 by ER α contributes to protection hepatocellular carcinoma from tumor proliferation and metastasis. <i>Journal of Experimental and Clinical Cancer Research</i> , 2015 , 34, 128	12.8	30
21	HIF-1 β induced expression of m6A reader YTHDF1 drives hypoxia-induced autophagy and malignancy of hepatocellular carcinoma by promoting ATG2A and ATG14 translation. <i>Signal Transduction and Targeted Therapy</i> , 2021 , 6, 76	21	28
20	Bidirectional transcription of Linc00441 and RB1 via H3K27 modification-dependent way promotes hepatocellular carcinoma. <i>Cell Death and Disease</i> , 2017 , 8, e2675	9.8	27
19	PTPRO plays a dual role in hepatic ischemia reperfusion injury through feedback activation of NF- κ B. <i>Journal of Hepatology</i> , 2014 , 60, 306-12	13.4	27
18	Kinesin family member 15 promotes cancer stem cell phenotype and malignancy via reactive oxygen species imbalance in hepatocellular carcinoma. <i>Cancer Letters</i> , 2020 , 482, 112-125	9.9	27
17	Listeria-based hepatocellular carcinoma vaccine facilitates anti-PD-1 therapy by regulating macrophage polarization. <i>Oncogene</i> , 2020 , 39, 1429-1444	9.2	22
16	Effect of Tumor Size on Cancer-Specific Survival in Small Hepatocellular Carcinoma. <i>Mayo Clinic Proceedings</i> , 2015 , 90, 1187-95	6.4	21
15	PTPRO-mediated autophagy prevents hepatosteatosis and tumorigenesis. <i>Oncotarget</i> , 2015 , 6, 9420-33	3.3	20
14	RP11-323N12.5 promotes the malignancy and immunosuppression of human gastric cancer by increasing YAP1 transcription. <i>Gastric Cancer</i> , 2021 , 24, 85-102	7.6	20
13	Chemokine CCL15 Mediates Migration of Human Bone Marrow-Derived Mesenchymal Stem Cells Toward Hepatocellular Carcinoma. <i>Stem Cells</i> , 2016 , 34, 1112-22	5.8	19
12	Aggravated Liver Injury but Attenuated Inflammation in PTPRO-Deficient Mice Following LPS/D-GalN Induced Fulminant Hepatitis. <i>Cellular Physiology and Biochemistry</i> , 2015 , 37, 214-24	3.9	14
11	Resistance to FGFR1-targeted therapy leads to autophagy via TAK1/AMPK activation in gastric cancer. <i>Gastric Cancer</i> , 2020 , 23, 988-1002	7.6	11
10	PTPROt maintains T cell immunity in the microenvironment of hepatocellular carcinoma. <i>Journal of Molecular Cell Biology</i> , 2015 , 7, 338-50	6.3	10
9	S100A4 hypomethylation affects epithelial-mesenchymal transition partially induced by LMP2A in nasopharyngeal carcinoma. <i>Molecular Carcinogenesis</i> , 2016 , 55, 1467-76	5	10
8	Survival and inflammation promotion effect of PTPRO in fulminant hepatitis is associated with NF- κ B activation. <i>Journal of Immunology</i> , 2014 , 193, 5161-70	5.3	10
7	Conjugated secondary 12 β -hydroxylated bile acids promote liver fibrogenesis. <i>EBioMedicine</i> , 2021 , 66, 103290	8.8	10
6	The attenuated hepatocellular carcinoma-specific Listeria vaccine Lmdd-MPFG prevents tumor occurrence through immune regulation of dendritic cells. <i>Oncotarget</i> , 2015 , 6, 8822-38	3.3	9

5	A bystander cell-based GM-CSF secreting vaccine synergized with a low dose of cyclophosphamide presents therapeutic immune responses against murine hepatocellular carcinoma. <i>Cellular and Molecular Immunology</i> , 2013 , 10, 349-59	15.4	9
4	Clinicopathological features and prognostic factors of young patients with surgically treated liver cancer. <i>Medicine (United States)</i> , 2015 , 94, e684	1.8	8
3	Evaluation of Epstein-Barr virus latent membrane protein 2 specific T-cell receptors driven by T-cell specific promoters using lentiviral vector. <i>Clinical and Developmental Immunology</i> , 2011 , 2011, 716926		8
2	Reply: To PMID 21674558. <i>Hepatology</i> , 2014 , 59, 1208	11.2	5
1	IL-22 Signaling in the Tumor Microenvironment. <i>Advances in Experimental Medicine and Biology</i> , 2021 , 1290, 81-88	3.6	2