

Abualbishr Alshreef

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8491588/publications.pdf>

Version: 2024-02-01

22
papers

473
citations

1162367

8
h-index

887659

17
g-index

23
all docs

23
docs citations

23
times ranked

657
citing authors

#	ARTICLE	IF	CITATIONS
1	Self-managed, computerised word finding therapy as an add-on to usual care for chronic aphasia post-stroke: An economic evaluation. <i>Clinical Rehabilitation</i> , 2021, 35, 703-717.	1.0	9
2	Cost Effectiveness of Ranibizumab vs Aflibercept vs Bevacizumab for the Treatment of Macular Oedema Due to Central Retinal Vein Occlusion: The LEAVO Study. <i>Pharmacoeconomics</i> , 2021, 39, 913-927.	1.7	6
3	Intravitreal ranibizumab versus aflibercept versus bevacizumab for macular oedema due to central retinal vein occlusion: the LEAVO non-inferiority three-arm RCT. <i>Health Technology Assessment</i> , 2021, 25, 1-196.	1.3	10
4	Increasing uptake of structured self-management education programmes for type 2 diabetes in a primary care setting: a feasibility study. <i>Pilot and Feasibility Studies</i> , 2020, 6, 71.	0.5	3
5	Mapping From Visual Acuity to EQ-5D, EQ-5D With Vision Bolt-On, and VFQ-UI in Patients With Macular Edema in the LEAVO Trial. <i>Value in Health</i> , 2020, 23, 928-935.	0.1	9
6	Computerised speech and language therapy or attention control added to usual care for people with long-term post-stroke aphasia: the Big CACTUS three-arm RCT. <i>Health Technology Assessment</i> , 2020, 24, 1-176.	1.3	24
7	Self-managed, computerised speech and language therapy for patients with chronic aphasia post-stroke compared with usual care or attention control (Big CACTUS): a multicentre, single-blinded, randomised controlled trial. <i>Lancet Neurology</i> , The, 2019, 18, 821-833.	4.9	116
8	The impact of an intervention to increase uptake to structured self-management education for people with type 2 diabetes mellitus in primary care (the embedding package), compared to usual care, on glycaemic control: study protocol for a mixed methods study incorporating a wait-list cluster randomised controlled trial. <i>BMC Family Practice</i> , 2019, 20, 152.	2.9	5
9	Statistical Methods for Adjusting Estimates of Treatment Effectiveness for Patient Nonadherence in the Context of Time-to-Event Outcomes and Health Technology Assessment: A Systematic Review of Methodological Papers. <i>Medical Decision Making</i> , 2019, 39, 910-925.	1.2	5
10	Cost-Effectiveness of Docetaxel and Paclitaxel for Adjuvant Treatment of Early Breast Cancer: Adaptation of a Model-Based Economic Evaluation From the United Kingdom to South Africa. <i>Value in Health Regional Issues</i> , 2019, 19, 65-74.	0.5	11
11	PIN156 INTRAVITREAL THERAPY WITH RANIBIZUMAB VS AFLIBERCEPT VS BEVACIZUMAB FOR MACULAR OEDEMA DUE TO CENTRAL RETINAL VEIN OCCLUSION: A WITHIN TRIAL COST-UTILITY ANALYSIS. <i>Value in Health</i> , 2019, 22, S664.	0.1	1
12	PSS25 MAPPING FROM VISUAL ACUITY TO UTILITY IN PATIENTS WITH MACULAR OEDEMA DUE TO CENTRAL RETINAL VEIN OCCLUSION USING DATA FROM THE LEAVO TRIAL. <i>Value in Health</i> , 2019, 22, S891.	0.1	0
13	Impact of pelvic floor muscle training on sexual function of women with urinary incontinence and a comparison of electrical stimulation versus standard treatment (IPSU trial): a randomised controlled trial. <i>Physiotherapy</i> , 2018, 104, 91-97.	0.2	22
14	Assessing the Appropriateness of Existing Model Adaptation Methods for Low and Middle-Income Countries. <i>Value in Health</i> , 2018, 21, S8.	0.1	2
15	Cost-Effectiveness of Haemorrhoidal Artery Ligation versus Rubber Band Ligation for the Treatment of Grade IIâ€“III Haemorrhoids: Analysis Using Evidence from the HubBLE Trial. <i>Pharmacoeconomics - Open</i> , 2017, 1, 175-184.	0.9	9
16	Haemorrhoidal Artery Ligation Procedure For The Treatment of Symptomatic Grade IIâ€“III Haemorrhoids: A Trial-Based And Long-Term Model-Based Economic Evaluation. <i>Value in Health</i> , 2017, 20, A666.	0.1	0
17	Conceptual Modelling for Cost-Effectiveness Analysis of Intravitreal Therapy with Ranibizumab, Aflibercept or Bevacizumab for Macular Oedema Due to Central Retinal Vein Occlusion. <i>Value in Health</i> , 2016, 19, A366.	0.1	0
18	Haemorrhoidal artery ligation versus rubber band ligation for the management of symptomatic second-degree and third-degree haemorrhoids (HubBLE): a multicentre, open-label, randomised controlled trial. <i>Lancet</i> , The, 2016, 388, 356-364.	6.3	168

#	ARTICLE	IF	CITATIONS
19	Review of Economic Submissions to NICE Medical Technologies Evaluation Programme. Applied Health Economics and Health Policy, 2016, 14, 623-634.	1.0	5
20	The HubBLLe Trial: haemorrhoidal artery ligation (HAL) versus rubber band ligation (RBL) for symptomatic second- and third-degree haemorrhoids: a multicentre randomised controlled trial and health-economic evaluation. Health Technology Assessment, 2016, 20, 1-150.	1.3	58
21	Methods to Account for Follow-Up Time Differences when Calculating Qalys from Randomised Controlled Trials. Value in Health, 2015, 18, A711.	0.1	0
22	Provider Payment Mechanisms: Effective Policy Tools for Achieving Universal and Sustainable Healthcare Coverage. , 0, , .		10