

Darragh Downey

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8500331/publications.pdf>

Version: 2024-02-01

25
papers

1,319
citations

394421

19
h-index

610901

24
g-index

25
all docs

25
docs citations

25
times ranked

2256
citing authors

#	ARTICLE	IF	CITATIONS
1	Vocal brain development in infants of mothers with serious mental illness (CAPRI-Voc): study protocol. <i>BMJ Open</i> , 2022, 12, e053598.	1.9	0
2	Cognitive function after electroconvulsive therapy for depression: relationship to clinical response. <i>Psychological Medicine</i> , 2021, 51, 1647-1656.	4.5	12
3	Neural pathways of maternal responding: systematic review and meta-analysis. <i>Archives of Women's Mental Health</i> , 2019, 22, 179-187.	2.6	32
4	Frontal haemodynamic responses in depression and the effect of electroconvulsive therapy. <i>Journal of Psychopharmacology</i> , 2019, 33, 1003-1014.	4.0	8
5	Regional default mode network connectivity in major depressive disorder: modulation by acute intravenous citalopram. <i>Translational Psychiatry</i> , 2019, 9, 116.	4.8	59
6	Ketamine augmentation of electroconvulsive therapy to improve neuropsychological and clinical outcomes in depression (Ketamine-ECT): a multicentre, double-blind, randomised, parallel-group, superiority trial. <i>Lancet Psychiatry</i> , 2017, 4, 365-377.	7.4	82
7	Randomised controlled trial of ketamine augmentation of electroconvulsive therapy to improve neuropsychological and clinical outcomes in depression (Ketamine-ECT study). <i>Efficacy and Mechanism Evaluation</i> , 2017, 4, 1-112.	0.7	6
8	Comparing the actions of lanicemine and ketamine in depression: key role of the anterior cingulate. <i>European Neuropsychopharmacology</i> , 2016, 26, 994-1003.	0.7	100
9	Natural variation in maternal sensitivity is reflected in maternal brain responses to infant stimuli. <i>Behavioral Neuroscience</i> , 2016, 130, 500-510.	1.2	41
10	Study protocol for the randomised controlled trial: Ketamine augmentation of ECT to improve outcomes in depression (Ketamine-ECT study). <i>BMC Psychiatry</i> , 2015, 15, 257.	2.6	11
11	fMRI and MRS measures of neuroplasticity in the pharyngeal motor cortex. <i>NeuroImage</i> , 2015, 117, 1-10.	4.2	22
12	Neuronal Nitric Oxide Synthase (NOS1) Polymorphisms Interact with Financial Hardship to Affect Depression Risk. <i>Neuropsychopharmacology</i> , 2014, 39, 2857-2866.	5.4	26
13	TOMM40 rs2075650 May Represent a New Candidate Gene for Vulnerability to Major Depressive Disorder. <i>Neuropsychopharmacology</i> , 2014, 39, 1743-1753.	5.4	21
14	Does oxytocin modulate variation in maternal caregiving in healthy new mothers?. <i>Brain Research</i> , 2014, 1580, 143-150.	2.2	41
15	The Neural Basis of Maternal Bonding. <i>PLoS ONE</i> , 2014, 9, e88436.	2.5	50
16	Increased Amygdala Responses to Sad But Not Fearful Faces in Major Depression: Relation to Mood State and Pharmacological Treatment. <i>American Journal of Psychiatry</i> , 2012, 169, 841-850.	7.2	163
17	Genetic variants in the catecholamine methyltransferase gene are associated with impulsivity and executive function: Relevance for major depression. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2012, 159B, 928-940.	1.7	16
18	Reversed Frontotemporal Connectivity During Emotional Face Processing in Remitted Depression. <i>Biological Psychiatry</i> , 2012, 72, 604-611.	1.3	55

#	ARTICLE	IF	CITATIONS
19	The CREB1-BDNF-NTRK2 Pathway in Depression: Multiple Gene-Cognition-Environment Interactions. <i>Biological Psychiatry</i> , 2011, 69, 762-771.	1.3	142
20	The effect of acute citalopram on face emotion processing in remitted depression: A pharmacMRI study. <i>European Neuropsychopharmacology</i> , 2011, 21, 140-148.	0.7	47
21	The HTR1A and HTR1B receptor genes influence stress-related information processing. <i>European Neuropsychopharmacology</i> , 2011, 21, 129-139.	0.7	33
22	State-dependent alteration in face emotion recognition in depression. <i>British Journal of Psychiatry</i> , 2011, 198, 302-308.	2.8	111
23	Risk-Taking Behavior in a Gambling Task Associated with Variations in the Tryptophan Hydroxylase 2 Gene: Relevance to Psychiatric Disorders. <i>Neuropsychopharmacology</i> , 2010, 35, 1109-1119.	5.4	35
24	CNR1 Gene is Associated with High Neuroticism and Low Agreeableness and Interacts with Recent Negative Life Events to Predict Current Depressive Symptoms. <i>Neuropsychopharmacology</i> , 2009, 34, 2019-2027.	5.4	153
25	Variations in the cannabinoid receptor 1 gene predispose to migraine. <i>Neuroscience Letters</i> , 2009, 461, 116-120.	2.1	53