

# Serge GuyÃ©tant

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8514121/publications.pdf>

Version: 2024-02-01

13  
papers

320  
citations

1040056

9  
h-index

1125743

13  
g-index

13  
all docs

13  
docs citations

13  
times ranked

426  
citing authors

#	ARTICLE	IF	CITATIONS
1	Merkel Cell Polyomavirusâ€Negative Merkel Cell Carcinoma Originating from In Situ Squamous Cell Carcinoma: A Keratinocytic Tumor with Neuroendocrine Differentiation. <i>Journal of Investigative Dermatology</i> , 2022, 142, 516-527.	0.7	21
2	Investigation of the RB1-SOX2 axis constitutes a tool for viral status determination and diagnosis in Merkel cell carcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2022, 480, 1239-1254.	2.8	6
3	Reevaluation of GLI1 Expression in Skin Tumors. <i>American Journal of Dermatopathology</i> , 2021, 43, 759-761.	0.6	1
4	Polyomavirus-Positive Merkel Cell Carcinoma Derived from a Trichoblastoma Suggests an Epithelial Origin of this Merkel Cell Carcinoma. <i>Journal of Investigative Dermatology</i> , 2020, 140, 976-985.	0.7	32
5	Merkel Cell Polyomavirus T Antigens Induce Merkel Cell-Like Differentiation in GLI1-Expressing Epithelial Cells. <i>Cancers</i> , 2020, 12, 1989.	3.7	22
6	p40 expression in Merkel cell carcinoma. <i>Human Pathology</i> , 2020, 100, 54-55.	2.0	1
7	Merkel cell carcinoma of lymph nodes without a skin primary tumor: A potential metastatic neoplasia associated with a brisk immune response. <i>Journal of the American Academy of Dermatology</i> , 2020, 83, 1789-1792.	1.2	1
8	Histogenesis of Merkel Cell Carcinoma: A Comprehensive Review. <i>Frontiers in Oncology</i> , 2019, 9, 451.	2.8	63
9	Morphologic and immunophenotypical features distinguishing Merkel cell polyomavirus-positive and negative Merkel cell carcinoma. <i>Modern Pathology</i> , 2019, 32, 1605-1616.	5.5	37
10	Diagnostic accuracy of a panel of immunohistochemical and molecular markers to distinguish Merkel cell carcinoma from other neuroendocrine carcinomas. <i>Modern Pathology</i> , 2019, 32, 499-510.	5.5	73
11	Differentiating Merkel cell carcinoma of lymph nodes without a detectable primary skin tumor from other metastatic neuroendocrine carcinomas: The ELECTHIP criteria. <i>Journal of the American Academy of Dermatology</i> , 2018, 78, 964-972.e3.	1.2	14
12	Merkel cell carcinomas infiltrated with CD33 + myeloid cells and CD8 + T cells are associated with improved outcome. <i>Journal of the American Academy of Dermatology</i> , 2018, 78, 973-982.e8.	1.2	33
13	Detection of the Merkel cell polyomavirus in the neuroendocrine component of combined Merkel cell carcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2018, 472, 825-837.	2.8	16