Louis M Luttrell

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8524443/louis-m-luttrell-publications-by-year.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

155	18,570	65	136
papers	citations	h-index	g-index
159	19,587	7.4	6.68
ext. papers	ext. citations	avg, IF	L-index

#	Paper	IF	Citations
155	Aging-related modifications to G protein-coupled receptor signaling diversity. <i>Pharmacology & Therapeutics</i> , 2021 , 223, 107793	13.9	3
154	EArrestin2 is a critical component of the GPCR-eNOS signalosome. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 11483-11492	11.5	11
153	Reply to Schierwagen et al.: EArrestins in liver disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 27085-27086	11.5	
152	Longitudinal Plasma Kallikrein Levels and Their Association With the Risk of Cardiovascular Disease Outcomes in Type 1 Diabetes in DCCT/EDIC. <i>Diabetes</i> , 2020 , 69, 2440-2445	0.9	0
151	SnapShot: EArrestin Functions. <i>Cell</i> , 2020 , 182, 1362-1362.e1	56.2	7
150	Conformational Sensors and Domain Swapping Reveal Structural and Functional Differences between EArrestin Isoforms. <i>Cell Reports</i> , 2019 , 28, 3287-3299.e6	10.6	30
149	Probing Arrestin Function Using Intramolecular FlAsH-BRET Biosensors. <i>Methods in Molecular Biology</i> , 2019 , 1957, 309-322	1.4	2
148	Islet Harvest in Carbon Monoxide-Saturated Medium for Chronic Pancreatitis Patients Undergoing Islet Autotransplantation. <i>Cell Transplantation</i> , 2019 , 28, 25S-36S	4	8
147	Transcriptomic characterization of signaling pathways associated with osteoblastic differentiation of MC-3T3E1 cells. <i>PLoS ONE</i> , 2019 , 14, e0204197	3.7	13
146	Plasma Connective Tissue Growth Factor (CTGF/CCN2) Levels Predict Myocardial Infarction in the Veterans Affairs Diabetes Trial (VADT) Cohort. <i>Diabetes Care</i> , 2018 , 41, 840-846	14.6	9
145	GIT2-A keystone in ageing and age-related disease. <i>Ageing Research Reviews</i> , 2018 , 43, 46-63	12	13
144	Translating in vitro ligand bias into in vivo efficacy. Cellular Signalling, 2018, 41, 46-55	4.9	28
143	The RXFP3-GIT2 signaling system represents a potential multidimensional therapeutic target in age-related disorders. <i>FASEB Journal</i> , 2018 , 32, 533.111	0.9	
142	Sphingosine 1 Phosphate Regulates Store-Operated Calcium Entry through binding to STIM1. <i>FASEB Journal</i> , 2018 , 32, 815.10	0.9	
141	Ligand-specific patterns of PTH1R and arrestin3 internalization and trafficking define a novel form of ligand Bias[IFASEB Journal, 2018, 32, 685.3]	0.9	
140	Autologous Mesenchymal Stem Cell and Islet Cotransplantation: Safety and Efficacy. <i>Stem Cells Translational Medicine</i> , 2018 , 7, 11-19	6.9	32
139	EArrestin Based Receptor Signaling Paradigms: Potential Therapeutic Targets for Complex Age-Related Disorders. <i>Frontiers in Pharmacology</i> , 2018 , 9, 1369	5.6	36

(2015-2018)

138	Analysis of longitudinal semicontinuous data using marginalized two-part model. <i>Journal of Translational Medicine</i> , 2018 , 16, 301	8.5	2
137	Manifold roles of Earrestins in GPCR signaling elucidated with siRNA and CRISPR/Cas9. <i>Science Signaling</i> , 2018 , 11,	8.8	116
136	The Diverse Roles of Arrestin Scaffolds in G Protein-Coupled Receptor Signaling. <i>Pharmacological Reviews</i> , 2017 , 69, 256-297	22.5	234
135	S1P in HDL promotes interaction between SR-BI and S1PR1 and activates S1PR1-mediated biological functions: calcium flux and S1PR1 internalization. <i>Journal of Lipid Research</i> , 2017 , 58, 325-338	3 ^{6.} 3	23
134	Arrestin-Dependent ERK Activation and Its Disruption 2017, 199-217		1
133	Is Signaling Specificity Encoded in Arrestin Conformation? 2017 , 235-253		
132	Informatic deconvolution of biased GPCR signaling mechanisms from in vivo pharmacological experimentation. <i>Methods</i> , 2016 , 92, 51-63	4.6	22
131	GPCR Signaling Rides a Wave of Conformational Changes. <i>Cell</i> , 2016 , 167, 602-603	56.2	4
130	Plasma Prekallikrein Is Associated With Carotid Intima-Media Thickness in Type 1 Diabetes. <i>Diabetes</i> , 2016 , 65, 498-502	0.9	11
129	The conformational signature of Earrestin2 predicts its trafficking and signalling functions. <i>Nature</i> , 2016 , 531, 665-8	50.4	144
128	Multivariate Generalized Linear Mixed Models With Random Intercepts To Analyze Cardiovascular Risk Markers in Type-1 Diabetic Patients. <i>Journal of Applied Statistics</i> , 2016 , 43, 1447-1464	1	7
127	Exploring G protein-coupled receptor signaling networks using SILAC-based phosphoproteomics. <i>Methods</i> , 2016 , 92, 36-50	4.6	19
126	Angiotensin II receptors and peritoneal dialysis-induced peritoneal fibrosis. <i>International Journal of Biochemistry and Cell Biology</i> , 2016 , 77, 240-50	5.6	7
125	Hyperparathyroidism-jaw Tumor Syndrome: An Overlooked Cause of Severe Hypercalcemia. <i>American Journal of the Medical Sciences</i> , 2016 , 352, 302-5	2.2	9
124	Delineation of a conserved arrestin-biased signaling repertoire in vivo. <i>Molecular Pharmacology</i> , 2015 , 87, 706-17	4.3	31
123	Inhibition of Sphingosine Kinase 1 Ameliorates Angiotensin II-Induced Hypertension and Inhibits Transmembrane Calcium Entry via Store-Operated Calcium Channel. <i>Molecular Endocrinology</i> , 2015 , 29, 896-908		20
122	Fulfilling the Promise of "Biased" G Protein-Coupled Receptor Agonism. <i>Molecular Pharmacology</i> , 2015 , 88, 579-88	4.3	153
121	Sphingosine Kinase 1 Mediates Transmembrane Calcium Entry via Store-Operated Calcium Channel. <i>FASEB Journal</i> , 2015 , 29, 715.34	0.9	

120	Stimulation of Cyclooxygenase 2 Expression in Rat Peritoneal Mesothelial Cells. <i>Nephron Experimental Nephrology</i> , 2014 ,		1
119	Epidermal growth factor-induced proliferation of collecting duct cells from Oak Ridge polycystic kidney mice involves activation of Na+/H+ exchanger. <i>American Journal of Physiology - Cell Physiology</i> , 2014 , 307, C554-60	5.4	9
118	Arrestin-dependent activation of ERK and Src family kinases. <i>Handbook of Experimental Pharmacology</i> , 2014 , 219, 225-57	3.2	27
117	Arrestin-dependent angiotensin AT1 receptor signaling regulates Akt and mTor-mediated protein synthesis. <i>Journal of Biological Chemistry</i> , 2014 , 289, 26155-26166	5.4	35
116	Minireview: More than just a hammer: ligand "bias" and pharmaceutical discovery. <i>Molecular Endocrinology</i> , 2014 , 28, 281-94		99
115	Biasing the parathyroid hormone receptor: relating in vitro ligand efficacy to in vivo biological activity. <i>Methods in Enzymology</i> , 2013 , 522, 229-62	1.7	20
114	The arrestin-selective angiotensin AT1 receptor agonist [Sar1,Ile4,Ile8]-AngII negatively regulates bradykinin B2 receptor signaling via AT1-B2 receptor heterodimers. <i>Journal of Biological Chemistry</i> , 2013 , 288, 18872-84	5.4	39
113	Relationship between vitamin D status and incidence of vascular events in the Veterans Affairs Diabetes Trial. <i>Atherosclerosis</i> , 2013 , 228, 502-7	3.1	19
112	Emergent biological properties of arrestin pathway-selective biased agonism. <i>Journal of Receptor and Signal Transduction Research</i> , 2013 , 33, 153-61	2.6	11
111	Earrestin-selective G protein-coupled receptor agonists engender unique biological efficacy in vivo. <i>Molecular Endocrinology</i> , 2013 , 27, 296-314		54
110	Angiotensin II activates NF-B through AT1A receptor recruitment of Earrestin in cultured rat vascular smooth muscle cells. <i>American Journal of Physiology - Cell Physiology</i> , 2013 , 304, C1176-86	5.4	16
109	Arrestins as regulators of kinases and phosphatases. <i>Progress in Molecular Biology and Translational Science</i> , 2013 , 118, 115-47	4	49
108	Arrestin pathways as drug targets. <i>Progress in Molecular Biology and Translational Science</i> , 2013 , 118, 469-97	4	20
107	A high-content, live-cell, and real-time approach to the quantitation of ligand-induced EArrestin2 and Class A/Class B GPCR mobilization. <i>Microscopy and Microanalysis</i> , 2013 , 19, 150-70	0.5	10
106	Textrous!: extracting semantic textual meaning from gene sets. <i>PLoS ONE</i> , 2013 , 8, e62665	3.7	18
105	Low-density lipoprotein induced expression of connective tissue growth factor via transactivation of sphingosine 1-phosphate receptors in mesangial cells. <i>Molecular Endocrinology</i> , 2012 , 26, 833-45		18
104	Endocrine function in aging. International Journal of Endocrinology, 2012, 2012, 872478	2.7	9
103	Functional signaling biases in G protein-coupled receptors: Game Theory and receptor dynamics. Mini-Reviews in Medicinal Chemistry, 2012, 12, 831-40	3.2	33

(2009-2011)

Refining efficacy: allosterism and bias in G protein-coupled receptor signaling. <i>Methods in Molecular Biology</i> , 2011 , 756, 3-35	1.4	66
'Biasing' the parathyroid hormone receptor: a novel anabolic approach to increasing bone mass?. <i>British Journal of Pharmacology</i> , 2011 , 164, 59-67	8.6	37
Increased expression of beta-arrestin 1 and 2 in murine models of rheumatoid arthritis: isoform specific regulation of inflammation. <i>Molecular Immunology</i> , 2011 , 49, 64-74	4.3	42
Refining efficacy: exploiting functional selectivity for drug discovery. <i>Advances in Pharmacology</i> , 2011 , 62, 79-107	5.7	24
Phospholipase C and protein kinase C-I2 mediate insulin-like growth factor II-dependent sphingosine kinase 1 activation. <i>Molecular Endocrinology</i> , 2011 , 25, 2144-56		16
The beta-arrestin pathway-selective type 1A angiotensin receptor (AT1A) agonist [Sar1,Ile4,Ile8]angiotensin II regulates a robust G protein-independent signaling network. <i>Journal of Biological Chemistry</i> , 2011 , 286, 19880-91	5.4	54
Beta-arrestin 2 negatively regulates sepsis-induced inflammation. <i>Immunology</i> , 2010 , 130, 344-51	7.8	60
HDL3, but not HDL2, stimulates plasminogen activator inhibitor-1 release from adipocytes: the role of sphingosine-1-phosphate. <i>Journal of Lipid Research</i> , 2010 , 51, 2619-28	6.3	42
Plasma kallikrein promotes epidermal growth factor receptor transactivation and signaling in vascular smooth muscle through direct activation of protease-activated receptors. <i>Journal of Biological Chemistry</i> , 2010 , 285, 35206-15	5.4	41
Bradykinin decreases podocyte permeability through ADAM17-dependent epidermal growth factor receptor activation and zonula occludens-1 rearrangement. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2010 , 334, 775-83	4.7	24
Beyond desensitization: physiological relevance of arrestin-dependent signaling. <i>Pharmacological Reviews</i> , 2010 , 62, 305-30	22.5	315
Genetic variant in the promoter of connective tissue growth factor gene confers susceptibility to nephropathy in type 1 diabetes. <i>Journal of Medical Genetics</i> , 2010 , 47, 391-7	5.8	21
Novel mechanisms in the regulation of G protein-coupled receptor trafficking to the plasma membrane. <i>Journal of Biological Chemistry</i> , 2010 , 285, 33816-25	5.4	10
Allosteric modulators of g protein-coupled receptors: future therapeutics for complex physiological disorders. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2009 , 331, 340-8	4.7	77
Angiotensin II-induced cyclooxygenase 2 expression in rat aorta vascular smooth muscle cells does not require heterotrimeric G protein activation. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2009 , 330, 118-24	4.7	12
c-Src-mediated phosphorylation of AP-2 reveals a general mechanism for receptors internalizing through the clathrin pathway. <i>Cellular Signalling</i> , 2009 , 21, 103-10	4.9	48
Diversity in arrestin function. Cellular and Molecular Life Sciences, 2009, 66, 2953-73	10.3	51
Essential role of c-Cbl in amphiregulin-induced recycling and signaling of the endogenous epidermal growth factor receptor. <i>Biochemistry</i> , 2009 , 48, 1462-73	3.2	39
	Biology, 2011, 756, 3-35 'Biasing' the parathyroid hormone receptor: a novel anabolic approach to increasing bone mass?. British Journal of Pharmacology, 2011, 164, 59-67 Increased expression of beta-arrestin 1 and 2 in murine models of rheumatoid arthritis: isoform specific regulation of Inflammation. Molecular Immunology, 2011, 49, 64-74 Refining efficacy: exploiting functional selectivity for drug discovery. Advances in Pharmacology, 2011, 62, 79-107 Phospholipase C and protein kinase C-12 mediate insulin-like growth factor II-dependent sphingosine kinase 1 activation. Molecular Endocrinology, 2011, 25, 2144-56 The beta-arrestin pathway-selective type 1A angiotensin receptor (A11A) agonist [Sar1,184,186] angiotensin II regulates a robust G protein-independent signaling network. Journal of Biological Chemistry, 2011, 286, 19880-91 Beta-arrestin 2 negatively regulates sepsis-induced inflammation. Immunology, 2010, 130, 344-51 HDL3, but not HDL2, stimulates plasminogen activator inhibitor-1 release from adipocytes: the role of sphingosine-1-phosphate. Journal of Lipid Research, 2010, 51, 2619-28 Plasma kallikrein promotes epidermal growth factor receptor transactivation and signaling in vascular smooth muscle through direct activation of protease-activated receptors. Journal of Biological Chemistry, 2010, 285, 35206-15 Bradykinin decreases podocyte permeability through ADAM17-dependent epidermal growth factor receptor activation and zonula octudens-1 rearrangement. Journal of Pharmacology and Experimental Therapeutics, 2010, 334, 775-83 Beyond desensitization: physiological relevance of arrestin-dependent signaling. Pharmacological Reviews, 2010, 62, 305-30 Genetic variant in the promoter of connective tissue growth factor gene confers susceptibility to nephropathy in type 1 diabetes. Journal of Medical Genetics, 2010, 47, 391-7 Novel mechanisms in the regulation of G protein-coupled receptors for complex physiological disorders. Journal of Biological Chemistry, 2010, 285, 33816-25 Allo	Biology, 2011, 756, 3-35 Biasing' the parathyroid hormone receptor: a novel anabolic approach to increasing bone mass?. Biasing' the parathyroid hormone receptor: a novel anabolic approach to increasing bone mass?. British Journal of Pharmacology, 2011, 164, 59-67 Increased expression of beta-arrestin 1 and 2 in murine models of rheumatoid arthritis: isoform specific regulation of inflammation. Molecular Immunology, 2011, 49, 64-74 Refining efficacy: exploiting functional selectivity for drug discovery. Advances in Pharmacology, 2011, 62, 79-107 Phospholipase C and protein kinase C-I2 mediate insulin-like growth factor II-dependent sphinogosine kinase 1 activation. Molecular Endocrinology, 2011, 25, 2144-56 The beta-arrestin pathway-selective type 1A angiotensin receptor (AT1A) agonist [Sar1, Ile4, Ile8] angiotensin II regulates a robust G protein-independent signaling network. Journal of Biological Chemistry, 2011, 286, 19880-91 Beta-arrestin 2 negatively regulates sepsis-induced inflammation. Immunology, 2010, 130, 344-51 7.8 HDL3, but not HDL2, stimulates plasminogen activator inhibitor-1 release from adipocytes: the role of sphingosine-1-phosphate. Journal of Lipid Research, 2010, 51, 2619-28 Plasma kallikrein promotes epidermal growth factor receptor transactivation and signalling in vascular smooth muscle through direct activation of protease-activated receptors. Journal of Biological Chemistry, 2010, 285, 35206-15 Bradykinin decreases podocyte permeability through ADAM17-dependent epidermal growth factor receptor activation and zonula occludens-1 rearrangement. Journal of Pharmacology and Experimental Therapeutics, 2010, 334, 775-83 Beyond desensitization: physiological relevance of arrestin-dependent signaling. Pharmacological Reviews, 2010, 62, 305-30 Genetic variant in the promoter of connective tissue growth factor gene confers susceptibility to rephropathy in type 1 diabetes. Journal of Medical Genetics, 2010, 47, 391-7 Novel mechanisms in the regulation of G protein-coupled receptors

84	A beta-arrestin-biased agonist of the parathyroid hormone receptor (PTH1R) promotes bone formation independent of G protein activation. <i>Science Translational Medicine</i> , 2009 , 1, 1ra1	17.5	165
83	Insulin-like growth factor-2/mannose-6 phosphate receptors. <i>Vitamins and Hormones</i> , 2009 , 80, 667-97	2.5	62
82	Biased agonism reveals new G protein-independent AT1a receptor signals. FASEB Journal, 2009, 23, 880	0.5 .9	
81	Heptahelical terpsichory. Who calls the tune?. <i>Journal of Receptor and Signal Transduction Research</i> , 2008 , 28, 39-58	2.6	19
80	The adiponectin receptors AdipoR1 and AdipoR2 activate ERK1/2 through a Src/Ras-dependent pathway and stimulate cell growth. <i>Biochemistry</i> , 2008 , 47, 11682-92	3.2	90
79	Role of beta-arrestin-mediated desensitization and signaling in the control of angiotensin AT1a receptor-stimulated transcription. <i>Journal of Biological Chemistry</i> , 2008 , 283, 2088-97	5.4	53
78	Connective tissue growth factor and susceptibility to renal and vascular disease risk in type 1 diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 1893-900	5.6	50
77	Reviews in molecular biology and biotechnology: transmembrane signaling by G protein-coupled receptors. <i>Molecular Biotechnology</i> , 2008 , 39, 239-64	3	105
76	5-HT induces threonine (T375) phosphorylation of ADAM17/TACE cytoplasmic tail. <i>FASEB Journal</i> , 2008 , 22, 829.7	0.9	
75	Identification of a putative nuclear localization sequence within ANG II AT(1A) receptor associated with nuclear activation. <i>American Journal of Physiology - Cell Physiology</i> , 2007 , 292, C1398-408	5.4	40
74	G protein-coupled receptor signaling complexity in neuronal tissue: implications for novel therapeutics. <i>Current Alzheimer Research</i> , 2007 , 4, 3-19	3	41
73	Ubiquitination of beta-arrestin links seven-transmembrane receptor endocytosis and ERK activation. <i>Journal of Biological Chemistry</i> , 2007 , 282, 29549-62	5.4	109
72	The insulin-like growth factor type 1 and insulin-like growth factor type 2/mannose-6-phosphate receptors independently regulate ERK1/2 activity in HEK293 cells. <i>Journal of Biological Chemistry</i> , 2007 , 282, 26150-7	5.4	44
71	Beta-arrestins 1 and 2 differentially regulate LPS-induced signaling and pro-inflammatory gene expression. <i>Molecular Immunology</i> , 2007 , 44, 3092-9	4.3	73
70	Transmembrane signaling by G protein-coupled receptors. <i>Methods in Molecular Biology</i> , 2006 , 332, 3-4	91.4	86
69	Constitutive ERK1/2 activation by a chimeric neurokinin 1 receptor-beta-arrestin1 fusion protein. Probing the composition and function of the G protein-coupled receptor "signalsome". <i>Journal of Biological Chemistry</i> , 2006 , 281, 19346-57	5.4	32
68	Arrestin-mediated ERK activation by gonadotropin-releasing hormone receptors: receptor-specific activation mechanisms and compartmentalization. <i>Journal of Biological Chemistry</i> , 2006 , 281, 2701-10	5.4	53
67	5-HT2A receptor induces ERK phosphorylation and proliferation through ADAM-17 tumor necrosis factor-alpha-converting enzyme (TACE) activation and heparin-bound epidermal growth factor-like growth factor (HB-EGE) shedding in mesangial cells. Journal of Biological Chemistry 2006, 281, 21004-2	5.4 1012	86

(2003-2006)

66	Distinct beta-arrestin- and G protein-dependent pathways for parathyroid hormone receptor-stimulated ERK1/2 activation. <i>Journal of Biological Chemistry</i> , 2006 , 281, 10856-64	5.4	383
65	Insulin-like growth factors mediate heterotrimeric G protein-dependent ERK1/2 activation by transactivating sphingosine 1-phosphate receptors. <i>Journal of Biological Chemistry</i> , 2006 , 281, 31399-4	o 7 ·4	77
64	Signal switching, crosstalk, and arrestin scaffolds: novel G protein-coupled receptor signaling in cardiovascular disease. <i>Hypertension</i> , 2006 , 48, 173-9	8.5	46
63	Insulin-like Growth Factors Mediate Heterotrimeric G Protein-dependent ERK1/2 Activation by Transactivating Sphingosine 1-Phosphate Receptors. <i>Journal of Biological Chemistry</i> , 2006 , 281, 31399-	3 ∮4⁄07	13
62	Composition and function of g protein-coupled receptor signalsomes controlling mitogen-activated protein kinase activity. <i>Journal of Molecular Neuroscience</i> , 2005 , 26, 253-64	3.3	93
61	The origins of diversity and specificity in g protein-coupled receptor signaling. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2005 , 314, 485-94	4.7	161
60	beta-Arrestin 2 expression determines the transcriptional response to lysophosphatidic acid stimulation in murine embryo fibroblasts. <i>Journal of Biological Chemistry</i> , 2005 , 280, 32157-67	5.4	42
59	Beta-arrestin- and G protein receptor kinase-mediated calcium-sensing receptor desensitization. <i>Molecular Endocrinology</i> , 2005 , 19, 1078-87		70
58	Regulators of GPCR Activity 2005 , 159-198		4
57	Ectodomain shedding-dependent transactivation of epidermal growth factor receptors in response to insulin-like growth factor type I. <i>Molecular Endocrinology</i> , 2004 , 18, 2727-39		41
56	Not so strange bedfellows: G-protein-coupled receptors and Src family kinases. <i>Oncogene</i> , 2004 , 23, 79	6 9. 78	187
55	Protein kinase A and G protein-coupled receptor kinase phosphorylation mediates beta-1 adrenergic receptor endocytosis through different pathways. <i>Journal of Biological Chemistry</i> , 2003 , 278, 35403-11	5.4	124
54	The stability of the G protein-coupled receptor-beta-arrestin interaction determines the mechanism and functional consequence of ERK activation. <i>Journal of Biological Chemistry</i> , 2003 , 278, 6258-67	5.4	282
53	Transactivation of the epidermal growth factor receptor mediates parathyroid hormone and prostaglandin F2 alpha-stimulated mitogen-activated protein kinase activation in cultured transgenic murine osteoblasts. <i>Molecular Endocrinology</i> , 2003 , 17, 1607-21		43
52	Signaling in time and space: G protein-coupled receptors and mitogen-activated protein kinases. <i>Assay and Drug Development Technologies</i> , 2003 , 1, 327-38	2.1	56
51	Independent beta-arrestin 2 and G protein-mediated pathways for angiotensin II activation of extracellular signal-regulated kinases 1 and 2. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 10782-7	11.5	569
50	Dual inhibition of beta-adrenergic and angiotensin II receptors by a single antagonist: a functional role for receptor-receptor interaction in vivo. <i>Circulation</i> , 2003 , 108, 1611-8	16.7	217
49	Phosphorylation of G Proteins 2003 , 609-612		

48	Protein kinase A-mediated phosphorylation of the beta 2-adrenergic receptor regulates its coupling to Gs and Gi. Demonstration in a reconstituted system. <i>Journal of Biological Chemistry</i> , 2002 , 277, 31249-56	5.4	148
47	beta-Arrestin scaffolding of the ERK cascade enhances cytosolic ERK activity but inhibits ERK-mediated transcription following angiotensin AT1a receptor stimulation. <i>Journal of Biological Chemistry</i> , 2002 , 277, 9429-36	5.4	314
46	Selective inhibition of heterotrimeric Gs signaling. Targeting the receptor-G protein interface using a peptide minigene encoding the Galpha(s) carboxyl terminus. <i>Journal of Biological Chemistry</i> , 2002 , 277, 28631-40	5.4	28
45	Src-dependent tyrosine phosphorylation regulates dynamin self-assembly and ligand-induced endocytosis of the epidermal growth factor receptor. <i>Journal of Biological Chemistry</i> , 2002 , 277, 26642-5	5 ⁵ t ⁴	118
44	Dancing with different partners: protein kinase a phosphorylation of seven membrane-spanning receptors regulates their G protein-coupling specificity. <i>Molecular Pharmacology</i> , 2002 , 62, 971-4	4.3	148
43	Activation and targeting of mitogen-activated protein kinases by G-protein-coupled receptors. <i>Canadian Journal of Physiology and Pharmacology</i> , 2002 , 80, 375-82	2.4	98
42	Big G, little G: G proteins and actin cytoskeletal reorganization. <i>Molecular Cell</i> , 2002 , 9, 1152-4	17.6	6
41	The role of Enrestins in the termination and transduction of G-protein-coupled receptor signals. <i>Journal of Cell Science</i> , 2002 , 115, 455-465	5.3	788
40	The role of beta-arrestins in the termination and transduction of G-protein-coupled receptor signals. <i>Journal of Cell Science</i> , 2002 , 115, 455-65	5.3	729
39	New mechanisms in heptahelical receptor signaling to mitogen activated protein kinase cascades. <i>Oncogene</i> , 2001 , 20, 1532-9	9.2	367
38	Epidermal growth factor (EGF) receptor-dependent ERK activation by G protein-coupled receptors: a co-culture system for identifying intermediates upstream and downstream of heparin-binding EGF shedding. <i>Journal of Biological Chemistry</i> , 2001 , 276, 23155-60	5.4	176
37	beta -Arrestin-mediated recruitment of the Src family kinase Yes mediates endothelin-1-stimulated glucose transport. <i>Journal of Biological Chemistry</i> , 2001 , 276, 43663-7	5.4	103
36	G protein-coupled receptors desensitize and down-regulate epidermal growth factor receptors in renal mesangial cells. <i>Journal of Biological Chemistry</i> , 2001 , 276, 27335-44	5.4	41
35	beta-arrestin1 interacts with the catalytic domain of the tyrosine kinase c-SRC. Role of beta-arrestin1-dependent targeting of c-SRC in receptor endocytosis. <i>Journal of Biological Chemistry</i> , 2000 , 275, 11312-9	5.4	167
34	Direct binding of activated c-Src to the beta 3-adrenergic receptor is required for MAP kinase activation. <i>Journal of Biological Chemistry</i> , 2000 , 275, 38131-4	5.4	161
33	Platelet-derived growth factor receptor association with Na(+)/H(+) exchanger regulatory factor potentiates receptor activity. <i>Molecular and Cellular Biology</i> , 2000 , 20, 8352-63	4.8	178
32	Pasteurella multocida toxin stimulates mitogen-activated protein kinase via G(q/11)-dependent transactivation of the epidermal growth factor receptor. <i>Journal of Biological Chemistry</i> , 2000 , 275, 2239	o ⁵ 45	74
31	Role of endocytosis in the activation of the extracellular signal-regulated kinase cascade by sequestering and nonsequestering G protein-coupled receptors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> 2000 , 97, 1489-94	11.5	200

(1997-2000)

30	The beta(2)-adrenergic receptor mediates extracellular signal-regulated kinase activation via assembly of a multi-receptor complex with the epidermal growth factor receptor. <i>Journal of Biological Chemistry</i> , 2000 , 275, 9572-80	5.4	341
29	Transactivation of the EGF receptor mediates IGF-1-stimulated shc phosphorylation and ERK1/2 activation in COS-7 cells. <i>Journal of Biological Chemistry</i> , 2000 , 275, 22583-9	5.4	199
28	MITOGENIC SIGNALING IN ANDROGEN SENSITIVE AND INSENSITIVE PROSTATE CANCER CELL LINES. <i>Journal of Urology</i> , 2000 , 163, 1027-1032	2.5	63
27	MITOGENIC SIGNALING IN ANDROGEN SENSITIVE AND INSENSITIVE PROSTATE CANCER CELL LINES. <i>Journal of Urology</i> , 2000 , 1027	2.5	1
26	Feedback regulation of beta-arrestin1 function by extracellular signal-regulated kinases. <i>Journal of Biological Chemistry</i> , 1999 , 274, 15971-4	5.4	113
25	Src-mediated tyrosine phosphorylation of dynamin is required for beta2-adrenergic receptor internalization and mitogen-activated protein kinase signaling. <i>Journal of Biological Chemistry</i> , 1999 , 274, 1185-8	5.4	223
24	The beta3-adrenergic receptor activates mitogen-activated protein kinase in adipocytes through a Gi-dependent mechanism. <i>Journal of Biological Chemistry</i> , 1999 , 274, 12017-22	5.4	146
23	Serotonin 5-HT1A receptor-mediated Erk activation requires calcium/calmodulin-dependent receptor endocytosis. <i>Journal of Biological Chemistry</i> , 1999 , 274, 4749-53	5.4	123
22	Pleiotropic coupling of G protein-coupled receptors to the mitogen-activated protein kinase cascade. Role of focal adhesions and receptor tyrosine kinases. <i>Journal of Biological Chemistry</i> , 1999 , 274, 13978-84	5.4	212
21	Regulation of tyrosine kinase cascades by G-protein-coupled receptors. <i>Current Opinion in Cell Biology</i> , 1999 , 11, 177-83	9	608
20	Beta-arrestin-dependent formation of beta2 adrenergic receptor-Src protein kinase complexes. <i>Science</i> , 1999 , 283, 655-61	33.3	1252
19	ACTIVATION OF EXTRACELLULAR SIGNAL-REGULATED KINASE IN HUMAN PROSTATE CANCER. <i>Journal of Urology</i> , 1999 , 162, 1537-1542	2.5	105
18	ACTIVATION OF EXTRACELLULAR SIGNAL-REGULATED KINASE IN HUMAN PROSTATE CANCER. Journal of Urology, 1999 , 1537-1542	2.5	3
17	Regulation of mitogen-activated protein kinase pathways by catecholamine receptors. <i>Advances in Pharmacology</i> , 1998 , 42, 466-70	5.7	5
16	Targeting the receptor-Gq interface to inhibit in vivo pressure overload myocardial hypertrophy. <i>Science</i> , 1998 , 280, 574-7	33.3	393
15	Essential role for G protein-coupled receptor endocytosis in the activation of mitogen-activated protein kinase. <i>Journal of Biological Chemistry</i> , 1998 , 273, 685-8	5.4	439
14	G protein-coupled receptors mediate two functionally distinct pathways of tyrosine phosphorylation in rat 1a fibroblasts. Shc phosphorylation and receptor endocytosis correlate with activation of Erk kinases. <i>Journal of Biological Chemistry</i> , 1997 , 272, 31648-56	5.4	180
13	Ras-dependent mitogen-activated protein kinase activation by G protein-coupled receptors. Convergence of Gi- and Gq-mediated pathways on calcium/calmodulin, Pyk2, and Src kinase. <i>Journal of Biological Chemistry</i> , 1997 , 272, 19125-32	5.4	376

12	Gbetagamma subunits mediate Src-dependent phosphorylation of the epidermal growth factor receptor. A scaffold for G protein-coupled receptor-mediated Ras activation. <i>Journal of Biological Chemistry</i> , 1997 , 272, 4637-44	5.4	366
11	Switching of the coupling of the beta2-adrenergic receptor to different G proteins by protein kinase A. <i>Nature</i> , 1997 , 390, 88-91	50.4	1090
10	21 G-protein-coupled receptors and their regulation. <i>Advances in Second Messenger and Phosphoprotein Research</i> , 1997 , 263-277		76
9	Mitogenic signaling via G protein-coupled receptors. <i>Endocrine Reviews</i> , 1996 , 17, 698-714	27.2	373
8	Role of c-Src tyrosine kinase in G protein-coupled receptor- and Gbetagamma subunit-mediated activation of mitogen-activated protein kinases. <i>Journal of Biological Chemistry</i> , 1996 , 271, 19443-50	5.4	435
7	G(o)-protein alpha-subunits activate mitogen-activated protein kinase via a novel protein kinase C-dependent mechanism. <i>Journal of Biological Chemistry</i> , 1996 , 271, 1266-9	5.4	175
6	Phosphatidylinositol 3-kinase is an early intermediate in the G beta gamma-mediated mitogen-activated protein kinase signaling pathway. <i>Journal of Biological Chemistry</i> , 1996 , 271, 12133-6	5 5.4	316
5	Receptor-tyrosine-kinase- and G beta gamma-mediated MAP kinase activation by a common signalling pathway. <i>Nature</i> , 1995 , 376, 781-4	50.4	526
4	Distinct pathways of Gi- and Gq-mediated mitogen-activated protein kinase activation. <i>Journal of Biological Chemistry</i> , 1995 , 270, 17148-53	5.4	357
3	G beta gamma subunits mediate mitogen-activated protein kinase activation by the tyrosine kinase insulin-like growth factor 1 receptor. <i>Journal of Biological Chemistry</i> , 1995 , 270, 16495-8	5.4	170
2	Effect of cellular expression of pleckstrin homology domains on Gi-coupled receptor signaling. Journal of Biological Chemistry, 1995 , 270, 12984-9	5.4	72
1	Partial insulin resistance in the mouse BC3H-1 cell line: absent hexose-independent actions of insulin. <i>Endocrinology</i> , 1986 , 119, 331-42	4.8	8