

Hans-Peter Sinn

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8539224/hans-peter-sinn-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

248
papers

10,673
citations

49
h-index

96
g-index

310
ext. papers

12,784
ext. citations

5.1
avg, IF

5.57
L-index

#	Paper	IF	Citations
248	Identification of a population of blood circulating tumor cells from breast cancer patients that initiates metastasis in a xenograft assay. <i>Nature Biotechnology</i> , 2013 , 31, 539-44	44.5	764
247	Tumour-infiltrating lymphocytes and prognosis in different subtypes of breast cancer: a pooled analysis of 3771 patients treated with neoadjuvant therapy. <i>Lancet Oncology, The</i> , 2018 , 19, 40-50	21.7	730
246	Tumor-infiltrating lymphocytes and response to neoadjuvant chemotherapy with or without carboplatin in human epidermal growth factor receptor 2-positive and triple-negative primary breast cancers. <i>Journal of Clinical Oncology</i> , 2015 , 33, 983-91	2.2	650
245	Associations of breast cancer risk factors with tumor subtypes: a pooled analysis from the Breast Cancer Association Consortium studies. <i>Journal of the National Cancer Institute</i> , 2011 , 103, 250-63	9.7	513
244	Pathophysiologic basis of contrast enhancement in breast tumors. <i>Journal of Magnetic Resonance Imaging</i> , 1999 , 10, 260-6	5.6	323
243	CD44 variant exon epitopes in primary breast cancer and length of survival. <i>Lancet, The</i> , 1995 , 345, 615-20	4.0	321
242	International expert panel on the use of primary (preoperative) systemic treatment of operable breast cancer: review and recommendations. <i>Journal of Clinical Oncology</i> , 2003 , 21, 2600-8	2.2	276
241	A common variant at the TERT-CLPTM1L locus is associated with estrogen receptor-negative breast cancer. <i>Nature Genetics</i> , 2011 , 43, 1210-4	36.3	253
240	A Brief Overview of the WHO Classification of Breast Tumors, 4th Edition, Focusing on Issues and Updates from the 3rd Edition. <i>Breast Care</i> , 2013 , 8, 149-54	2.4	194
239	PIK3CA mutations are associated with lower rates of pathologic complete response to anti-human epidermal growth factor receptor 2 (her2) therapy in primary HER2-overexpressing breast cancer. <i>Journal of Clinical Oncology</i> , 2014 , 32, 3212-20	2.2	189
238	Novel, improved grading system(s) for IDH-mutant astrocytic gliomas. <i>Acta Neuropathologica</i> , 2018 , 136, 153-166	14.3	162
237	Standardized evaluation of tumor-infiltrating lymphocytes in breast cancer: results of the ring studies of the international immuno-oncology biomarker working group. <i>Modern Pathology</i> , 2016 , 29, 1155-64	9.8	154
236	Dose-Response Association of CD8+ Tumor-Infiltrating Lymphocytes and Survival Time in High-Grade Serous Ovarian Cancer. <i>JAMA Oncology</i> , 2017 , 3, e173290	13.4	152
235	A meta-analysis of genome-wide association studies of breast cancer identifies two novel susceptibility loci at 6q14 and 20q11. <i>Human Molecular Genetics</i> , 2012 , 21, 5373-84	5.6	143
234	Comparison of immunohistochemistry and RT-PCR for detection of CD44v-expression, a new prognostic factor in human breast cancer. <i>International Journal of Cancer</i> , 1995 , 60, 471-7	7.5	143
233	Low penetrance breast cancer susceptibility loci are associated with specific breast tumor subtypes: findings from the Breast Cancer Association Consortium. <i>Human Molecular Genetics</i> , 2011 , 20, 3289-303	5.6	140
232	Gene expression signature predicting pathologic complete response with gemcitabine, epirubicin, and docetaxel in primary breast cancer. <i>Journal of Clinical Oncology</i> , 2006 , 24, 1839-45	2.2	129

231	The histone acetyltransferase hMOF is frequently downregulated in primary breast carcinoma and medulloblastoma and constitutes a biomarker for clinical outcome in medulloblastoma. <i>International Journal of Cancer</i> , 2008 , 122, 1207-13	7.5	128
230	Prognostic value of codon 918 (ATG-->ACG) RET proto-oncogene mutations in sporadic medullary thyroid carcinoma. <i>International Journal of Cancer</i> , 2001 , 95, 62-6	7.5	106
229	Common breast cancer susceptibility loci are associated with triple-negative breast cancer. <i>Cancer Research</i> , 2011 , 71, 6240-9	10.1	100
228	Ductal epithelial proliferations of the breast: a biological continuum? Comparative genomic hybridization and high-molecular-weight cytokeratin expression patterns. <i>Journal of Pathology</i> , 2001 , 195, 415-21	9.4	98
227	Accuracy of tumor size measurement in breast cancer using MRI is influenced by histological regression induced by neoadjuvant chemotherapy. <i>European Radiology</i> , 2003 , 13, 1213-23	8	97
226	19p13.1 is a triple-negative-specific breast cancer susceptibility locus. <i>Cancer Research</i> , 2012 , 72, 1795-803	10.1	93
225	Plasma microRNA panel for minimally invasive detection of breast cancer. <i>PLoS ONE</i> , 2013 , 8, e76729	3.7	86
224	Synthetic antitumor vaccines containing MUC1 glycopeptides with two immunodominant domains-induction of a strong immune response against breast tumor tissues. <i>Angewandte Chemie - International Edition</i> , 2011 , 50, 9977-81	16.4	86
223	Clinical response after two cycles compared to HER2, Ki-67, p53, and bcl-2 in independently predicting a pathological complete response after preoperative chemotherapy in patients with operable carcinoma of the breast. <i>Breast Cancer Research</i> , 2008 , 10, R30	8.3	85
222	The role of early expression of inducible nitric oxide synthase in human breast cancer. <i>European Journal of Cancer</i> , 2005 , 41, 265-71	7.5	80
221	Aberrant cytoplasmic expression of the p16 protein in breast cancer is associated with accelerated tumour proliferation. <i>British Journal of Cancer</i> , 1998 , 78, 1661-8	8.7	79
220	Deubiquitination of EGFR by Cezanne-1 contributes to cancer progression. <i>Oncogene</i> , 2012 , 31, 4599-608	8.2	73
219	Genome-wide methylation screen in low-grade breast cancer identifies novel epigenetically altered genes as potential biomarkers for tumor diagnosis. <i>FASEB Journal</i> , 2012 , 26, 4937-50	0.9	72
218	Evaluation of neoadjuvant chemotherapeutic response of breast cancer using dynamic MRI with high temporal resolution. <i>European Radiology</i> , 2003 , 13, 80-7	8	71
217	Prognosis of breast cancer molecular subtypes in routine clinical care: A large prospective cohort study. <i>BMC Cancer</i> , 2016 , 16, 734	4.8	70
216	Safety and pharmacokinetics of bivatuzumab mertansine in patients with CD44v6-positive metastatic breast cancer: final results of a phase I study. <i>Anti-Cancer Drugs</i> , 2007 , 18, 477-85	2.4	66
215	Can Routine Imaging After Neoadjuvant Chemotherapy in Breast Cancer Predict Pathologic Complete Response?. <i>Annals of Surgical Oncology</i> , 2016 , 23, 789-95	3.1	65
214	Co-expression of MET and CD47 is a novel prognosticator for survival of luminal breast cancer patients. <i>Oncotarget</i> , 2014 , 5, 8147-60	3.3	64

213	The cancer-associated microprotein CASIMO1 controls cell proliferation and interacts with squalene epoxidase modulating lipid droplet formation. <i>Oncogene</i> , 2018 , 37, 4750-4768	9.2	64
212	Prognostic significance of tumour necrosis factor-related apoptosis-inducing ligand (TRAIL) receptor expression in patients with breast cancer. <i>Journal of Molecular Medicine</i> , 2009 , 87, 995-1007	5.5	60
211	Expression of endothelial and inducible nitric oxide synthase in benign and malignant lesions of the breast and measurement of nitric oxide using electron paramagnetic resonance spectroscopy. <i>Cancer</i> , 2002 , 95, 1191-8	6.4	59
210	Human mammary carcinomas express homologues of rat metastasis-associated variants of CD44. <i>Breast Cancer Research and Treatment</i> , 1995 , 36, 307-313	4.4	58
209	Comparison of molecular abnormalities in vulvar and vaginal melanomas. <i>Modern Pathology</i> , 2014 , 27, 1386-93	9.8	55
208	AGO Recommendations for the Diagnosis and Treatment of Patients with Early Breast Cancer: Update 2019. <i>Breast Care</i> , 2019 , 14, 224-245	2.4	53
207	Intratumoral cytokines and tumor cell biology determine spontaneous breast cancer-specific immune responses and their correlation to prognosis. <i>Cancer Research</i> , 2009 , 69, 8420-8	10.1	53
206	Tumor mutational burden and immune infiltration as independent predictors of response to neoadjuvant immune checkpoint inhibition in early TNBC in GeparNuevo. <i>Annals of Oncology</i> , 2020 , 31, 1216-1222	10.3	52
205	Breast cancer in young women (<i>International Journal of Cancer</i> , 2003 , 107, 583-92)	7.5	52
204	Mutational profiles in triple-negative breast cancer defined by ultradeep multigene sequencing show high rates of PI3K pathway alterations and clinically relevant entity subgroup specific differences. <i>Oncotarget</i> , 2014 , 5, 9952-65	3.3	52
203	Metastasis-initiating cells induce and exploit a fibroblast niche to fuel malignant colonization of the lungs. <i>Nature Communications</i> , 2020 , 11, 1494	17.4	51
202	Centrosomal aberrations in primary invasive breast cancer are associated with nodal status and hormone receptor expression. <i>International Journal of Cancer</i> , 2003 , 107, 346-52	7.5	49
201	Stress signaling in breast cancer cells induces matrix components that promote chemoresistant metastasis. <i>EMBO Molecular Medicine</i> , 2018 , 10,	12	49
200	Distribution of MED12 mutations in fibroadenomas and phyllodes tumors of the breast--implications for tumor biology and pathological diagnosis. <i>Genes Chromosomes and Cancer</i> , 2015 , 54, 444-52	5	48
199	Comparison of EndoPredict and Oncotype DX test results in hormone receptor positive invasive breast cancer. <i>PLoS ONE</i> , 2013 , 8, e58483	3.7	48
198	Interobserver agreement of proliferation index (Ki-67) outperforms mitotic count in pulmonary carcinoids. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2013 , 462, 507-13	5.1	46
197	Multigene Assays for Classification, Prognosis, and Prediction in Breast Cancer: a Critical Review on the Background and Clinical Utility. <i>Geburtshilfe Und Frauenheilkunde</i> , 2013 , 73, 932-940	2	46
196	Eliminating the breast cancer surgery paradigm after neoadjuvant systemic therapy: current evidence and future challenges. <i>Annals of Oncology</i> , 2020 , 31, 61-71	10.3	46

195	Circulating fibronectin controls tumor growth. <i>Neoplasia</i> , 2013 , 15, 925-38	6.4	45
194	Prognostic value of automated KI67 scoring in breast cancer: a centralised evaluation of 8088 patients from 10 study groups. <i>Breast Cancer Research</i> , 2016 , 18, 104	8.3	44
193	Clonality of lobular carcinoma in situ (LCIS) and metachronous invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2008 , 107, 331-5	4.4	44
192	Interdisciplinary Screening, Diagnosis, Therapy and Follow-up of Breast Cancer. Guideline of the DGGG and the DKG (S3-Level, AWMF Registry Number 032/045OL, December 2017) - Part 2 with Recommendations for the Therapy of Primary, Recurrent and Advanced Breast Cancer. <i>Geburtshilfe Und Frauenheilkunde</i> , 2018 , 78, 1056-1088	2	44
191	c-myc amplifications in primary breast carcinomas and their local recurrences. <i>Journal of Clinical Pathology</i> , 2006 , 59, 424-8	3.9	43
190	Efficient engraftment of human primary breast cancer transplants in nonconditioned NOD/Scid mice. <i>International Journal of Cancer</i> , 2003 , 105, 444-53	7.5	43
189	Comparison of immunohistochemistry with PCR for assessment of ER, PR, and Ki-67 and prediction of pathological complete response in breast cancer. <i>BMC Cancer</i> , 2017 , 17, 124	4.8	42
188	Treatment of advanced metastasized breast cancer with bone marrow-derived tumour-reactive memory T cells: a pilot clinical study. <i>Cancer Immunology, Immunotherapy</i> , 2009 , 58, 887-900	7.4	42
187	Survival and tumor characteristics of German hereditary breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2000 , 59, 185-92	4.4	42
186	The branched-chain amino acid transaminase 1 sustains growth of antiestrogen-resistant and ER-negative breast cancer. <i>Oncogene</i> , 2017 , 36, 4124-4134	9.2	41
185	The role of molecular analysis in breast cancer. <i>Pathology</i> , 2009 , 41, 77-88	1.6	41
184	Association of p16 expression with prognosis varies across ovarian carcinoma histotypes: an Ovarian Tumor Tissue Analysis consortium study. <i>Journal of Pathology: Clinical Research</i> , 2018 , 4, 250-261	5.3	38
183	Prediction of underestimated invasiveness in patients with ductal carcinoma in situ of the breast on percutaneous biopsy as rationale for recommending concurrent sentinel lymph node biopsy. <i>Breast</i> , 2013 , 22, 537-42	3.6	38
182	Invasive tubular carcinoma of the breast frequently is clonally related to flat epithelial atypia and low-grade ductal carcinoma in situ. <i>American Journal of Surgical Pathology</i> , 2009 , 33, 1646-53	6.7	38
181	Extensive and predominant in situ component in breast carcinoma: their influence on treatment results after breast-conserving therapy. <i>European Journal of Cancer</i> , 1998 , 34, 646-53	7.5	38
180	C-myc oncogene amplification in ductal carcinoma in situ of the breast. <i>Breast Cancer Research and Treatment</i> , 2002 , 74, 25-31	4.4	38
179	Prognostic gene expression signature for high-grade serous ovarian cancer. <i>Annals of Oncology</i> , 2020 , 31, 1240-1250	10.3	37
178	Can a pathological complete response of breast cancer after neoadjuvant chemotherapy be diagnosed by minimal invasive biopsy?. <i>European Journal of Cancer</i> , 2016 , 69, 142-150	7.5	37

177	Interdisciplinary consensus recommendations for the use of vacuum-assisted breast biopsy under sonographic guidance: first update 2012. <i>Ultraschall in Der Medizin</i> , 2012 , 33, 366-71	3.8	37
176	Changes in chemotherapy usage and outcome of early breast cancer patients in the last decade. <i>Breast Cancer Research and Treatment</i> , 2016 , 160, 491-499	4.4	36
175	Role of TP53 mutations in triple negative and HER2-positive breast cancer treated with neoadjuvant anthracycline/taxane-based chemotherapy. <i>Oncotarget</i> , 2016 , 7, 67686-67698	3.3	36
174	Interdisciplinary Screening, Diagnosis, Therapy and Follow-up of Breast Cancer. Guideline of the DGGG and the DKG (S3-Level, AWMF Registry Number 032/045OL, December 2017) - Part 1 with Recommendations for the Screening, Diagnosis and Therapy of Breast Cancer. <i>Geburtshilfe Und Frauenheilkunde</i> , 2018 , 78, 927-948	2	36
173	Intrinsic breast cancer subtypes defined by estrogen receptor signalling-prognostic relevance of progesterone receptor loss. <i>Modern Pathology</i> , 2013 , 26, 1161-71	9.8	35
172	Outcome analysis of patients with primary breast cancer initially treated at a certified academic breast unit. <i>Breast</i> , 2012 , 21, 303-8	3.6	33
171	Chromogranin A as tumor marker in medullary thyroid carcinoma. <i>Thyroid</i> , 1992 , 2, 5-10	6.2	33
170	Reprogramming of the ER α and ER β target gene landscape triggers tamoxifen resistance in breast cancer. <i>Cancer Research</i> , 2015 , 75, 720-31	10.1	32
169	Biological subtypes of triple-negative breast cancer are associated with distinct morphological changes and clinical behaviour. <i>Breast</i> , 2013 , 22, 986-92	3.6	31
168	Triple-Negative Breast Cancer: Clinical and Histological Correlations. <i>Breast Care</i> , 2011 , 6, 273-278	2.4	31
167	Reduction of CD44(+)/CD24(-) breast cancer cells by conventional cytotoxic chemotherapy. <i>Human Pathology</i> , 2010 , 41, 574-81	3.7	31
166	CTCF gene mutations in invasive ductal breast cancer. <i>Breast Cancer Research and Treatment</i> , 2003 , 80, 347-52	4.4	31
165	Clinical relevance of genomic aberrations in homogeneously treated high-risk stage II/III breast cancer patients. <i>International Journal of Cancer</i> , 2001 , 93, 80-4	7.5	30
164	Efficacy of nab-paclitaxel does not seem to be associated with SPARC expression in metastatic breast cancer. <i>Anticancer Research</i> , 2014 , 34, 6609-15	2.3	30
163	Breast cancers with a BRCA1-like DNA copy number profile recur less often than expected after high-dose alkylating chemotherapy. <i>Clinical Cancer Research</i> , 2015 , 21, 763-70	12.9	29
162	Obesity as risk factor for subtypes of breast cancer: results from a prospective cohort study. <i>BMC Cancer</i> , 2018 , 18, 616	4.8	29
161	Differentiation and histogenesis of syringomatous tumour of the nipple and low-grade adenosquamous carcinoma: evidence for a common origin. <i>Histopathology</i> , 2014 , 65, 9-23	7.3	29
160	Variable expression of the Fragile X Mental Retardation 1 (FMR1) gene in patients with premature ovarian failure syndrome is not dependent on number of (CGG) n triplets in exon 1. <i>Human Reproduction</i> , 2011 , 26, 1241-51	5.7	29

159	Reduced incidence of distant metastases and lower mortality in 1072 patients with breast cancer with a history of hormone replacement therapy. <i>American Journal of Obstetrics and Gynecology</i> , 2007 , 196, 342.e1-9	6.4	29
158	P53 is the strongest predictor of survival in high-risk primary breast cancer patients undergoing high-dose chemotherapy with autologous blood stem cell support. <i>International Journal of Cancer</i> , 2002 , 100, 290-6	7.5	27
157	Estrogen-related receptor alpha expression and function is associated with the transcriptional coregulator AIB1 in breast carcinoma. <i>Cancer Research</i> , 2009 , 69, 5186-93	10.1	25
156	Assessing HER2 testing quality in breast cancer: variables that influence HER2 positivity rate from a large, multicenter, observational study in Germany. <i>Modern Pathology</i> , 2017 , 30, 217-226	9.8	24
155	FGFR4 Arg388 genotype is associated with pathological complete response to neoadjuvant chemotherapy for primary breast cancer. <i>Annals of Oncology</i> , 2010 , 21, 1636-1642	10.3	24
154	AGO Recommendations for the Diagnosis and Treatment of Patients with Locally Advanced and Metastatic Breast Cancer: Update 2019. <i>Breast Care</i> , 2019 , 14, 247-255	2.4	23
153	Initial Treatment of Patients with Primary Breast Cancer: Evidence, Controversies, Consensus: Spectrum of Opinion of German Specialists at the 15th International St. Gallen Breast Cancer Conference (Vienna 2017). <i>Geburtshilfe Und Frauenheilkunde</i> , 2017 , 77, 633-644	2	23
152	Endometrial Cancer Molecular Risk Stratification is Equally Prognostic for Endometrioid Ovarian Carcinoma. <i>Clinical Cancer Research</i> , 2020 , 26, 5400-5410	12.9	23
151	Quality management and accreditation of research tissue banks: experience of the National Center for Tumor Diseases (NCT) Heidelberg. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2010 , 457, 741-7	5.1	22
150	Magnetic resonance imaging of nude mice with heterotransplanted high-grade squamous cell carcinomas: use of a low-loaded, covalently bound Gd-Hsa conjugate as contrast agent with high tumor affinity. <i>Investigative Radiology</i> , 2002 , 37, 193-8	10.1	22
149	A combination of the immunohistochemical markers CK7 and SATB2 is highly sensitive and specific for distinguishing primary ovarian mucinous tumors from colorectal and appendiceal metastases. <i>Modern Pathology</i> , 2019 , 32, 1834-1846	9.8	21
148	Development and Validation of the Gene Expression Predictor of High-grade Serous Ovarian Carcinoma Molecular SubTYPE (PrOTYPE). <i>Clinical Cancer Research</i> , 2020 , 26, 5411-5423	12.9	21
147	Complement Activation in Peritoneal Dialysis-Induced Arteriopathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2018 , 29, 268-282	12.7	21
146	RESPONDER - diagnosis of pathological complete response by vacuum-assisted biopsy after neoadjuvant chemotherapy in breast Cancer - a multicenter, confirmative, one-armed, intra-individually-controlled, open, diagnostic trial. <i>BMC Cancer</i> , 2018 , 18, 851	4.8	21
145	Early Breast Cancer Precursor Lesions: Lessons Learned from Molecular and Clinical Studies. <i>Breast Care</i> , 2010 , 5, 218-226	2.4	21
144	Role of natural killer cells in the pathogenesis of human acute graft-versus-host disease. <i>Transplantation</i> , 1993 , 56, 113-20	1.8	21
143	Summary of head-to-head comparisons of patient risk classifications by the 21-gene Recurrence Score [®] (RS) assay and other genomic assays for early breast cancer. <i>International Journal of Cancer</i> , 2019 , 145, 882-893	7.5	21
142	An international reproducibility study validating quantitative determination of ERBB2, ESR1, PGR, and MKI67 mRNA in breast cancer using MammaTyper [®] . <i>Breast Cancer Research</i> , 2017 , 19, 55	8.3	20

141	Gains of chromosome region 3q26 in intraepithelial neoplasia and invasive squamous cell carcinoma of the vulva are frequent and independent of HPV status. <i>Journal of Clinical Pathology</i> , 2008 , 61, 1034-7	3.9	20
140	Similar contributions of BRCA1 and BRCA2 germline mutations to early-onset breast cancer in Germany. <i>European Journal of Human Genetics</i> , 2003 , 11, 464-7	5.3	20
139	Gemcitabine, epirubicin and docetaxel as primary systemic therapy in patients with early breast cancer: results of a multicentre phase I/II study. <i>European Journal of Cancer</i> , 2004 , 40, 2432-8	7.5	20
138	Hyperoncotic dextran and systemic aprotinin in necrotizing rodent pancreatitis. <i>Scandinavian Journal of Gastroenterology</i> , 1995 , 30, 812-6	2.4	20
137	Nucleolar organizer regions in myogenic stromal tumours of the stomach. <i>Virchows Archiv A, Pathological Anatomy and Histopathology</i> , 1989 , 415, 317-21		20
136	Mutations in genes encoding PI3K-AKT and MAPK signaling define anogenital papillary hidradenoma. <i>Genes Chromosomes and Cancer</i> , 2016 , 55, 113-9	5	20
135	Omission of axillary dissection according to ACOSOG Z0011: impact on adjuvant treatment recommendations. <i>Annals of Surgical Oncology</i> , 2013 , 20, 1538-44	3.1	19
134	DNA methylation-based profiling of uterine neoplasms: a novel tool to improve gynecologic cancer diagnostics. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020 , 146, 97-104	4.9	19
133	AGO Recommendations for the Diagnosis and Treatment of Patients with Early Breast Cancer: Update 2021. <i>Breast Care</i> , 2021 , 16, 214-227	2.4	19
132	Deep Learning for the Classification of Small-Cell and Non-Small-Cell Lung Cancer. <i>Cancers</i> , 2020 , 12,	6.6	18
131	Do Patients After Reexcision Due to Involved or Close Margins Have the Same Risk of Local Recurrence as Those After One-Step Breast-Conserving Surgery?. <i>Annals of Surgical Oncology</i> , 2016 , 23, 1831-7	3.1	18
130	Only grading has independent impact on breast cancer survival after adjustment for pathological response to preoperative chemotherapy. <i>Anti-Cancer Drugs</i> , 2004 , 15, 127-35	2.4	18
129	CD24 Ala57Val polymorphism predicts pathologic complete response to sequential anthracycline- and taxane-based neoadjuvant chemotherapy for primary breast cancer. <i>Breast Cancer Research and Treatment</i> , 2012 , 132, 819-31	4.4	17
128	High nuclear poly(adenosine diphosphate-ribose) polymerase expression is predictive for BRCA1- and BRCA2-deficient breast cancer. <i>Journal of Clinical Oncology</i> , 2011 , 29, 4586-8; author reply 4588	2.2	17
127	Clinical and pathological associations of PTEN expression in ovarian cancer: a multicentre study from the Ovarian Tumour Tissue Analysis Consortium. <i>British Journal of Cancer</i> , 2020 , 123, 793-802	8.7	16
126	High-throughput automated scoring of Ki67 in breast cancer tissue microarrays from the Breast Cancer Association Consortium. <i>Journal of Pathology: Clinical Research</i> , 2016 , 2, 138-53	5.3	16
125	Reverse phase protein array based tumor profiling identifies a biomarker signature for risk classification of hormone receptor-positive breast cancer. <i>Translational Proteomics</i> , 2014 , 2, 52-59		16
124	Bortezomib sensitizes primary meningioma cells to TRAIL-induced apoptosis by enhancing formation of the death-inducing signaling complex. <i>Journal of Neuropathology and Experimental Neurology</i> , 2014 , 73, 1034-46	3.1	16

123	Anti-p53 in breast cancer: concordance of different assay procedures and association with p53 antigen expression. <i>Oncology</i> , 2002 , 63, 297-305	3.6	16
122	Intraoperative venous blood sampling to localize a small androgen-producing ovarian tumor. <i>Gynecological Endocrinology</i> , 2005 , 21, 138-41	2.4	15
121	MyD88 and TLR4 Expression in Epithelial Ovarian Cancer. <i>Mayo Clinic Proceedings</i> , 2018 , 93, 307-320	6.4	14
120	Clinical relevance and concordance of HER2 status in local and central testing-an analysis of 1581 HER2-positive breast carcinomas over 12 years. <i>Modern Pathology</i> , 2018 , 31, 607-615	9.8	14
119	Disseminated Tumor Cells in the Bone Marrow of Patients with Operable Primary Breast Cancer: Prognostic Impact in Immunophenotypic Subgroups and Clinical Implication for Bisphosphonate Treatment. <i>Annals of Surgical Oncology</i> , 2016 , 23, 757-66	3.1	14
118	Targeted next-generation sequencing enables reliable detection of HER2 (ERBB2) status in breast cancer and provides ancillary information of clinical relevance. <i>Genes Chromosomes and Cancer</i> , 2017 , 56, 255-265	5	14
117	Invasive ductal breast cancer within a malignant phyllodes tumor: case report and assessment of clonality. <i>Human Pathology</i> , 2010 , 41, 293-6	3.7	14
116	A randomized phase II trial of doxorubicin plus pemetrexed followed by docetaxel versus doxorubicin plus cyclophosphamide followed by docetaxel as neoadjuvant treatment of early breast cancer. <i>Annals of Oncology</i> , 2011 , 22, 609-617	10.3	14
115	S100P and HYAL2 as prognostic markers for patients with triple-negative breast cancer. <i>Experimental and Molecular Pathology</i> , 2015 , 99, 180-7	4.4	13
114	Pre-diagnostic smoking behaviour and poorer prognosis in a German breast cancer patient cohort - Differential effects by tumour subtype, NAT2 status, BMI and alcohol intake. <i>Cancer Epidemiology</i> , 2014 , 38, 419-26	2.8	13
113	AGO Recommendations for the Diagnosis and Treatment of Patients with Early Breast Cancer: Update 2017. <i>Breast Care</i> , 2017 , 12, 172-183	2.4	13
112	CD44 isoforms in prognosis of breast cancer. <i>Lancet, The</i> , 1995 , 346, 502	40	13
111	Automatic tumor-stroma separation in fluorescence TMAs enables the quantitative high-throughput analysis of multiple cancer biomarkers. <i>PLoS ONE</i> , 2011 , 6, e28048	3.7	12
110	Therapy response and prognosis of patients with early breast cancer with low positivity for hormone receptors - An analysis of 2765 patients from neoadjuvant clinical trials. <i>European Journal of Cancer</i> , 2021 , 148, 159-170	7.5	12
109	Heterogeneous Responses of Axillary Lymph Node Metastases to Neoadjuvant Chemotherapy are Common and Depend on Breast Cancer Subtype. <i>Annals of Surgical Oncology</i> , 2019 , 26, 4381-4389	3.1	11
108	AGO Recommendations for the Diagnosis and Treatment of Patients with Advanced and Metastatic Breast Cancer: Update 2017. <i>Breast Care</i> , 2017 , 12, 184-191	2.4	11
107	Neoadjuvant epirubicin, gemcitabine and docetaxel for primary breast cancer: long-term survival data and major prognostic factors based on two consecutive neoadjuvant phase I/II trials. <i>International Journal of Cancer</i> , 2013 , 133, 1006-15	7.5	11
106	Prediction of the axillary lymph node status in mammary cancer on the basis of clinicopathological data and flow cytometry. <i>Medical and Biological Engineering and Computing</i> , 2004 , 42, 733-9	3.1	11

105	Dose-dense primary systemic chemotherapy with gemcitabine plus epirubicin sequentially followed by docetaxel for early breast cancer: final results of a phase I/II trial. <i>Anti-Cancer Drugs</i> , 2005 , 16, 1023-8	2.4	11
104	Immunohistochemical and cytogenetic characterization of acantholytic squamous cell carcinoma of the breast. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2005 , 446, 305-9	5.1	11
103	Tandem high-dose chemotherapy in high-risk primary breast cancer: a multivariate analysis and a matched-pair comparison with standard-dose chemotherapy. <i>Biology of Blood and Marrow Transplantation</i> , 2001 , 7, 332-42	4.7	11
102	Abstract S1-06: Increased tumor-associated lymphocytes predict benefit from addition of carboplatin to neoadjuvant therapy for triple-negative and HER2-positive early breast cancer in the GeparSixto trial (GBG 66) 2013 ,		11
101	Endometrial stromal sarcomas with BCOR-rearrangement harbor MDM2 amplifications. <i>Journal of Pathology: Clinical Research</i> , 2020 , 6, 178-184	5.3	10
100	WHO grade related expression of TRAIL-receptors and apoptosis regulators in meningioma. <i>Pathology Research and Practice</i> , 2015 , 211, 109-16	3.4	10
99	Characteristics associated with long-term progression-free survival following high-dose chemotherapy in metastatic breast cancer and influence of chemotherapy dose. <i>Annals of Oncology</i> , 2002 , 13, 679-88	10.3	10
98	Metastatic potential of small and minimally invasive breast carcinomas. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 1994 , 425, 237-41	5.1	10
97	Analysis of the Lewisx epitope in human pancreas and pancreatic adenocarcinomas. <i>International Journal of Gastrointestinal Cancer</i> , 1992 , 11, 125-35		10
96	Gonadoblastoma Y locus genes expressed in germ cells of individuals with dysgenetic gonads and a Y chromosome in their karyotypes include DDX3Y and TSPY. <i>Human Reproduction</i> , 2019 , 34, 770-779	5.7	10
95	Immune-related Gene Expression Predicts Response to Neoadjuvant Chemotherapy but not Additional Benefit from PD-L1 Inhibition in Women with Early Triple-negative Breast Cancer. <i>Clinical Cancer Research</i> , 2021 , 27, 2584-2591	12.9	10
94	Etiology of hormone receptor positive breast cancer differs by levels of histologic grade and proliferation. <i>International Journal of Cancer</i> , 2018 , 143, 746-757	7.5	9
93	Transitions between flat epithelial atypia and low-grade ductal carcinoma in situ of the breast. <i>American Journal of Surgical Pathology</i> , 2012 , 36, 1247-52	6.7	9
92	Oblimersen combined with docetaxel, adriamycin and cyclophosphamide as neo-adjuvant systemic treatment in primary breast cancer: final results of a multicentric phase I study. <i>Annals of Oncology</i> , 2008 , 19, 1698-705	10.3	9
91	A multicentre analytical comparison study of inter-reader and inter-assay agreement of four programmed death-ligand 1 immunohistochemistry assays for scoring in triple-negative breast cancer. <i>Histopathology</i> , 2021 , 78, 567-577	7.3	9
90	Identification of a characteristic vascular belt zone in human colorectal cancer. <i>PLoS ONE</i> , 2017 , 12, e0171378	3.78	8
89	IgG4-related sclerosing mastitis in a 49-year-old patient with multiple, tumor-like nodules-Diagnostic accuracy of core needle biopsy. <i>Breast Journal</i> , 2019 , 25, 1251-1253	1.2	8
88	Genistein and enterolactone in relation to Ki-67 expression and HER2 status in postmenopausal breast cancer patients. <i>Molecular Nutrition and Food Research</i> , 2017 , 61, 1700449	5.9	8

87	Effects of passive immunization against parathyroid hormone-related protein: PTHrP is the responsible factor in mediating hypercalcemia in the Walker carcinosarcoma 256 rat model. <i>Journal of Bone and Mineral Research</i> , 1995 , 10, 7-16	6.3	8
86	Simulation of heat transfer in zone plate optics irradiated by X-ray free electron laser radiation. <i>Nuclear Instruments and Methods in Physics Research, Section A: Accelerators, Spectrometers, Detectors and Associated Equipment</i> , 2010 , 621, 620-626	1.2	8
85	Stem cell dose and tumorbiologic parameters as prognostic markers for patients with metastatic breast cancer undergoing high-dose chemotherapy with autologous blood stem cell support. <i>Stem Cells</i> , 2002 , 20, 32-40	5.8	8
84	Immunohistochemical evaluation of endothelial nitric oxide synthase expression in primary breast cancer. <i>Breast</i> , 2005 , 14, 230-5	3.6	8
83	Diagnosing Pathologic Complete Response in the Breast After Neoadjuvant Systemic Treatment of Breast Cancer Patients by Minimal Invasive Biopsy: Oral Presentation at the San Antonio Breast Cancer Symposium on Friday, December 13, 2019, Program Number GS5-03. <i>Annals of Surgery</i> , 2020 , 275,	7.8	8
82	Mismatch Repair Deficiency Drives Durable Complete Remission by Targeting Programmed Death Receptor 1 in a Metastatic Luminal Breast Cancer Patient. <i>Breast Care</i> , 2019 , 14, 53-59	2.4	8
81	Predictors of Residual Tumor in Breast-Conserving Therapy. <i>Annals of Surgical Oncology</i> , 2015 , 22 Suppl 3, S451-8	3.1	7
80	Tumor biomarker conversion between primary and metastatic breast cancer: mRNA assessment and its concordance with immunohistochemistry. <i>Oncotarget</i> , 2017 , 8, 51416-51428	3.3	7
79	Extent of primary breast cancer surgery: standards and individualized concepts. <i>Breast Care</i> , 2012 , 7, 364-9	2.4	7
78	Robust gridding of TMAs after whole-slide imaging using template matching. <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2010 , 77, 1169-76	4.6	7
77	Trypsinogen activation peptides (TAP) in peritoneal fluid as predictors of late histopathologic injury in necrotizing pancreatitis of the rat. <i>Digestive Diseases and Sciences</i> , 1999 , 44, 823-9	4	7
76	Exome analysis of oncogenic pathways and tumor mutational burden (TMB) in triple-negative breast cancer (TNBC): Results of the translational biomarker program of the neoadjuvant double-blind placebo controlled GeparNuevo trial.. <i>Journal of Clinical Oncology</i> , 2019 , 37, 509-509	2.2	7
75	AGO Recommendations for the Diagnosis and Treatment of Patients with Locally Advanced and Metastatic Breast Cancer: Update 2021. <i>Breast Care</i> , 2021 , 16, 228-235	2.4	7
74	Protein phosphatase 1, regulatory subunit 15B is a survival factor for ER β -positive breast cancer. <i>International Journal of Cancer</i> , 2013 , 132, 2714-9	7.5	6
73	Onkologie. Konsensusempfehlung zu Anwendung und Indikationen der Vakuumbiopsie der Brust unter Ultraschallsicht. <i>Geburtshilfe Und Frauenheilkunde</i> , 2005 , 65, 526-529	2	6
72	Laser-induced fluorescence diagnosis and photodynamic therapy of human renal cell carcinoma. <i>Urologia Internationalis</i> , 1995 , 55, 197-201	1.9	6
71	Abstract S4-06: PIK3CA mutation predicts resistance to anti-HER2/chemotherapy in primary HER2-positive/hormone-receptor-positive breast cancer [Prospective analysis of 737 participants of the GeparSixto and GeparQuinto studies 2013 ,		6
70	Abstract GS5-03: Diagnosing residual disease and pathologic complete response after neoadjuvant chemotherapy in breast cancer patients by image-guided vacuum-assisted breast biopsy: Results of a prospective multicenter trial 2020 ,		6

69	Sustained prognostic impact of circulating tumor cell status and kinetics upon further progression of metastatic breast cancer. <i>Breast Cancer Research and Treatment</i> , 2019 , 173, 155-165	4.4	6
68	Prediction of pathological complete response in breast cancer patients during neoadjuvant chemotherapy: Is shear wave elastography a useful tool in clinical routine?. <i>European Journal of Radiology</i> , 2020 , 128, 109025	4.7	5
67	Ultra-High-Value Kurtosis Imaging for Noninvasive Tissue Characterization of Ovarian Lesions. <i>Radiology</i> , 2020 , 296, 358-369	20.5	5
66	Predictors of resectability in breast-conserving therapy. <i>Archives of Gynecology and Obstetrics</i> , 2012 , 286, 1023-31	2.5	5
65	Histological alterations of the preampullary common bile and pancreatic duct in acute biliary and nonbiliary pancreatitis. <i>Digestion</i> , 1986 , 34, 93-100	3.6	5
64	CATCH: A Prospective Precision Oncology Trial in Metastatic Breast Cancer. <i>JCO Precision Oncology</i> , 2021 , 5,	3.6	5
63	Clinicopathologic and molecular analysis of embryonal rhabdomyosarcoma of the genitourinary tract: evidence for a distinct DICER1-associated subgroup. <i>Modern Pathology</i> , 2021 , 34, 1558-1569	9.8	5
62	Molecular Subtype Conversion between Primary and Metastatic Breast Cancer Corresponding to the Dynamics of Apoptotic and Intact Circulating Tumor Cells. <i>Cancers</i> , 2019 , 11,	6.6	4
61	Determination of paraneoplastic autoimmune responses by tumor cell biology and intratumoral IFN-alpha/IL-12 in breast cancer patients. <i>Cancer Immunology, Immunotherapy</i> , 2011 , 60, 401-11	7.4	4
60	Doxorubicin/pemetrexed followed by docetaxel versus doxorubicin/ cyclophosphamide followed by docetaxel as neoadjuvant treatment for early-stage breast cancer: a randomized phase II trial. <i>Clinical Breast Cancer</i> , 2007 , 7, 555-8	3	4
59	Stage IIa cervix carcinoma with metastasis to the heart: report of a case with immunohistochemistry, flow cytometry, and virology findings. <i>Gynecologic Oncology</i> , 2000 , 76, 133-8	4.9	4
58	Efficacy of intraoperative specimen radiography as margin assessment tool in breast conserving surgery. <i>Breast Cancer Research and Treatment</i> , 2020 , 179, 425-433	4.4	4
57	Chromothripsis in Human Breast Cancer. <i>Cancer Research</i> , 2020 , 80, 4918-4931	10.1	4
56	Estrogen, progesterone, and human epidermal growth factor receptor 2 discordance between primary and metastatic breast cancer. <i>Breast Cancer Research and Treatment</i> , 2020 , 183, 137-144	4.4	3
55	Immunohistological expression of oestrogen receptor, progesterone receptor, mammaglobin, human epidermal growth factor receptor 2 and GATA-binding protein 3 in non-small-cell lung cancer. <i>Histopathology</i> , 2020 , 77, 900-914	7.3	3
54	Änderungen der Stadiengruppierung und Überlebensraten bei Anwendung der neuen TNM-Klassifikation für das Mammakarzinom. <i>Geburtshilfe Und Frauenheilkunde</i> , 2004 , 64, 962-967	2	3
53	Neoadjuvante Chemotherapie bei lokal fortgeschrittenen Mammakarzinomen: Tumorregression und perioperative Komplikationen. <i>Geburtshilfe Und Frauenheilkunde</i> , 1998 , 58, 44-50	2	3
52	Early myocardial lesions in mice with EMC virus myocarditis. Electrocardiographic and morphological findings. <i>European Heart Journal</i> , 1987 , 8, 411-415	9.5	3

51	Validated biomarker assays confirm ARID1A loss is confounded with MMR deficiency, CD8 TIL infiltration, and provides no independent prognostic value in endometriosis-associated ovarian carcinomas.. <i>Journal of Pathology</i> , 2021 ,	9.4	3
50	MGMT promoter methylation in triple negative breast cancer of the GeparSixto trial. <i>PLoS ONE</i> , 2020 , 15, e0238021	3.7	3
49	Reconstructing tumor history in breast cancer: signatures of mutational processes and response to neoadjuvant chemotherapy. <i>Annals of Oncology</i> , 2021 , 32, 500-511	10.3	3
48	CaM Kinase II- β s Required for Diabetic Hyperglycemia and Retinopathy but Not Nephropathy. <i>Diabetes</i> , 2021 , 70, 616-626	0.9	3
47	Diagnostic accuracy of axillary staging by ultrasound in early breast cancer patients. <i>European Journal of Radiology</i> , 2021 , 135, 109468	4.7	3
46	Intrinsic subtypes and risk scores in ER+/HER2-Breast Cancer: a comparison of Prosigna and OncotypeDX risk categories with Ki67. <i>Breast</i> , 2017 , 32, S105	3.6	2
45	Vacuum-Assisted Biopsy to Diagnose a Pathological Complete Response in Breast Cancer Patients After Neoadjuvant Systemic Therapy. <i>Annals of Surgery</i> , 2018 , 268, e60-e61	7.8	2
44	Ki-67 and p53 expression of the fallopian tube mucosa in breast cancer patients with hereditary risk. <i>Archives of Gynecology and Obstetrics</i> , 2014 , 289, 1079-85	2.5	2
43	Comparison of immunohistological parameters in primary breast cancers and corresponding locoregional recurrences. <i>Pathology Research and Practice</i> , 2006 , 202, 125-30	3.4	2
42	Consensus recommendations for the use of vacuum-assisted breast biopsy under sonographic guidance. <i>Gynecological Surgery</i> , 2006 , 3, 309-314	1.7	2
41	Intronic TP53 Germline Sequence Variants Modify the Risk in German Breast/Ovarian Cancer Families. <i>Hereditary Cancer in Clinical Practice</i> , 2004 , 2, 139-45	2.3	2
40	Diagnosis and Surgical Treatment of Gastric Sarcoma. <i>Oncology Research and Treatment</i> , 1994 , 17, 391-396	3.6	2
39	Abstract P3-06-08: Ki-67 mRNA as a predictor for response to neoadjuvant chemotherapy in primary breast cancer 2012 ,		2
38	MCM3 is a novel proliferation marker associated with longer survival for patients with tubo-ovarian high-grade serous carcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2021 ,	5.1	2
37	AGO Recommendations for the Surgical Therapy of the Axilla After Neoadjuvant Chemotherapy: 2021 Update. <i>Geburtshilfe Und Frauenheilkunde</i> , 2021 , 81, 1112-1120	2	2
36	Abstract PD2-07: mRNA signatures predict response to durvalumab therapy in triple negative breast cancer (TNBC) [Results of the translational biomarker programme of the neoadjuvant double-blind placebo controlled GeparNuevo trial 2019 ,		2
35	Benefits of various dextrans after delayed therapy in necrotizing pancreatitis of the rat 1996 , 22, 1207		2
34	CXCR3-expressing metastasis-initiating cells induce and exploit a fibroblast niche in the lungs to fuel metastatic colonization		2

33	Immunohistological Expression of SOX-10 in Triple-Negative Breast Cancer: A Descriptive Analysis of 113 Samples. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	2
32	The Lack of Evidence for an Association between Cancer Biomarker Conversion Patterns and CTC-Status in Patients with Metastatic Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	2
31	Statistical modelling of HER2-positivity in breast cancer: Final analyses from two large, multicentre, non-interventional studies in Germany. <i>Breast</i> , 2020 , 49, 246-253	3.6	1
30	Giant struma ovarii. <i>Archives of Gynecology and Obstetrics</i> , 2013 , 287, 399-400	2.5	1
29	Reply to L. Ozretić et al. <i>Journal of Clinical Oncology</i> , 2011 , 29, 4588-4589	2.2	1
28	Micrometastatic bone marrow cells at diagnosis have no impact on survival of primary breast cancer patients with extensive axillary lymph node involvement treated with stem cell-supported high-dose chemotherapy. <i>Annals of Oncology</i> , 2004 , 15, 1627-32	10.3	1
27	Cyclins and Breast Cancer. <i>Oncology Research and Treatment</i> , 1998 , 21, 28-34	2.8	1
26	Characterization of new monoclonal antibodies directed against normal human exocrine pancreas and pancreatic adenocarcinomas. <i>Pancreas</i> , 1993 , 8, 279-88	2.6	1
25	Assessing HER2 testing quality in breast cancer (BC): Variables that influence HER2-positivity from a large, multicenter, observational study in Germany.. <i>Journal of Clinical Oncology</i> , 2015 , 33, 11062-11062	2.2	1
24	Lean Umbilical Cord - a Case Report. <i>Geburtshilfe Und Frauenheilkunde</i> , 2016 , 76, 1186-1188	2	1
23	Renal Tubular Dysgenesis in a Case of Fetus Acardius Amorphus. <i>Case Reports in Pathology</i> , 2019 , 2019, 5416936	0.9	1
22	JUNB suppresses distant metastasis by influencing the initial metastatic stage. <i>Clinical and Experimental Metastasis</i> , 2021 , 38, 411-423	4.7	1
21	Prognostic and Immunological Significance of ARID1A Status in Endometriosis-Associated Ovarian Carcinoma		1
20	Breast cancer characteristics and surgery among women with Li-Fraumeni syndrome in Germany-A retrospective cohort study. <i>Cancer Medicine</i> , 2021 , 10, 7747-7758	4.8	1
19	uPA-PAI-1 heteromerization promotes breast cancer progression by attracting tumorigenic neutrophils. <i>EMBO Molecular Medicine</i> , 2021 , 13, e13110	12	0
18	Vacuum-Assisted Breast Biopsy After Neoadjuvant Systemic Treatment for Reliable Exclusion of Residual Cancer in Breast Cancer Patients. <i>Annals of Surgical Oncology</i> , 2021 , 1	3.1	0
17	Sex chromosome DSD individuals with mosaic 45,X0 and aberrant Y chromosomes in 46,XY cells: distinct gender phenotypes and germ cell tumour risks.. <i>Systems Biology in Reproductive Medicine</i> , 2022 , 1-11	2.9	0
16	Author's reply to: Comparing the performance of gene expression assays in breast cancer. <i>International Journal of Cancer</i> , 2019 , 145, 1163-1164	7.5	

15	Zervixkarzinom und seine Vorstufen. <i>Onkologe</i> , 2016 , 22, 737-746	0.1
14	Sonographische Beurteilung des Endometriums. <i>Gynakologische Endokrinologie</i> , 2013 , 11, 62-64	0.1
13	Molekulare Pathologie beim Mammakarzinom. <i>Onkopipeline</i> , 2009 , 2, 141-145	
12	Angioneogenese beim Mammakarzinom - Quantifizierung der GefäÙichte mittels Morphometrie als neuer Prognosefaktor. <i>Geburtshilfe Und Frauenheilkunde</i> , 1997 , 57, 321-327	2
11	Prognosefaktoren bei der Planung organübergreifender Debulkingoperation und intraoperativer Strahlentherapie bei Beckenwandrezidiven gynäkologischer Karzinome. <i>Geburtshilfe Und Frauenheilkunde</i> , 2002 , 62, 554-559	2
10	Verminderte Inzidenz von Brustkrebsmetastasen bei Patientinnen mit präoperativer Hormonersatztherapie (HRT). <i>Geburtshilfe Und Frauenheilkunde</i> , 2002 , 62, 1175-1182	2
9	Effect of N-methyl-Nnitro-N-nitrosoguanidine on carbohydrate profiles of non-metaplastic rat gastric mucosa. <i>Journal of Cancer Research and Clinical Oncology</i> , 1993 , 119, 155-9	4.9
8	A program nucleus for encoding and decoding calendar dates. <i>Computer Methods and Programs in Biomedicine</i> , 1986 , 23, 317-8	6.9
7	ASO Visual Abstract: Vacuum-Assisted Breast Biopsy After Neoadjuvant Systemic Treatment to Reliably Exclude Residual Cancer in Breast Cancer Patients. <i>Annals of Surgical Oncology</i> , 2021 , 1	3.1
6	HER2 overexpression is associated with shorter disease-free survival (DFS) following neoadjuvant systemic therapy (NST) of primary breast cancer (PBC) patients with gemcitabine (G), epirubicin (E) and docetaxel (Doc). <i>Journal of Clinical Oncology</i> , 2007 , 25, 11087-11087	2.2
5	Expression of SPARC and response to nab-paclitaxel (nab-p) in patients (pts) with metastatic breast cancer (MBC).. <i>Journal of Clinical Oncology</i> , 2014 , 32, e12009-e12009	2.2
4	Premalignant and Malignant Breast Pathology 2016 , 179-194	
3	Predictive value of ultra-high ESR1 mRNA expression in early breast cancer.. <i>Journal of Clinical Oncology</i> , 2016 , 34, e12045-e12045	2.2
2	Aus dem Tumorboard ¶ 2-jährige Patientin mit einem benignen intraduktalen Phylloidentumor. <i>Senologie - Zeitschrift Für Mammadiagnostik Und -therapie</i> , 2016 , 13, 23-25	0
1	AGO-Empfehlungen zur operativen Therapie der Axilla nach neoadjuvanter Chemotherapie: Update 2021. <i>Senologie - Zeitschrift Für Mammadiagnostik Und -therapie</i> , 2022 , 19, 56-64	0