

# Nevsun Ä°nanÄ§

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8551523/publications.pdf>

Version: 2024-02-01

50  
papers

683  
citations

623734

14  
h-index

610901

24  
g-index

51  
all docs

51  
docs citations

51  
times ranked

1113  
citing authors

#	ARTICLE	IF	CITATIONS
1	The assessment of presenteeism and activity impairment in Beh�set�'s syndrome and recurrent aphthous stomatitis: a multicentre study. <i>Rheumatology</i> , 2022, 61, 1538-1547.	1.9	6
2	Development of a new ultrasound scoring system to evaluate glandular inflammation in Sj�gren's syndrome: an OMERACT reliability exercise. <i>Rheumatology</i> , 2022, 61, 3341-3350.	1.9	16
3	Balloon Pulmonary Angioplasty in Patients With Inoperable or Recurrent/Residual Chronic Thromboembolic Pulmonary Hypertension: A Single-Centre Initial Experience. <i>Heart Lung and Circulation</i> , 2022, 31, 520-529.	0.4	6
4	Pulmonary Endarterectomy in Patients with Antiphospholipid Syndrome-Associated Chronic Thromboembolic Pulmonary Hypertension. , 2022, 26, 394-400.		3
5	Novel and known periodontal pathogens residing in gingival crevicular fluid are associated with rheumatoid arthritis. <i>Journal of Periodontology</i> , 2021, 92, 359-370.	3.4	18
6	Risk of tuberculosis is increased in Beh�set�'s disease compared to other rheumatological disorders after anti-TNF� treatments: a case series and review of the literature. <i>Turkish Journal of Medical Sciences</i> , 2021, 51, 1689-1694.	0.9	2
7	Elevated serum TREM-1 is associated with periodontitis and disease activity in rheumatoid arthritis. <i>Scientific Reports</i> , 2021, 11, 2888.	3.3	12
8	Evaluation of rheumatoid arthritis and connective tissue disease-related interstitial lung disease with pulmonary physiologic test, HRCT, and patient-based measures of dyspnea and functional disability. <i>Clinical Rheumatology</i> , 2021, 40, 3797-3805.	2.2	3
9	Post-COVID-19 syndrome in patients with primary Sj�gren's syndrome after acute SARS-CoV-2 infection. <i>Clinical and Experimental Rheumatology</i> , 2021, , .	0.8	0
10	Influence of the age at diagnosis in the disease expression of primary Sj�gren syndrome. Analysis of 12,753 patients from the Sj�gren Big Data Consortium. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 166-174.	0.8	12
11	Influence of the age at diagnosis in the disease expression of primary Sj�gren syndrome. Analysis of 12,753 patients from the Sj�gren Big Data Consortium.. <i>Clinical and Experimental Rheumatology</i> , 2021, 39 Suppl 133, 166-174.	0.8	0
12	Predictive factors for work�day loss in Beh�set's syndrome: A multi�center study. <i>International Journal of Rheumatic Diseases</i> , 2020, 23, 240-246.	1.9	7
13	EULAR/eumusc.net standards of care for rheumatoid arthritis: cross-sectional analyses of importance, level of implementation and care gaps experienced by patients and rheumatologists across 35 European countries. <i>Annals of the Rheumatic Diseases</i> , 2020, 79, 1423-1431.	0.9	11
14	Are Salivary Gland Ultrasonography Scores Associated with Salivary Flow Rates and Oral Health-related Quality of Life in Sj�gren Syndrome?. <i>Journal of Rheumatology</i> , 2020, 47, 1774-1779.	2.0	9
15	Moderation analysis exploring associations between age and mucocutaneous activity in Beh�set�'s syndrome: A multicenter study from Turkey. <i>Journal of Dermatology</i> , 2020, 47, 1403-1410.	1.2	4
16	The assessment of malnutrition and severity of gastrointestinal disease by using symptom-based questionnaires in systemic sclerosis: is it related to severe organ involvement or capillary rarefaction at microcirculation?. <i>Clinical and Experimental Rheumatology</i> , 2020, 38 Suppl 125, 127-131.	0.8	1
17	Oral health is a mediator for disease severity in patients with Beh�set�'s disease: A multiple mediation analysis study. <i>Journal of Oral Rehabilitation</i> , 2019, 46, 349-354.	3.0	14
18	Patients with rheumatoid arthritis facing sick leave or work disability meet varying regulations: a study among rheumatologists and patients from 44 European countries. <i>Annals of the Rheumatic Diseases</i> , 2019, 78, 1472-1479.	0.9	7

#	ARTICLE	IF	CITATIONS
19	Predictors and the optimal duration of sustained remission in rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2019, 38, 3033-3039.	2.2	5
20	Musculoskeletal Ultrasound in Systemic Lupus Erythematosus: Systematic Literature Review by the Lupus Task Force of the OMERACT Ultrasound Working Group. <i>Journal of Rheumatology</i> , 2019, 46, 1379-1387.	2.0	17
21	AB1242 - COLCHICINE USE DURING PREGNANCY: CASE REPORTS. , 2019, , .		2
22	AB0223 - ASSESSMENT OF DISEASE ACTIVITY IN SYSTEMIC SCLEROSIS: THE COMPARISON BETWEEN REVISED EUSTAR AND ESCSG ACTIVITY INDEXES. , 2019, , .		0
23	THU0081 - ELEVATED SERUM TREM-1 LEVELS ARE ASSOCIATED WITH DAS28 AND PERIODONTITIS IN RHEUMATOID ARTHRITIS. , 2019, , .		0
24	Does illness perception associate with disease symptoms in Behçet's disease?. <i>Internal and Emergency Medicine</i> , 2019, 14, 691-697.	2.0	4
25	Evaluation of salivary gland ultrasonography in primary Sjögren's syndrome: does it reflect clinical activity and outcome of the disease?. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 118, 140-145.	0.8	7
26	Oral ulcer activity assessment with the composite index according to different treatment modalities in Behçet's syndrome: a multicentre study. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 121, 98-104.	0.8	3
27	The assessment of contributing factors to oral ulcer presence in Behçet's disease: Dietary and non-dietary factors. <i>European Journal of Rheumatology</i> , 2018, 5, 240-243.	0.6	19
28	Intercentre variance in patient reported outcomes is lower than objective rheumatoid arthritis activity measures: a cross-sectional study. <i>Rheumatology</i> , 2017, 56, 1395-1400.	1.9	4
29	Association of ERAP1, IL23R and PTGER4 Polymorphisms with Radiographic Severity of Ankylosing Spondylitis. <i>Open Rheumatology Journal</i> , 2017, 11, 1-9.	0.2	10
30	Oral ulcer activity in Behçet's disease: Poor medication adherence is an underestimated risk factor. <i>European Journal of Rheumatology</i> , 2017, 4, 109-112.	0.6	6
31	The 2013 ACC/AHA 10-year atherosclerotic cardiovascular disease risk index is better than SCORE and QRisk II in rheumatoid arthritis: is it enough?. <i>Rheumatology</i> , 2016, 55, kev363.	1.9	16
32	Is ankle involvement underestimated in rheumatoid arthritis? Results of a multicenter ultrasound study. <i>Clinical Rheumatology</i> , 2016, 35, 2669-2678.	2.2	18
33	Response rate of initial conventional treatments, disease course, and related factors of patients with adult-onset Still's disease: Data from a large multicenter cohort. <i>Journal of Autoimmunity</i> , 2016, 69, 59-63.	6.5	62
34	Assessment of the New 2012 EULAR/ACR Clinical Classification Criteria for Polymyalgia Rheumatica: A Prospective Multicenter Study. <i>Journal of Rheumatology</i> , 2016, 43, 893-900.	2.0	36
35	Ultrasonographic Assessment of Fifth Metatarsophalangeal Joint Erosion in Rheumatoid Arthritis: Which Aspect Is Better?. <i>Ultrasound in Medicine and Biology</i> , 2016, 42, 864-869.	1.5	6
36	Biomarkers in Remission According to Different Criteria in Patients with Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2015, 42, 2066-2070.	2.0	6

#	ARTICLE	IF	CITATIONS
37	Use of musculoskeletal ultrasound in rheumatoid arthritis in Turkey among rheumatologists: A national targeted ultrasound initiative survey. <i>European Journal of Rheumatology</i> , 2015, 2, 43-44.	0.6	4
38	The Role of Depression, Anxiety, Fatigue, and Fibromyalgia on the Evaluation of the Remission Status in Patients with Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2014, 41, 1755-1760.	2.0	44
39	Unmet need in Behcet's disease: most patients in routine follow-up continue to have oral ulcers. <i>Clinical Rheumatology</i> , 2014, 33, 1773-1776.	2.2	35
40	A new Mucocutaneous Activity Index for Behcet's disease. <i>Clinical and Experimental Rheumatology</i> , 2014, 32, S80-6.	0.8	13
41	The association of fatigue, comorbidity burden, disease activity, disability and gross domestic product in patients with rheumatoid arthritis. Results from 34 countries participating in the Quest-RA program. <i>Clinical and Experimental Rheumatology</i> , 2014, 32, 869-77.	0.8	42
42	Salivary levels of HNP 1-3 are related to oral ulcer activity in Behcet's disease. <i>International Journal of Dermatology</i> , 2013, 52, 1198-1201.	1.0	11
43	Psoriasis Aotendu induit par le rituximab chez un patient atteint de polyarthrite rhumatoide: effet indesirable inattendu. <i>Revue Du Rhumatisme (Edition Francaise)</i> , 2013, 80, 530-531.	0.0	0
44	Effects of azithromycin on intracellular cytokine responses and mucocutaneous manifestations in Behcet's disease. <i>International Journal of Dermatology</i> , 2013, 52, 1561-1566.	1.0	17
45	Patient's global assessment of disease activity and patient's assessment of general health for rheumatoid arthritis activity assessment: are they equivalent?. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 1942-1949.	0.9	59
46	Salivary levels of antimicrobial peptides Hnp 1-3, LI-37 and S100 in Behcet's disease. <i>Archives of Oral Biology</i> , 2012, 57, 642-646.	1.8	30
47	A composite index for determining the impact of oral ulcer activity in Behcet's disease and recurrent aphthous stomatitis. <i>Journal of Oral Pathology and Medicine</i> , 2010, 39, 511-511.	2.7	0
48	Agreement Between Quantiferon-TB Gold Test and Tuberculin Skin Test in the Identification of Latent Tuberculosis Infection in Patients with Rheumatoid Arthritis and Ankylosing Spondylitis. <i>Journal of Rheumatology</i> , 2009, 36, 2675-2681.	2.0	37
49	Serum mannose-binding lectin levels are decreased in behcet's disease and associated with disease severity. <i>Journal of Rheumatology</i> , 2005, 32, 287-91.	2.0	34
50	Low Serum Mannose-binding Lectin Levels in Behcet's Disease. , 2003, 528, 287-289.		5