

# Andrey I Egorov

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8555390/publications.pdf>

Version: 2024-02-01

15  
papers

392  
citations

1040056

9  
h-index

996975

15  
g-index

15  
all docs

15  
docs citations

15  
times ranked

626  
citing authors

#	ARTICLE	IF	CITATIONS
1	Latent <i>Toxoplasma gondii</i> infections are associated with elevated biomarkers of inflammation and vascular injury. <i>BMC Infectious Diseases</i> , 2021, 21, 188.	2.9	12
2	A Multiplex Noninvasive Salivary Antibody Assay for SARS-CoV-2 Infection and Its Application in a Population-Based Survey by Mail. <i>Microbiology Spectrum</i> , 2021, 9, e0069321.	3.0	9
3	Recreational water exposure and waterborne infections in a prospective salivary antibody study at a Lake Michigan beach. <i>Scientific Reports</i> , 2021, 11, 20540.	3.3	2
4	Human Cytomegalovirus Infections Are Associated With Elevated Biomarkers of Vascular Injury. <i>Frontiers in Cellular and Infection Microbiology</i> , 2020, 10, 334.	3.9	7
5	Greater tree cover near residence is associated with reduced allostatic load in residents of central North Carolina. <i>Environmental Research</i> , 2020, 186, 109435.	7.5	18
6	Application of a multiplex salivary immunoassay to detect sporadic incident norovirus infections. <i>Scientific Reports</i> , 2019, 9, 19576.	3.3	13
7	Environmental risk factors for <i>Toxoplasma gondii</i> infections and the impact of latent infections on allostatic load in residents of Central North Carolina. <i>BMC Infectious Diseases</i> , 2018, 18, 421.	2.9	19
8	Asymptomatic norovirus infection associated with swimming at a tropical beach: A prospective cohort study. <i>PLoS ONE</i> , 2018, 13, e0195056.	2.5	27
9	Application of a salivary immunoassay in a prospective community study of waterborne infections. <i>Water Research</i> , 2018, 142, 289-300.	11.3	14
10	Vegetated land cover near residence is associated with reduced allostatic load and improved biomarkers of neuroendocrine, metabolic and immune functions. <i>Environmental Research</i> , 2017, 158, 508-521.	7.5	113
11	Use of Pathogen-Specific Antibody Biomarkers to Estimate Waterborne Infections in Population-Based Settings. <i>Current Environmental Health Reports</i> , 2016, 3, 322-334.	6.7	22
12	Application of salivary antibody immunoassays for the detection of incident infections with Norwalk virus in a group of volunteers. <i>Journal of Immunological Methods</i> , 2015, 424, 53-63.	1.4	27
13	Development of a multiplex microsphere immunoassay for the quantitation of salivary antibody responses to selected waterborne pathogens. <i>Journal of Immunological Methods</i> , 2011, 364, 83-93.	1.4	51
14	Mortality in Russian Penitentiaries and the General Population. <i>Journal of Public Health Policy</i> , 2005, 26, 69-74.	2.0	1
15	Daily variations in effluent water turbidity and diarrhoeal illness in a Russian city. <i>International Journal of Environmental Health Research</i> , 2003, 13, 81-94.	2.7	57