

Constantinos Giaginis

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8559383/constantinos-giaginis-publications-by-citations.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

140
papers

3,017
citations

31
h-index

45
g-index

146
ext. papers

3,439
ext. citations

3.4
avg, IF

5.4
L-index

#	Paper	IF	Citations
140	DNA repair systems as targets of cadmium toxicity. <i>Toxicology and Applied Pharmacology</i> , 2006 , 213, 282-90	4.6	171
139	Alternative measures of lipophilicity: from octanol-water partitioning to IAM retention. <i>Journal of Pharmaceutical Sciences</i> , 2008 , 97, 2984-3004	3.9	104
138	Current State of the Art in HPLC Methodology for Lipophilicity Assessment of Basic Drugs. A Review. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007 , 31, 79-96	1.3	81
137	Clinical Value of Nutritional Status in Cancer: What is its Impact and how it Affects Disease Progression and Prognosis?. <i>Nutrition and Cancer</i> , 2017 , 69, 1151-1176	2.8	77
136	MCM proteins as diagnostic and prognostic tumor markers in the clinical setting. <i>Histology and Histopathology</i> , 2010 , 25, 351-70	1.4	76
135	Current clinical status on the preventive effects of cranberry consumption against urinary tract infections. <i>Nutrition Research</i> , 2013 , 33, 595-607	4	69
134	Clinical significance of MCM-2 and MCM-5 expression in colon cancer: association with clinicopathological parameters and tumor proliferative capacity. <i>Digestive Diseases and Sciences</i> , 2009 , 54, 282-91	4	68
133	Effects of curcumin consumption on human chronic diseases: A narrative review of the most recent clinical data. <i>Phytotherapy Research</i> , 2018 , 32, 957-975	6.7	63
132	Clinical significance of HuR expression in human malignancy. <i>Medical Oncology</i> , 2014 , 31, 161	3.7	55
131	NF- κ B signaling at the crossroads of inflammation and atherogenesis: searching for new therapeutic links. <i>Expert Opinion on Therapeutic Targets</i> , 2014 , 18, 1089-101	6.4	53
130	Different retention behavior of structurally diverse basic and neutral drugs in immobilized artificial membrane and reversed-phase high performance liquid chromatography: comparison with octanol-water partitioning. <i>Journal of Chromatography A</i> , 2006 , 1116, 158-64	4.5	53
129	Heat Shock Protein-27, -60 and -90 expression in gastric cancer: association with clinicopathological variables and patient survival. <i>BMC Gastroenterology</i> , 2009 , 9, 14	3	50
128	Peroxisome proliferator-activated receptor-gamma (PPAR-gamma) ligands as potential therapeutic agents to treat arthritis. <i>Pharmacological Research</i> , 2009 , 60, 160-9	10.2	50
127	Octanol/water partitioning simulation by reversed-phase high performance liquid chromatography for structurally diverse acidic drugs: Effect of n-octanol as mobile phase additive. <i>Journal of Chromatography A</i> , 2007 , 1166, 116-25	4.5	50
126	Lipophilicity and biomimetic properties to support drug discovery. <i>Expert Opinion on Drug Discovery</i> , 2017 , 12, 885-896	6.2	48
125	Ephrin (Eph) receptor A1, A4, A5 and A7 expression in human non-small cell lung carcinoma: associations with clinicopathological parameters, tumor proliferative capacity and patientsS survival. <i>BMC Clinical Pathology</i> , 2014 , 14, 8	3	43
124	Peroxisome proliferator-activated receptors (PPARs) in the control of bone metabolism. <i>Fundamental and Clinical Pharmacology</i> , 2007 , 21, 231-44	3.1	42

123	Application of quantitative structure-activity relationships for modeling drug and chemical transport across the human placenta barrier: a multivariate data analysis approach. <i>Journal of Applied Toxicology</i> , 2009 , 29, 724-33	4.1	41
122	Epithelial-Mesenchymal Transition in Non Small-cell Lung Cancer. <i>Anticancer Research</i> , 2017 , 37, 1773-1778		41
121	Histone deacetylase inhibitors and pancreatic cancer: are there any promising clinical trials?. <i>World Journal of Gastroenterology</i> , 2013 , 19, 1173-81	5.6	41
120	Direct injection liquid chromatography/positive ion electrospray ionization mass spectrometric quantification of methotrexate, folinic acid, folic acid and ondansetron in human serum. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2009 , 877, 3850-6	3.2	39
119	Clinical significance of ephrin (eph)-A1, -A2, -a4, -a5 and -a7 receptors in pancreatic ductal adenocarcinoma. <i>Pathology and Oncology Research</i> , 2010 , 16, 267-76	2.6	39
118	Contribution to the standardization of the chromatographic conditions for the lipophilicity assessment of neutral and basic drugs. <i>Analytica Chimica Acta</i> , 2006 , 573-574, 311-8	6.6	39
117	Histone deacetylase-1 and -2 expression in mobile tongue squamous cell carcinoma: associations with clinicopathological parameters and patients survival. <i>Journal of Oral Pathology and Medicine</i> , 2011 , 40, 706-14	3.3	37
116	MCM-2 and MCM-5 expression in gastric adenocarcinoma: clinical significance and comparison with Ki-67 proliferative marker. <i>Digestive Diseases and Sciences</i> , 2011 , 56, 777-85	4	37
115	Peroxisome proliferator-activated receptor-gamma (PPAR-gamma) ligands: novel pharmacological agents in the treatment of ischemia reperfusion injury. <i>Current Molecular Medicine</i> , 2008 , 8, 562-79	2.5	36
114	Electrostatic interactions and ionization effect in immobilized artificial membrane retention. A comparative study with octanol-water partitioning. <i>Journal of Chromatography A</i> , 2008 , 1187, 67-78	4.5	36
113	Expression of peroxisome proliferator-activated receptor-gamma in colon cancer: correlation with histopathological parameters, cell cycle-related molecules, and patients survival. <i>Digestive Diseases and Sciences</i> , 2007 , 52, 2305-11	4	35
112	Current evidence on the anticancer potential of Chios mastic gum. <i>Nutrition and Cancer</i> , 2011 , 63, 1174-84	2.4	34
111	Current toxicological aspects on drug and chemical transport and metabolism across the human placental barrier. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2012 , 8, 1263-75	5.5	31
110	The role of peroxisome proliferator-activated receptor- γ in breast cancer. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2012 , 12, 1025-44	2.2	31
109	Current evidence for histone deacetylase inhibitors in pancreatic cancer. <i>World Journal of Gastroenterology</i> , 2013 , 19, 813-28	5.6	31
108	Clinical significance of histone deacetylase (HDAC)-1, HDAC-2, HDAC-4, and HDAC-6 expression in human malignant and benign thyroid lesions. <i>Tumor Biology</i> , 2014 , 35, 61-71	2.9	30
107	Histone deacetylase (HDAC)-1, -2, -4 and -6 expression in human pancreatic adenocarcinoma: associations with clinicopathological parameters, tumor proliferative capacity and patients survival. <i>BMC Gastroenterology</i> , 2015 , 15, 148	3	29
106	Metallothionein expression in mobile tongue squamous cell carcinoma: associations with clinicopathological parameters and patient survival. <i>Histopathology</i> , 2011 , 59, 514-25	7.3	29

105	Retention of structurally diverse drugs in human serum albumin chromatography and its potential to simulate plasma protein binding. <i>Journal of Chromatography A</i> , 2010 , 1217, 5761-8	4.5	29
104	Pregnane X receptor (PXR) at the crossroads of human metabolism and disease. <i>Current Drug Metabolism</i> , 2013 , 14, 341-50	3.5	28
103	Farnesoid-X Receptor (FXR) as a Promising Pharmaceutical Target in Atherosclerosis. <i>Current Medicinal Chemistry</i> , 2017 , 24, 1147-1157	4.3	27
102	Clinical implication of plasma neutrophil gelatinase-associated lipocalin (NGAL) concentrations in patients with advanced carotid atherosclerosis. <i>Clinical Chemistry and Laboratory Medicine</i> , 2010 , 48, 1035-41	5.9	26
101	Peroxisome proliferator-activated receptor-gamma ligands as investigational modulators of angiogenesis. <i>Expert Opinion on Investigational Drugs</i> , 2007 , 16, 1561-72	5.9	26
100	Expression and clinical significance of focal adhesion kinase in the two distinct histological types, intestinal and diffuse, of human gastric adenocarcinoma. <i>Pathology and Oncology Research</i> , 2009 , 15, 173-81	2.6	25
99	Quantitative Structure-Retention Relationships as Useful Tool to Characterize Chromatographic Systems and Their Potential to Simulate Biological Processes. <i>Chromatographia</i> , 2013 , 76, 211-226	2.1	23
98	Peroxisome Proliferator-Activated Receptor-gamma Ligands: Potential Pharmacological Agents for Targeting the Angiogenesis Signaling Cascade in Cancer. <i>PPAR Research</i> , 2008 , 2008, 431763	4.3	23
97	Investigation of the lipophilic behaviour of some thiazolidinediones. Relationships with PPAR-gamma activity. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2007 , 857, 181-7	3.2	23
96	Hu-antigen receptor (HuR) and cyclooxygenase-2 (COX-2) expression in human non-small-cell lung carcinoma: associations with clinicopathological parameters, tumor proliferative capacity and patients survival. <i>Tumor Biology</i> , 2015 , 36, 315-27	2.9	22
95	Coxsackievirus and adenovirus receptor expression in human endometrial adenocarcinoma: possible clinical implications. <i>World Journal of Surgical Oncology</i> , 2008 , 6, 59	3.4	22
94	Physicochemical profile of nimesulide. Exploring the interplay of lipophilicity, solubility and ionization. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2007 , 44, 57-62	3.5	22
93	Evaluating Mediterranean diet adherence in university student populations: Does this dietary pattern affect students academic performance and mental health?. <i>International Journal of Health Planning and Management</i> , 2020 , 35, 5-21	2.2	22
92	Wine: An Aspiring Agent in Promoting Longevity and Preventing Chronic Diseases. <i>Diseases (Basel, Switzerland)</i> , 2018 , 6,	4.4	21
91	Expression of peroxisome proliferator activated receptor-gamma (PPAR- γ) in human non-small cell lung carcinoma: correlation with clinicopathological parameters, proliferation and apoptosis related molecules and patients survival. <i>Pathology and Oncology Research</i> , 2012 , 18, 875-83	2.6	21
90	Correlation of Peroxisome Proliferator-Activated Receptor-gamma (PPAR-gamma) and Retinoid X Receptor-alpha (RXR-alpha) expression with clinical risk factors in patients with advanced carotid atherosclerosis. <i>Medical Science Monitor</i> , 2011 , 17, CR381-91	3.2	21
89	Peroxisome proliferator-activated receptor- γ in mobile tongue squamous cell carcinoma: associations with clinicopathological parameters and patients survival. <i>Journal of Cancer Research and Clinical Oncology</i> , 2011 , 137, 251-9	4.9	20
88	Clinical significance of tumor-associated antigen RCAS1 expression in human pancreatic ductal adenocarcinoma. <i>Digestive Diseases and Sciences</i> , 2008 , 53, 1728-34	4	20

87	Expression of DNA repair proteins, MSH2, MLH1 and MGMT in mobile tongue squamous cell carcinoma: associations with clinicopathological parameters and patients survival. <i>Journal of Oral Pathology and Medicine</i> , 2011 , 40, 218-26	3.3	19
86	Primary extramammary invasive Paget's vulvar disease: what is the standard, what are the challenges and what is the future for radiotherapy?. <i>BMC Cancer</i> , 2016 , 16, 563	4.8	19
85	Evaluation of FAK and Src expression in human benign and malignant thyroid lesions. <i>Pathology and Oncology Research</i> , 2010 , 16, 497-507	2.6	18
84	High Farnesoid X Receptor (FXR) expression is a strong and independent prognosticator in invasive breast carcinoma. <i>Neoplasia</i> , 2017 , 64, 633-639	3.3	17
83	Involvement of hepatic stimulator substance in experimentally induced fibrosis and cirrhosis in the rat. <i>Digestive Diseases and Sciences</i> , 2009 , 54, 2367-76	4	17
82	Evaluation of the clinical significance of focal adhesion kinase and SRC expression in human pancreatic ductal adenocarcinoma. <i>Pancreas</i> , 2010 , 39, 930-6	2.6	17
81	Cyclin D1 and Rb protein expression and their correlation with prognosis in patients with colon cancer. <i>World Journal of Surgical Oncology</i> , 2006 , 4, 5	3.4	17
80	Peroxisome proliferator-activated receptor-gamma and retinoid X receptor-alpha expression in pancreatic ductal adenocarcinoma: association with clinicopathological parameters, tumor proliferative capacity, and patients survival. <i>Medical Science Monitor</i> , 2009 , 15, BR148-56	3.2	17
79	Platelet-activating factor involvement in thioacetamide-induced experimental liver fibrosis and cirrhosis. <i>Digestive Diseases and Sciences</i> , 2010 , 55, 276-84	4	16
78	PPAR-gamma signaling pathway in placental development and function: a potential therapeutic target in the treatment of gestational diseases. <i>Expert Opinion on Therapeutic Targets</i> , 2008 , 12, 1049-63	6.4	16
77	Farnesoid X Receptor (FXR) from normal to malignant state. <i>Histology and Histopathology</i> , 2012 , 27, 835-53	5.3	16
76	Nutritional status is associated with the degree of cognitive impairment and depressive symptoms in a Greek elderly population. <i>Nutritional Neuroscience</i> , 2020 , 23, 201-209	3.6	16
75	Farnesoid x receptor in human metabolism and disease: the interplay between gene polymorphisms, clinical phenotypes and disease susceptibility. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2015 , 11, 523-32	5.5	15
74	Salivaomics for oral diseases biomarkers detection. <i>Expert Review of Molecular Diagnostics</i> , 2016 , 16, 285-95	3.8	15
73	Quantitative structure-activity relationship (QSAR) methodology in forensic toxicology: modeling postmortem redistribution of structurally diverse drugs using multivariate statistics. <i>Forensic Science International</i> , 2009 , 190, 9-15	2.6	15
72	A consideration of PPAR-gamma ligands with respect to lipophilicity: current trends and perspectives. <i>Expert Opinion on Investigational Drugs</i> , 2007 , 16, 413-7	5.9	15
71	The impact of physicochemical and molecular properties in drug design: navigation in the "drug-like" chemical space. <i>Advances in Experimental Medicine and Biology</i> , 2015 , 822, 187-94	3.6	14
70	Investigation of the retention behavior of structurally diverse drugs on alpha(1) acid glycoprotein column: insight on the molecular factors involved and correlation with protein binding data. <i>European Journal of Pharmaceutical Sciences</i> , 2014 , 60, 24-31	5.1	14

69	FAK and Src expression in mobile tongue squamous cell carcinoma: associations with clinicopathological parameters and patients survival. <i>Journal of Cancer Research and Clinical Oncology</i> , 2012 , 138, 1369-77	4.9	14
68	Eph-A2 and Eph-A4 expression in human benign and malignant thyroid lesions: an immunohistochemical study. <i>Medical Science Monitor</i> , 2011 , 17, BR257-65	3.2	14
67	Assessing drug transport across the human placental barrier: from in vivo and in vitro measurements to the ex vivo perfusion method and in silico techniques. <i>Current Pharmaceutical Biotechnology</i> , 2011 , 12, 804-13	2.6	14
66	Current State of the Art on the Antioxidant Activity of Sage (<i>Salvia</i> spp.) and Its Bioactive Components. <i>Planta Medica</i> , 2020 , 86, 224-238	3.1	13
65	Applying quantitative structure-activity relationship (QSAR) methodology for modeling postmortem redistribution of benzodiazepines and tricyclic antidepressants. <i>Journal of Analytical Toxicology</i> , 2014 , 38, 242-8	2.9	13
64	Extracellular signal-regulated kinase (ERK) expression and activation in mobile tongue squamous cell carcinoma: associations with clinicopathological parameters and patients survival. <i>Tumor Biology</i> , 2014 , 35, 6455-65	2.9	13
63	Quantitative structure-activity relationships for PPAR-gamma binding and gene transactivation of tyrosine-based agonists using multivariate statistics. <i>Chemical Biology and Drug Design</i> , 2008 , 72, 257-64	2.9	13
62	Clinical utility of microRNAs in renal cell carcinoma: current evidence and future perspectives. <i>Expert Review of Molecular Diagnostics</i> , 2018 , 18, 981-991	3.8	13
61	Clinical Significance of Hu-Antigen Receptor (HuR) and Cyclooxygenase-2 (COX-2) Expression in Human Malignant and Benign Thyroid Lesions. <i>Pathology and Oncology Research</i> , 2016 , 22, 189-96	2.6	12
60	Evaluation of cannabinoid CB1 and CB2 receptors expression in mobile tongue squamous cell carcinoma: associations with clinicopathological parameters and patients survival. <i>Tumor Biology</i> , 2016 , 37, 3647-56	2.9	12
59	Elevated Farnesoid X Receptor (FXR) and Retinoid X Receptors (RXRs) expression is associated with less tumor aggressiveness and favourable prognosis in patients with pancreatic adenocarcinoma. <i>Neoplasia</i> , 2015 , 62, 332-41	3.3	12
58	Toll-like receptor 4 immunohistochemical expression is enhanced in macrophages of symptomatic carotid atherosclerotic plaques. <i>Cerebrovascular Diseases</i> , 2011 , 31, 29-36	3.2	12
57	Mediterranean diet adherence is associated with better cognitive status and less depressive symptoms in a Greek elderly population. <i>Aging Clinical and Experimental Research</i> , 2021 , 33, 1033-1040	4.8	12
56	Ephrin receptor (Eph) -A1, -A2, -A4 and -A7 expression in mobile tongue squamous cell carcinoma: associations with clinicopathological parameters and patients survival. <i>Pathology and Oncology Research</i> , 2014 , 20, 277-84	2.6	11
55	Correlation of plasma osteoprotegerin (OPG) and receptor activator of the nuclear factor κ B ligand (RANKL) levels with clinical risk factors in patients with advanced carotid atherosclerosis. <i>Medical Science Monitor</i> , 2012 , 18, CR597-604	3.2	11
54	Evaluation of coxsackievirus and adenovirus receptor expression in human benign and malignant thyroid lesions. <i>Apmis</i> , 2010 , 118, 210-21	3.4	11
53	Application of the ion pair concept to the n-octanol-water partitioning of cefepime and cefpirome. <i>International Journal of Pharmaceutics</i> , 2006 , 316, 52-7	6.5	10
52	Dietary Supplements on Controlling Multiple Sclerosis Symptoms and Relapses: Current Clinical Evidence and Future Perspectives. <i>Medicines (Basel, Switzerland)</i> , 2019 , 6,	4.1	9

51	Higher adherence to Mediterranean diet prior to pregnancy is associated with decreased risk for deviation from the maternal recommended gestational weight gain. <i>International Journal of Food Sciences and Nutrition</i> , 2018 , 69, 84-92	3.7	9
50	Retention behavior of flavonoids on immobilized artificial membrane chromatography and correlation with cell-based permeability. <i>Biomedical Chromatography</i> , 2018 , 32, e4108	1.7	9
49	Histone deacetylase inhibitors in oral squamous cell carcinoma treatment. <i>Expert Opinion on Investigational Drugs</i> , 2015 , 24, 69-78	5.9	9
48	Platelet-activating factor (PAF) receptor expression is associated with histopathological stage and grade and patients survival in gastric adenocarcinoma. <i>Neoplasia</i> , 2014 , 61, 309-17	3.3	9
47	Peroxisome proliferator-activated receptor-gamma ligands as bone turnover modulators. <i>Expert Opinion on Investigational Drugs</i> , 2007 , 16, 195-207	5.9	9
46	Cranberry Consumption Against Urinary Tract Infections: Clinical Stateof- the-Art and Future Perspectives. <i>Current Pharmaceutical Biotechnology</i> , 2018 , 19, 1049-1063	2.6	9
45	Serum tissue inhibitor of metalloproteinase 1 and 2 (TIMP-1 and TIMP-2) levels in colorectal cancer patients: associations with clinicopathological variables and patient survival. <i>International Journal of Biological Markers</i> , 2009 , 24, 245-52	2.8	9
44	Clinical Evaluation of the Effect of Blueberries Consumption on Chronic Diseases, Illness Prevention and Health Promotion. <i>Natural Products Journal</i> , 2018 , 8, 45-53	0.6	8
43	Structural basis for the design of PPAR-gamma ligands: a survey on quantitative structure- activity relationships. <i>Mini-Reviews in Medicinal Chemistry</i> , 2009 , 9, 1075-83	3.2	8
42	A QSAR Study on Indole-Based PPAR- γ Agonists in Respect to Receptor Binding and Gene Transactivation Data. <i>QSAR and Combinatorial Science</i> , 2009 , 28, 802-805		8
41	Diagnostic and Prognostic Utility of Serum Receptor-Binding Cancer Antigen Expressed on SiSo Cells (RCAS1) Levels in Colon Cancer Patients. <i>International Journal of Biological Markers</i> , 2009 , 24, 70-76	2.8	8
40	Effect of the aqueous component of the mobile phase on RP-TLC retention and its implication in the determination of lipophilicity for a series of structurally diverse drugs. <i>Journal of Planar Chromatography - Modern TLC</i> , 2006 , 19, 151-156	0.9	8
39	Expression of DNA repair proteins MSH2, MLH1 and MGMT in human benign and malignant thyroid lesions: an immunohistochemical study. <i>Medical Science Monitor</i> , 2011 , 17, BR81-90	3.2	8
38	Angiogenesis in salivary gland tumors: from clinical significance to treatment. <i>Expert Opinion on Therapeutic Targets</i> , 2015 , 19, 807-19	6.4	7
37	Clinical value of nutritional status in neurodegenerative diseases: What is its impact and how it affects disease progression and management?. <i>Nutritional Neuroscience</i> , 2018 , 21, 162-175	3.6	7
36	Clinical Significance of EphB4 and EphB6 Expression in Human Malignant and Benign Thyroid Lesions. <i>Pathology and Oncology Research</i> , 2016 , 22, 269-75	2.6	7
35	Analysis of PPAR- γ Activity by Combining 2-D QSAR and Molecular Simulation. <i>Molecular Informatics</i> , 2013 , 32, 431-45	3.8	7
34	Expression and Clinical Significance of Concomitant FAK/SRC and p-Paxillin in Mobile Tongue Squamous Cell Carcinoma. <i>Anticancer Research</i> , 2017 , 37, 1313-1319	2.3	7

33	Elevated Hu-Antigen Receptor (HuR) Expression is Associated with Tumor Aggressiveness and Poor Prognosis but not with COX-2 Expression in Invasive Breast Carcinoma Patients. <i>Pathology and Oncology Research</i> , 2018 , 24, 631-640	2.6	7
32	Clinical significance of farnesoid X receptor expression in thyroid neoplasia. <i>Future Oncology</i> , 2017 , 13, 1785-1792	3.6	6
31	The Impact of Lipophilicity in Drug Discovery: Rapid Measurements by Means of Reversed-Phase HPLC. <i>Methods in Molecular Biology</i> , 2018 , 1824, 217-228	1.4	6
30	Estimating the agreement between the metabolic rate calculated from prediction equations and from a portable indirect calorimetry device: an effort to develop a new equation for predicting resting metabolic rate. <i>Nutrition and Metabolism</i> , 2018 , 15, 41	4.6	6
29	Detection of dental plaque with disclosing agents in the context of preventive oral hygiene training programs. <i>Heliyon</i> , 2019 , 5, e02064	3.6	6
28	Octanol/water partitioning simulation by RP-HPLC for structurally diverse acidic drugs: comparison of three columns in the presence and absence of n-octanol as the mobile phase additive. <i>Journal of Separation Science</i> , 2013 , 36, 3830-6	3.4	6
27	Phosphorylated Epidermal Growth Factor Receptor Expression Is Associated With Clinicopathologic Parameters and Patient Survival in Mobile Tongue Squamous Cell Carcinoma. <i>Journal of Oral and Maxillofacial Surgery</i> , 2017 , 75, 632-640	1.8	6
26	Pregnane X Receptor Expression in Human Pancreatic Adenocarcinoma: Associations With Clinicopathologic Parameters, Tumor Proliferative Capacity, Patients Survival, and Retinoid X Receptor Expression. <i>Pancreas</i> , 2015 , 44, 1134-40	2.6	6
25	Increased RCAS1 expression is associated with advanced histopathological stage and poor prognosis in patients with gastric adenocarcinoma. <i>Disease Markers</i> , 2013 , 35, 213-9	3.2	6
24	Serum Tissue Inhibitor of Metalloproteinase 1 and 2 (TIMP-1 and TIMP-2) Levels in Colorectal Cancer Patients: Associations with Clinicopathological Variables and Patient Survival. <i>International Journal of Biological Markers</i> , 2009 , 24, 245-252	2.8	6
23	Cranberry as promising natural source of potential anticancer agents: current evidence and future perspectives. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2012 , 12, 619-30	2.2	6
22	RPTLC retention indices of basic and neutral drugs as surrogates of octanol-water distribution coefficients. Effect of buffer constituents and pH. <i>Journal of Planar Chromatography - Modern TLC</i> , 2009 , 22, 217-224	0.9	6
21	Chromatographic behavior of zwitterionic enalapril-exploring the conditions for lipophilicity assessment. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2008 , 48, 739-43	3.5	6
20	Diagnostic and prognostic utility of serum receptor-binding cancer antigen expressed on SiSo cells (RCAS1) levels in colon cancer patients. <i>International Journal of Biological Markers</i> , 2009 , 24, 70-6	2.8	6
19	Simple physicochemical properties related with lipophilicity, polarity, molecular size and ionization status exert significant impact on the transfer of drugs and chemicals into human breast milk. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2016 , 12, 1273-1278	5.5	5
18	Clinical Significance of Cannabinoid Receptors CB1 and CB2 Expression in Human Malignant and Benign Thyroid Lesions. <i>BioMed Research International</i> , 2015 , 2015, 839403	3	5
17	Expression and clinical significance of FAK and Src proteins in human endometrial adenocarcinoma. <i>Pathology and Oncology Research</i> , 2011 , 17, 277-85	2.6	5
16	The performance of 1-ethyl-3-methylimidazolium tetrafluoroborate ionic liquid as mobile phase additive in HPLC-based lipophilicity assessment. <i>Biomedical Chromatography</i> , 2011 , 25, 606-12	1.7	5

15	Cranberry: A Promising Natural Source of Potential Nutraceuticals with Anticancer Activity. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2019 , 19, 1672-1686	2.2	5
14	RCAS1 expression in mobile tongue squamous cell carcinoma: an immunohistochemical study. <i>Medical Science Monitor</i> , 2011 , 17, BR228-234	3.2	4
13	Current Advances on the Extraction and Identification of Bioactive Components of Sage (<i>Salvia</i> spp.). <i>Current Pharmaceutical Biotechnology</i> , 2019 , 20, 845-857	2.6	4
12	Receptor-binding cancer antigen expressed on SiSo cells (RCAS1) expression in human benign and malignant thyroid lesions. <i>Medical Science Monitor</i> , 2012 , 18, BR123-9	3.2	4
11	Medicinal plants consumption against urinary tract infections: a narrative review of the current evidence. <i>Expert Review of Anti-Infective Therapy</i> , 2021 , 19, 519-528	5.5	4
10	Insight into pain-inducing and -related gene expression: a challenge for development of novel targeted therapeutic approaches. <i>Fundamental and Clinical Pharmacology</i> , 2011 , 25, 48-62	3.1	3
9	Current Clinical Status on the Estimation of Energy Requirement: Searching for a Reliable Equation to Predict Energy Requirement in Multiple Populations. <i>Current Nutrition and Food Science</i> , 2019 , 14, 375-385	0.7	3
8	Histone deacetylase inhibitors as potential therapeutic agents for the treatment of malignant mesothelioma. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2013 , 13, 476-82	2.2	3
7	Can Bioelectrical Impedance Analysis and BMI be a Prognostic Tool in Head and Neck Cancer Patients? A Review of the Evidence. <i>Cancers</i> , 2020 , 12,	6.6	2
6	Endocannabinoid System: A Promising Therapeutic Target for the Treatment of Haematological Malignancies?. <i>Current Medicinal Chemistry</i> , 2016 , 23, 2350-62	4.3	2
5	QSAR/QSPR Modeling in the Design of Drug Candidates with Balanced Pharmacodynamic and Pharmacokinetic Properties. <i>Challenges and Advances in Computational Chemistry and Physics</i> , 2017 , 339-384	9.7	2
4	Histone Deacetylase Inhibitors: A Promising Therapeutic Alternative for Endometrial Carcinoma. <i>Disease Markers</i> , 2021 , 2021, 7850688	3.2	1
3	HuR as Therapeutic Target in Cancer: What the Future Holds. <i>Current Medicinal Chemistry</i> , 2021 ,	4.3	1
2	The clinical and pathological significance of RCAS1 expression as a prognostic biomarker in non-small cell lung cancer. <i>In Vivo</i> , 2014 , 28, 375-81	2.3	1
1	The clinical significance of ephrin receptors expression in non-small cell lung cancers.. <i>Journal of Clinical Oncology</i> , 2013 , 31, e22160-e22160	2.2	