
Chee-Wee Tan

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8560/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Effectiveness of the Alfredson Protocol Compared With a Lower Repetition-Volume Protocol for Midportion Achilles Tendinopathy: A Randomized Controlled Trial. Journal of Orthopaedic and Sports Physical Therapy, 2014, 44, 59-67.	1.7	77
2	Measurement of lower limb volume: Agreement between the vertically oriented perometer and a tape measure method. Physiotherapy, 2013, 99, 247-251.	0.2	42
3	Comparison of Breast Cancer-Related Lymphedema (Upper Limb Swelling) Prevalence Estimated Using Objective and Subjective Criteria and Relationship with Quality of Life. BioMed Research International, 2013, 2013, 1-8.	0.9	38
4	A Morbidity Screening Tool for identifying fatigue, pain, upper limb dysfunction and lymphedema after breast cancer treatment: A validity study. European Journal of Oncology Nursing, 2014, 18, 218-227.	0.9	37
5	Discrimination of Real and Sham Acupuncture Needles Using the Park Sham Device: A Preliminary Study. Archives of Physical Medicine and Rehabilitation, 2009, 90, 2141-2145.	0.5	18
6	Falls efficacy instruments for community-dwelling older adults: a COSMIN-based systematic review. BMC Geriatrics, 2021, 21, 21.	1.1	16
7	Falls efficacy: Extending the understanding of self-efficacy in older adults towards managing falls. Journal of Frailty, Sarcopenia and Falls, 2021, 06, 131-138.	0.4	16
8	Prevalence and Impacts of Upper Limb Morbidity after Treatment for Breast Cancer: A Cross-Sectional Study of Lymphedema and Function. Cancer and Oncology Research, 2013, 1, 30-39.	0.2	16
9	Modification of Experimental, Lower Limb Ischemic Pain With Transcutaneous Electrical Nerve Stimulation. Clinical Journal of Pain, 2012, 28, 693-699.	0.8	11
10	Discrimination Accuracy between Real and Sham Needles using the Park Sham Device in the Upper and Lower Limbs. Acupuncture in Medicine, 2011, 29, 208-214.	0.4	10
11	Transcutaneous Electrical Nerve Stimulation Improves Walking Performance in Patients With Intermittent Claudication. Journal of Cardiovascular Nursing, 2016, 31, 323-330.	0.6	9
12	Perometry limb volume measurement: protocol development and reliability. European Journal of Physiotherapy, 2013, 15, 193-200.	0.7	7
13	Detection theory analysis of scaling and discrimination tasks: Responses to noxious thermal stimuli. Perception & Psychophysics, 2007, 69, 994-1001.	2.3	4
14	25. Upper limb morbidity after treatment for breast cancer: A cross-sectional study of lymphedema and function. European Journal of Surgical Oncology, 2012, 38, 1141.	0.5	3
15	Primary care lymphoedema service reduces incidence of infection, antibiotic use, hospitalisation and decreased limb volume: a service evaluation. Physiotherapy, 2015, 101, e1358-e1359.	0.2	1
16	The influence of upper limb dominance on limb volume measurements made by perometry and water displacement volumetry. Physiotherapy, 2015, 101, e1359.	0.2	1
17	90 Falls Efficacy Related Instruments for Community-Dwellig Older Adults – Development and Content Validity: A Cosmin-Based Systematic Review. Age and Ageing, 2019, 48, iv18-iv27.	0.7	1
18	Contribution of Vision, Touch, and Hearing to the Use of Sham Devices in Acupuncture-Related Studies. JAMS Journal of Acupuncture and Meridian Studies, 2020, 13, 25-32.	0.3	1

#	Article	IF	CITATIONS
19	Validation of a new patient-reported outcome measure of balance recovery confidence (BRC) for community-dwelling older adults: a study protocol. Physical Therapy Reviews, 2021, 26, 457-466.	0.3	1
20	The effect of stimulus number on the stability of responses for an extensive heat pain test. Somatosensory & Motor Research, 2012, 29, 62-69.	0.4	0