

# Eeva-Mari Jouhilahti

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8560008/publications.pdf>

Version: 2024-02-01

21  
papers

912  
citations

686830

13  
h-index

713013

21  
g-index

25  
all docs

25  
docs citations

25  
times ranked

1594  
citing authors

#	ARTICLE	IF	CITATIONS
1	CRISPR activation enables high-fidelity reprogramming into human pluripotent stem cells. <i>Stem Cell Reports</i> , 2022, 17, 413-426.	2.3	13
2	DUX4 is a multifunctional factor priming human embryonic genome activation. <i>IScience</i> , 2022, 25, 104137.	1.9	20
3	Transient DUX4 expression in human embryonic stem cells induces blastomere-like expression program that is marked by SLC34A2. <i>Stem Cell Reports</i> , 2022, 17, 1743-1756.	2.3	11
4	Embryonic LTR retrotransposons supply promoter modules to somatic tissues. <i>Genome Research</i> , 2021, 31, 1983-1993.	2.4	7
5	Characterization of the human RFX transcription factor family by regulatory and target gene analysis. <i>BMC Genomics</i> , 2018, 19, 181.	1.2	73
6	Phylogenetic and mutational analyses of human LEUTX, a homeobox gene implicated in embryogenesis. <i>Scientific Reports</i> , 2018, 8, 17421.	1.6	17
7	Human pluripotent reprogramming with CRISPR activators. <i>Nature Communications</i> , 2018, 9, 2643.	5.8	128
8	The human PRD-like homeobox gene <i>LEUTX</i> has a central role in embryo genome activation. <i>Development (Cambridge)</i> , 2016, 143, 3459-3469.	1.2	42
9	An approach to comprehensive genome and proteome expression analyses in Schwann cells and neurons during peripheral nerve myelin formation. <i>Journal of Neurochemistry</i> , 2016, 138, 830-844.	2.1	10
10	Characterization and target genes of nine human PRD-like homeobox domain genes expressed exclusively in early embryos. <i>Scientific Reports</i> , 2016, 6, 28995.	1.6	33
11	Globin mRNA reduction for whole-blood transcriptome sequencing. <i>Scientific Reports</i> , 2016, 6, 31584.	1.6	42
12	Gene expression analysis of skin grafts and cultured keratinocytes using synthetic RNA normalization reveals insights into differentiation and growth control. <i>BMC Genomics</i> , 2015, 16, 476.	1.2	21
13	Novel PRD-like homeodomain transcription factors and retrotransposon elements in early human development. <i>Nature Communications</i> , 2015, 6, 8207.	5.8	100
14	Hypoxic conditions stimulate the release of B $\alpha$ type natriuretic peptide from human retinal pigment epithelium cell culture. <i>Acta Ophthalmologica</i> , 2014, 92, 740-744.	0.6	6
15	Neurofibromatosis Type 1 Gene Mutation Analysis Using Sequence Capture and High-throughput Sequencing. <i>Acta Dermato-Venereologica</i> , 2014, 94, 663-666.	0.6	8
16	Oral soft tissue alterations in patients with neurofibromatosis. <i>Clinical Oral Investigations</i> , 2012, 16, 551-558.	1.4	37
17	Molecular and Cellular Basis of Human Cutaneous Neurofibromas and Their Development. , 2012, , 393-403.		2
18	The Development of Cutaneous Neurofibromas. <i>American Journal of Pathology</i> , 2011, 178, 500-505.	1.9	63

#	ARTICLE	IF	CITATIONS
19	The Pathoetiology of Neurofibromatosis 1. American Journal of Pathology, 2011, 178, 1932-1939.	1.9	145
20	Reevaluation of the Normal Epidermal Calcium Gradient, and Analysis of Calcium Levels and ATP Receptors in Haileyâ€“Hailey and Darier Epidermis. Journal of Investigative Dermatology, 2009, 129, 1379-1387.	0.3	55
21	Class III $\beta$ -Tubulin Is a Component of the Mitotic Spindle in Multiple Cell Types. Journal of Histochemistry and Cytochemistry, 2008, 56, 1113-1119.	1.3	64