

# Hidemasa Nakaminami

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8564833/publications.pdf>

Version: 2024-02-01

99  
papers

1,469  
citations

361045

20  
h-index

414034

32  
g-index

103  
all docs

103  
docs citations

103  
times ranked

1507  
citing authors

#	ARTICLE	IF	CITATIONS
1	Familial infections caused by nasal colonization of Panton-Valentine leukocidin-positive methicillin-resistant <i>Staphylococcus aureus</i> USA300 clone. <i>Journal of Dermatology</i> , 2022, 49, .	0.6	1
2	Comparative Genomic Analysis of the Human Variant of Methicillin-Resistant <i>Staphylococcus aureus</i> CC398 in Japan and Korea. <i>Microbial Drug Resistance</i> , 2022, , .	0.9	3
3	High-Level Quinolone-Resistant <i>Haemophilus haemolyticus</i> in Pediatric Patient with No History of Quinolone Exposure. <i>Emerging Infectious Diseases</i> , 2022, 28, 105-111.	2.0	4
4	Multidrug-resistant <i>Cutibacterium avidum</i> isolated from patients with acne vulgaris and other infections. <i>Journal of Global Antimicrobial Resistance</i> , 2022, 28, 151-157.	0.9	10
5	Clinical Efficacy and Safety of Arbekacin against Pneumonia in Febrile Neutropenia: A Retrospective Study in Patients with Hematologic Malignancies. <i>Infection and Chemotherapy</i> , 2022, 54, 80.	1.0	1
6	Quinolone Resistance Is Transferred Horizontally via Uptake Signal Sequence Recognition in <i>Haemophilus influenzae</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2022, 66, AAC0196721.	1.4	4
7	Identification of a Transferable Linear Plasmid Carrying the Macrolide-Clindamycin Resistance Gene <i>erm</i> (X) in a <i>Cutibacterium acnes</i> Isolate from a Patient with Acne Vulgaris in Japan. <i>Microbiology Resource Announcements</i> , 2022, 11, e0009422.	0.3	4
8	Identification and characterisation of a novel multidrug-resistant streptococcus, <i>Streptococcus toyakuensis</i> sp. nov., from a blood sample. <i>Journal of Global Antimicrobial Resistance</i> , 2022, 29, 316-322.	0.9	8
9	Increased Prevalence of Minocycline-Resistant <i>Staphylococcus epidermidis</i> with <i>tet</i> (M) by Tetracycline Use for Acne Treatment. <i>Microbial Drug Resistance</i> , 2022, 28, 861-866.	0.9	2
10	A case of <i>Shewanella algae</i> -induced bacteremia in Japan: Case report and literature review. <i>Journal of Infection and Chemotherapy</i> , 2022, 28, 1430-1432.	0.8	4
11	In vitro anti-biofilm effect of anti-methicillin-resistant <i>Staphylococcus aureus</i> (anti-MRSA) agents against the USA300 clone. <i>Journal of Global Antimicrobial Resistance</i> , 2021, 24, 63-71.	0.9	14
12	Possible Dissemination of a Panton-Valentine Leukocidin-Positive Livestock-Associated Methicillin-Resistant <i>Staphylococcus aureus</i> CC398 Clone in Tokyo, Japan. <i>Japanese Journal of Infectious Diseases</i> , 2021, 74, 82-84.	0.5	8
13	Prevalence of antimicrobial-resistant staphylococci in nares and affected sites of pet dogs with superficial pyoderma. <i>Journal of Veterinary Medical Science</i> , 2021, 83, 214-219.	0.3	8
14	A family case of community-acquired methicillin-resistant <i>Staphylococcus aureus</i> infection with USA300 clone. <i>Journal of Dermatology</i> , 2021, 48, e210-e211.	0.6	3
15	In vitro growth-inhibitory effects of <i>Portulaca oleracea</i> L. formulation on intestinal pathogens. <i>Access Microbiology</i> , 2021, 3, 000208.	0.2	4
16	Increased prevalence of doxycycline low-susceptible <i>Cutibacterium acnes</i> isolated from acne patients in Japan caused by antimicrobial use and diversification of tetracycline resistance factors. <i>Journal of Dermatology</i> , 2021, 48, 1365-1371.	0.6	9
17	<i>Escherichia coli</i> GyrA Tower Domain Interacts with QnrB1 Loop B and Plays an Important Role in QnrB1 Protection from Quinolone Inhibition. <i>Antimicrobial Agents and Chemotherapy</i> , 2021, 65, e0040221.	1.4	0
18	Comparison of the bactericidal effects of quinolones against low-susceptible <i>Haemophilus influenzae</i> . <i>Journal of Medical Microbiology</i> , 2021, 70, .	0.7	0

#	ARTICLE	IF	CITATIONS
19	Antimicrobial activity and additive effect of the modified Gingyo-san with antimicrobials against <i>Helicobacter pylori</i> . <i>Journal of Infection and Chemotherapy</i> , 2021, 27, 957-961.	0.8	1
20	Dissemination of quinolone low-susceptible <i>Haemophilus influenzae</i> ST422 in Tokyo, Japan. <i>Journal of Infection and Chemotherapy</i> , 2021, 27, 962-966.	0.8	4
21	Recurrent furunculosis by Panton-Valentine leukocidin-positive methicillin-resistant <i>Staphylococcus aureus</i> USA300 clone in Tokyo: A report of two cases. <i>Journal of Dermatology</i> , 2021, 48, E522-E523.	0.6	3
22	Chinese herbal medicines and nutraceuticals inhibit <i>Pseudomonas aeruginosa</i> biofilm formation. <i>Access Microbiology</i> , 2021, 3, 000254.	0.2	1
23	<i>Cutibacterium acnes</i> phylogenetic type IC and II isolated from patients with non-acne diseases exhibit high-level biofilm formation. <i>International Journal of Medical Microbiology</i> , 2021, 311, 151538.	1.5	11
24	Molecular characterisation of carbapenem- and tigecycline-resistant <i>Klebsiella pneumoniae</i> strains isolated from blood and bile samples. <i>Journal of Infection and Chemotherapy</i> , 2021, , .	0.8	1
25	Trends in Panton-Valentine Leukocidin (PVL)-Positive Methicillin-Resistant <i>Staphylococcus aureus</i> (MRSA) in Outpatients of a University Hospital. <i>Iryo Yakugaku (Japanese Journal of)</i> Tj ETQq1 1 0.784314 rgBT /Overlo		
26	An outbreak of severe infectious diseases caused by methicillin-resistant <i>Staphylococcus aureus</i> USA300 clone among hospitalized patients and nursing staff in a tertiary care university hospital. <i>Journal of Infection and Chemotherapy</i> , 2020, 26, 76-81.	0.8	23
27	Combination effects of modified Gingyo-san extract and antimicrobial agents. <i>European Journal of Integrative Medicine</i> , 2020, 33, 101016.	0.8	1
28	Detection of Panton-Valentine leukocidin-positive livestock-associated <i>Staphylococcus aureus</i> CC398 clone in a Vietnamese patient in Japan. <i>Journal of Global Antimicrobial Resistance</i> , 2020, 23, 72-73.	0.9	2
29	Arthritis Caused by MRSA CC398 in a Patient without Animal Contact, Japan. <i>Emerging Infectious Diseases</i> , 2020, 26, 3015-3105.	2.0	2
30	First isolation of an IMP-1 metallo- $\beta$ -lactamase-producing <i>Kluyvera ascorbata</i> in Japan. <i>Journal of Global Antimicrobial Resistance</i> , 2020, 23, 228-231.	0.9	1
31	Current status of Panton-Valentine leukocidin-positive methicillin-resistant <i>Staphylococcus aureus</i> isolated from patients with skin and soft tissue infections in Japan. <i>Journal of Dermatology</i> , 2020, 47, 1280-1286.	0.6	23
32	Whole-genome sequence of <i>Haemophilus influenzae</i> ST422 outbreak clone strain 2018-Y40 with low quinolone susceptibility isolated from a paediatric patient. <i>Journal of Global Antimicrobial Resistance</i> , 2020, 22, 759-761.	0.9	6
33	A novel community-acquired MRSA clone, USA300-LV/J, uniquely evolved in Japan. <i>Journal of Antimicrobial Chemotherapy</i> , 2020, 75, 3131-3134.	1.3	12
34	Phosphatidylinositol-specific phospholipase C enhances epidermal penetration by <i>Staphylococcus aureus</i> . <i>Scientific Reports</i> , 2020, 10, 17845.	1.6	7
35	Kampo medicines suppress the production of exfoliative toxins causing impetigo in <i>Staphylococcus aureus</i> . <i>Journal of Dermatology</i> , 2020, 47, 714-719.	0.6	2
36	pspK acquisition contributes to the loss of capsule in pneumococci: molecular characterisation of non-encapsulated pneumococci. <i>Microbes and Infection</i> , 2020, 22, 451-456.	1.0	4

#	ARTICLE	IF	CITATIONS
37	Arthritis Caused by MRSA CC398 in a Patient without Animal Contact, Japan. <i>Emerging Infectious Diseases</i> , 2020, 26, 795-797.	2.0	16
38	A class A $\beta$ -lactamase produced by borderline oxacillin-resistant <i>Staphylococcus aureus</i> hydrolyses oxacillin. <i>Journal of Global Antimicrobial Resistance</i> , 2020, 22, 244-247.	0.9	18
39	A risk as an infection route: Nasal colonization of methicillin-resistant <i>Staphylococcus aureus</i> USA300 clone among contact sport athletes in Japan. <i>Journal of Infection and Chemotherapy</i> , 2020, 26, 862-864.	0.8	9
40	The effects of surface properties of liposomes on their activity against <i>Pseudomonas aeruginosa</i> PAO-1 biofilm. <i>Journal of Drug Delivery Science and Technology</i> , 2020, 57, 101754.	1.4	21
41	Transferable Multidrug-Resistance Plasmid Carrying a Novel Macrolide-Clindamycin Resistance Gene, <i>ermA</i> (50), in <i>Cutibacterium acnes</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2020, 64, .	1.4	24
42	First Report of Fatal Infection Caused by Community-acquired Methicillin-resistant <i>Staphylococcus aureus</i> USA300 Clone in a Collegiate Athlete. <i>JMA Journal</i> , 2020, 3, 78-82.	0.6	4
43	Development of effective antimicrobial cocktails to prevent bacterial contamination of allograft tissues under low temperature conditions. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2019, 28, 128-136.	0.5	4
44	Relationship between quinolone use and resistance of <i>Staphylococcus epidermidis</i> in patients with acne vulgaris. <i>Journal of Dermatology</i> , 2019, 46, 782-786.	0.6	11
45	Decreased Prevalence of <i>qacA</i> -Positive Methicillin-Resistant <i>Staphylococcus aureus</i> in Hospitalized Patients in Tokyo, Japan. <i>Microbial Drug Resistance</i> , 2019, 25, 1032-1040.	0.9	2
46	Tokiinshi, a traditional Japanese medicine (Kampo), suppresses Panton-Valentine leukocidin production in the methicillin-resistant <i>Staphylococcus aureus</i> USA300 clone. <i>PLoS ONE</i> , 2019, 14, e0214470.	1.1	4
47	Clonal change of methicillin-resistant <i>Staphylococcus aureus</i> isolated from patients with impetigo in Kagawa, Japan. <i>Journal of Dermatology</i> , 2019, 46, 301-307.	0.6	10
48	A case of acute septic arthritis of the hip joint caused by Panton-Valentine leukocidin-positive ST772 community-acquired methicillin-resistant <i>Staphylococcus aureus</i> . <i>Journal of Infection and Chemotherapy</i> , 2019, 25, 212-214.	0.8	3
49	Identification and detection of USA300 methicillin-resistant <i>Staphylococcus aureus</i> clones with a partial deletion in the <i>ccrB2</i> gene on the type IV SCCmec element. <i>Diagnostic Microbiology and Infectious Disease</i> , 2019, 94, 86-87.	0.8	16
50	Evaluation of <i>in vitro</i> Bactericidal Activity of 1.5% Olanexidine Gluconate, a Novel Biguanide Antiseptic Agent. <i>Biological and Pharmaceutical Bulletin</i> , 2019, 42, 512-515.	0.6	11
51	Comparative analysis of methicillin-resistant <i>Staphylococcus aureus</i> isolated from outpatients of dermatology unit in hospitals and clinics. <i>Journal of Infection and Chemotherapy</i> , 2019, 25, 233-237.	0.8	8
52	Fast-acting bactericidal activity of olanexidine gluconate against <i>qacA/B</i> -positive methicillin-resistant <i>Staphylococcus aureus</i> . <i>Journal of Medical Microbiology</i> , 2019, 68, 957-960.	0.7	9
53	Impact of the introduction of a 13-valent pneumococcal vaccine on pneumococcal serotypes in non-invasive isolates from 2007 to 2016 at a teaching hospital in Japan. <i>Journal of Medical Microbiology</i> , 2019, 68, 903-909.	0.7	10
54	Glyceraldehyde-3-phosphate dehydrogenase of <i>Mycoplasma pneumoniae</i> induces infection-related glomerulonephritis. <i>Clinical Nephrology</i> , 2019, 92, 263-272.	0.4	11

#	ARTICLE	IF	CITATIONS
55	<i>Propionibacterium acnes</i> Has Low Susceptibility to Chlorhexidine Digluconate. Surgical Infections, 2018, 19, 298-302.	0.7	15
56	Dissemination of Panton-Valentine leukocidin-positive methicillin-resistant <i>Staphylococcus aureus</i> USA300 clone in multiple hospitals in Tokyo, Japan. Clinical Microbiology and Infection, 2018, 24, 1211.e1-1211.e7.	2.8	53
57	Change in genotype of methicillin-resistant <i>Staphylococcus aureus</i> (MRSA) affects the antibiogram of hospital-acquired MRSA. Journal of Infection and Chemotherapy, 2018, 24, 563-569.	0.8	36
58	Specific clones of <i>Staphylococcus lugdunensis</i> may be associated with colon carcinoma. Journal of Infection and Public Health, 2018, 11, 39-42.	1.9	13
59	651. Non-encapsulation of Pneumococci as a Potential Evasion Mechanism From Vaccines. Open Forum Infectious Diseases, 2018, 5, S236-S236.	0.4	0
60	Characterization of SCCmec type IV methicillin-resistant <i>Staphylococcus aureus</i> clones increased in Japanese hospitals. Journal of Medical Microbiology, 2018, 67, 769-774.	0.7	22
61	Prevalence of skin infections caused by Panton-Valentine leukocidin-positive methicillin-resistant <i>Staphylococcus aureus</i> in Japan, particularly in Ishigaki, Okinawa. Journal of Infection and Chemotherapy, 2017, 23, 800-803.	0.8	35
62	Efflux Transporter of Siderophore Staphyloferrin A in <i>Staphylococcus aureus</i> Contributes to Bacterial Fitness in Abscesses and Epithelial Cells. Infection and Immunity, 2017, 85, .	1.0	14
63	Antibiotic That Inhibits the ATPase Activity of an ATP-Binding Cassette Transporter by Binding to a Remote Extracellular Site. Journal of the American Chemical Society, 2017, 139, 10597-10600.	6.6	18
64	Evaluation of <i>In Vitro</i> Antiamoebic Activity of Antimicrobial Agents Against Clinical <i>Acanthamoeba</i> Isolates. Journal of Ocular Pharmacology and Therapeutics, 2017, 33, 629-634.	0.6	12
65	Rise in <i>Haemophilus influenzae</i> With Reduced Quinolone Susceptibility and Development of a Simple Screening Method. Pediatric Infectious Disease Journal, 2017, 36, 263-266.	1.1	17
66	First report of sasX-positive methicillin-resistant <i>Staphylococcus aureus</i> in Japan. FEMS Microbiology Letters, 2017, 364, .	0.7	9
67	Genetic diversity of pvl- positive community-onset methicillin-resistant <i>Staphylococcus aureus</i> isolated at a university hospital in Japan. Journal of Infection and Chemotherapy, 2017, 23, 856-858.	0.8	16
68	Determination of the Mutant Prevention Concentration and the Mutant Selection Window of Topical Antimicrobial Agents against <i>Propionibacterium acnes</i> . Journal of Infection and Chemotherapy, 2017, 62, 94-99.	0.8	6
69	Amino Acid Substitution in the Major Multidrug Efflux Transporter Protein AcrB Contributes to Low Susceptibility to Azithromycin in <i>Haemophilus influenzae</i> . Antimicrobial Agents and Chemotherapy, 2017, 61, .	1.4	8
70	<i>Propionibacterium acnes</i> is developing gradual increase in resistance to oral tetracyclines. Journal of Medical Microbiology, 2017, 66, 8-12.	0.7	24
71	Methicillin-Resistant <i>Staphylococcus epidermidis</i> Is Part of the Skin Flora on the Hands of Both Healthy Individuals and Hospital Workers. Biological and Pharmaceutical Bulletin, 2016, 39, 1868-1875.	0.6	11
72	A novel 23S rRNA mutation in <i>Propionibacterium acnes</i> confers resistance to 14-membered macrolides. Journal of Global Antimicrobial Resistance, 2016, 6, 160-161.	0.9	11

#	ARTICLE	IF	CITATIONS
73	Emergence of fluoroquinolone-resistant <i>Propionibacterium acnes</i> caused by amino acid substitutions of DNA gyrase but not DNA topoisomerase IV. <i>Anaerobe</i> , 2016, 42, 166-171.	1.0	21
74	The modified Gingyo-san, a Chinese herbal medicine, has direct antibacterial effects against respiratory pathogens. <i>BMC Complementary and Alternative Medicine</i> , 2016, 16, 463.	3.7	12
75	Prevalence of macrolide-non-susceptible isolates among $\beta$ -lactamase-negative ampicillin-resistant <i>Haemophilus influenzae</i> in a tertiary care hospital in Japan. <i>Journal of Global Antimicrobial Resistance</i> , 2016, 6, 22-26.	0.9	16
76	Clarithromycin Resistance Mechanisms of Epidemic $\beta$ -Lactamase-Nonproducing Ampicillin-Resistant <i>Haemophilus influenzae</i> Strains in Japan. <i>Antimicrobial Agents and Chemotherapy</i> , 2016, 60, 3207-3210.	1.4	14
77	Impact of calcium concentration in Muller-Hinton medium on the antimicrobial activity of daptomycin. <i>Journal of Global Antimicrobial Resistance</i> , 2016, 4, 76-77.	0.9	2
78	Increase in SCCmec type IV strains affects trends in antibiograms of methicillin-resistant <i>Staphylococcus aureus</i> at a tertiary-care hospital. <i>Journal of Medical Microbiology</i> , 2015, 64, 745-751.	0.7	22
79	In Vitro Antimicrobial Activity of Fibrin Sealants Containing Antimicrobial Agents. <i>Surgical Infections</i> , 2014, 15, 29-35.	0.7	10
80	Discovery of Natural Products Possessing Selective Eukaryotic Readthrough Activity: Deoxyneogamycin and Its Leucine Adduct. <i>ChemMedChem</i> , 2014, 9, 2233-2237.	1.6	18
81	Characterization of methicillin-resistant <i>Staphylococcus aureus</i> isolated from tertiary care hospitals in Tokyo, Japan. <i>Journal of Infection and Chemotherapy</i> , 2014, 20, 512-515.	0.8	36
82	Comprehensive evaluation of fibrin glue as a local drug-delivery system: efficacy and safety of sustained release of vancomycin by fibrin glue against local methicillin-resistant <i>Staphylococcus aureus</i> infection. <i>Journal of Artificial Organs</i> , 2014, 17, 42-49.	0.4	16
83	A novel GyrB mutation in methicillin-resistant <i>Staphylococcus aureus</i> (MRSA) confers a high level of resistance to third-generation quinolones. <i>International Journal of Antimicrobial Agents</i> , 2014, 43, 478-479.	1.1	11
84	Relationship between the severity of acne vulgaris and antimicrobial resistance of bacteria isolated from acne lesions in a hospital in Japan. <i>Journal of Medical Microbiology</i> , 2014, 63, 721-728.	0.7	65
85	In vitro antiseptic susceptibilities for <i>Staphylococcus pseudintermedius</i> isolated from canine superficial pyoderma in Japan. <i>Veterinary Dermatology</i> , 2013, 24, 126.	0.4	27
86	Novel Hybrid-Type Antimicrobial Agents Targeting the Switch Region of Bacterial RNA Polymerase. <i>ACS Medicinal Chemistry Letters</i> , 2013, 4, 220-224.	1.3	20
87	Antimicrobial Spectrum of Alcohol-Based Hand-Rubbing Solutions Containing 1 w/v% Chlorhexidine Gluconate. <i>Iryo Yakugaku (Japanese Journal of Pharmaceutical Health Care and Sciences)</i> , 2013, 39, 304-308.	0.0	3
88	First report of high levels of clindamycin-resistant <i>Propionibacterium acnes</i> carrying erm(X) in Japanese patients with acne vulgaris. <i>Journal of Dermatology</i> , 2012, 39, 794-796.	0.6	38
89	Effect of pretreatment with <i>Lactobacillus gasseri</i> OLL2716 on first-line <i>Helicobacter pylori</i> eradication therapy. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2012, 27, 888-892.	1.4	60
90	Susceptibility of <i>Propionibacterium acnes</i> isolated from patients with acne vulgaris to zinc ascorbate and antibiotics. <i>Clinical, Cosmetic and Investigational Dermatology</i> , 2011, 4, 161.	0.8	11

#	ARTICLE	IF	CITATIONS
91	Characterization of Enterococcus Strains Contained in Probiotic Products. Biological and Pharmaceutical Bulletin, 2011, 34, 1469-1473.	0.6	19
92	ç—...é™çâ†...ã@é«~é»âº æŽYèš è;éçã«ãããã,ç°èçEã†çš,,ç'°âçfèª;æŸ». Japanese Journal of Environmental Infections, 2011, 26, 362-366.	0.6	19
93	Fluoroquinolone Efflux by the Plasmid-Mediated Multidrug Efflux Pump QacB Variant QacBIII in <i>Staphylococcus aureus</i> . Antimicrobial Agents and Chemotherapy, 2010, 54, 4107-4111.	1.4	58
94	Antimicrobial susceptibilities of <i>Propionibacterium acnes</i> isolated from patients with acne vulgaris. Microbiology and Immunology, 2008, 52, 621-624.	0.7	54
95	Characterization of the pTZ2162 encoding multidrug efflux gene qacB from <i>Staphylococcus aureus</i> . Plasmid, 2008, 60, 108-117.	0.4	37
96	Molecular epidemiology and antimicrobial susceptibilities of 273 exfoliative toxin-encoding-gene-positive <i>Staphylococcus aureus</i> isolates from patients with impetigo in Japan. Journal of Medical Microbiology, 2008, 57, 1251-1258.	0.7	53
97	Anti-infectious Effect of S-Benzylisothiourea Compound A22, Which Inhibits the Actin-Like Protein, MreB, in <i>Shigella flexneri</i> . Biological and Pharmaceutical Bulletin, 2008, 31, 1327-1332.	0.6	16
98	Transduction of the Plasmid Encoding Antiseptic Resistance Gene qacB in <i>Staphylococcus aureus</i> . Biological and Pharmaceutical Bulletin, 2007, 30, 1412-1415.	0.6	26
99	Antimicrobial Agent of Susceptibilities and Antiseptic Resistance Gene Distribution among Methicillin-Resistant <i>Staphylococcus aureus</i> Isolates from Patients with Impetigo and Staphylococcal Scalded Skin Syndrome. Journal of Clinical Microbiology, 2006, 44, 2119-2125.	1.8	88