

# Eduardo De Carli

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8570168/publications.pdf>

Version: 2024-02-01

16  
papers

247  
citations

1307366

7  
h-index

1058333

14  
g-index

16  
all docs

16  
docs citations

16  
times ranked

318  
citing authors

#	ARTICLE	IF	CITATIONS
1	Low Adherence to the EAT-Lancet Sustainable Reference Diet in the Brazilian Population: Findings from the National Dietary Survey 2017-2018. <i>Nutrients</i> , 2022, 14, 1187.	1.7	23
2	Unmetabolized folic acid is associated with TNF- $\alpha$ , IL-1 $\beta$ and IL-12 concentrations in a population exposed to mandatory food fortification with folic acid: a cross-sectional population-based study in Sao Paulo, Brazil. <i>European Journal of Nutrition</i> , 2021, 60, 1071-1079.	1.8	7
3	Development and Validation of an Index Based on EAT-Lancet Recommendations: The Planetary Health Diet Index. <i>Nutrients</i> , 2021, 13, 1698.	1.7	57
4	Prevalence of inadequate intake of folate in the post-fortification era: data from the Brazilian National Dietary Surveys 2008-2009 and 2017-2018. <i>British Journal of Nutrition</i> , 2021, , 1-27.	1.2	2
5	Evolução da ingestão de energia e nutrientes no Brasil entre 2008-2009 e 2017-2018. <i>Revista De Saude Publica</i> , 2021, 55, 1-22.	0.7	10
6	Limitações na comparação dos Inquéritos Nacionais de Alimentação de 2008-2009 e 2017-2018. <i>Revista De Saude Publica</i> , 2021, 55, 1-10.	0.7	4
7	Increased adiposity by feeding growing rats a high-fat diet results in iron decompartmentalisation. <i>British Journal of Nutrition</i> , 2020, 123, 1094-1108.	1.2	4
8	Magnesium intake in a Longitudinal Study of Adult Health: associated factors and the main food sources. <i>Ciencia E Saude Coletiva</i> , 2020, 25, 2541-2550.	0.1	0
9	12th IFDC 2017 Special issue - Brazilian Nutrient Intake Evaluation Database: An essential tool for estimating nutrient intake data. <i>Journal of Food Composition and Analysis</i> , 2019, 83, 103286.	1.9	8
10	Increased serum iron in preeclamptic women is likely due to low hepcidin levels. <i>Nutrition Research</i> , 2018, 53, 32-39.	1.3	18
11	Dietary Iron Bioavailability: Agreement between Estimation Methods and Association with Serum Ferritin Concentrations in Women of Childbearing Age. <i>Nutrients</i> , 2018, 10, 650.	1.7	11
12	Association between Serum Unmetabolized Folic Acid Concentrations and Folic Acid from Fortified Foods. <i>Journal of the American College of Nutrition</i> , 2017, 36, 572-578.	1.1	21
13	A Daily Dose of 5 mg Folic Acid for 90 Days Is Associated with Increased Serum Unmetabolized Folic Acid and Reduced Natural Killer Cell Cytotoxicity in Healthy Brazilian Adults. <i>Journal of Nutrition</i> , 2017, 147, 1677-1685.	1.3	48
14	Fructo-oligosaccharides and iron bioavailability in anaemic rats: the effects on iron species distribution, ferroportin-1 expression, crypt bifurcation and crypt cell proliferation in the caecum. <i>British Journal of Nutrition</i> , 2014, 112, 1286-1295.	1.2	32
15	Short-term dietary magnesium restriction lowers spleen iron concentrations in growing rats fed a high-fat diet. <i>LWT - Food Science and Technology</i> , 2014, 59, 1298-1303.	2.5	2
16	Biomarkers of Fruit Intake Using a Targeted Metabolomics Approach: An Observational Cross-sectional Analysis of the ELSA-Brasil Study. <i>Journal of Nutrition</i> , 0, , .	1.3	0