Melek Sacan

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8570909/melek-sacan-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

676 16 25 39 g-index h-index citations papers 748 4.42 39 5.4 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
39	Chemometric Modeling of Algal Toxicity 2021 , 275-291		
38	Assessments of Algal Toxicity and PBT Behaviour of Pesticides with No Eco-toxicological Data: Predictive Ability of QSA/(T)R Models. <i>Molecular Informatics</i> , 2019 , 38, e1800137	3.8	3
37	Predicting Cytotoxicity and Enzymatic Activity of Diverse Chemicals Using Goldfish Scale Tissue and Topminnow Hepatoma Cell Line-based Data. <i>Molecular Informatics</i> , 2019 , 38, e1800127	3.8	2
36	Revisiting fish toxicity of active pharmaceutical ingredients: Mechanistic insights from integrated ligand-/structure-based assessments on acetylcholinesterase. <i>Ecotoxicology and Environmental Safety</i> , 2019 , 170, 548-558	7	5
35	Understanding the toxic potencies of xenobiotics inducing TCDD/TCDF-like effects. <i>SAR and QSAR in Environmental Research</i> , 2018 , 29, 117-131	3.5	7
34	Toxicity of contaminants of emerging concern to Dugesia japonica: QSTR modeling and toxicity relationship with Daphnia magna. <i>Journal of Hazardous Materials</i> , 2018 , 351, 20-28	12.8	20
33	A multipronged QSAR approach to predict algal low-toxic-effect concentrations of substituted phenols and anilines. <i>Journal of Hazardous Materials</i> , 2018 , 344, 893-901	12.8	15
32	On the prediction of cytotoxicity of diverse chemicals for topminnow (Poeciliopsis lucida) hepatoma cell line, PLHC-1. <i>SAR and QSAR in Environmental Research</i> , 2018 , 29, 675-691	3.5	O
31	Bacterial cellulose production by Komagataeibacter hansenii using algae-based glucose. <i>Environmental Science and Pollution Research</i> , 2017 , 24, 11154-11162	5.1	23
30	On the aquatic toxicity of substituted phenols to Chlorella vulgaris: QSTR with an extended novel data set and interspecies models. <i>Journal of Hazardous Materials</i> , 2017 , 339, 122-130	12.8	36
29	Renewable fuels from pyrolysis of Dunaliella tertiolecta: An alternative approach to biochemical conversions of microalgae. <i>Energy</i> , 2017 , 120, 907-914	7.9	35
28	Impact of geometry optimization methods on QSAR modelling: A case study for predicting human serum albumin binding affinity. <i>SAR and QSAR in Environmental Research</i> , 2017 , 28, 491-509	3.5	9
27	An in silico algal toxicity model with a wide applicability potential for industrial chemicals and pharmaceuticals. <i>Environmental Toxicology and Chemistry</i> , 2017 , 36, 1012-1019	3.8	9
26	An in silico approach to cytotoxicity of pharmaceuticals and personal care products on the rainbow trout liver cell line RTL-W1. <i>Environmental Toxicology and Chemistry</i> , 2017 , 36, 1162-1169	3.8	6
25	Single and mixture toxicity of pharmaceuticals and chlorophenols to freshwater algae Chlorella vulgaris. <i>Ecotoxicology and Environmental Safety</i> , 2016 , 129, 189-98	7	86
24	QSAR models for antioxidant activity of new coumarin derivatives. <i>SAR and QSAR in Environmental Research</i> , 2015 , 26, 721-37	3.5	9
23	Comparative performance of descriptors in a multiple linear and Kriging models: a case study on the acute toxicity of organic chemicals to algae. <i>Environmental Science and Pollution Research</i> , 2014 , 21, 11924-32	5.1	4

(2001-2014)

22	Molecular structure-adsorption study on current textile dyes. <i>SAR and QSAR in Environmental Research</i> , 2014 , 25, 983-98	3.5	3	
21	Assessment and modeling of the novel toxicity data set of phenols to Chlorella vulgaris. <i>Ecotoxicology and Environmental Safety</i> , 2013 , 90, 61-8	7	24	
20	Quantitative structure-activity relationships (QSARs) using the novel marine algal toxicity data of phenols. <i>Journal of Molecular Graphics and Modelling</i> , 2012 , 38, 90-100	2.8	30	
19	First toxicity data of chlorophenols on marine alga Dunaliella tertiolecta: correlation of marine algal toxicity with hydrophobicity and interspecies toxicity relationships. <i>Environmental Toxicology and Chemistry</i> , 2012 , 31, 1113-20	3.8	30	
18	QSTR modelling of the acute toxicity of pharmaceuticals to fish. <i>SAR and QSAR in Environmental Research</i> , 2012 , 23, 297-310	3.5	25	
17	Modelling the relative toxicity of metals on respiration of nitrifiers using ion characteristics. <i>SAR</i> and <i>QSAR in Environmental Research</i> , 2009 , 20, 727-40	3.5	1	
16	Water quality and heavy metal monitoring in water and sediment samples of the KMBkmece Lagoon, Turkey (2002-2003). <i>Environmental Monitoring and Assessment</i> , 2009 , 151, 345-62	3.1	17	
15	Merits of ozonation and catalytic ozonation pre-treatment in the algal treatment of pulp and paper mill effluents. <i>Journal of Environmental Management</i> , 2007 , 85, 918-26	7.9	35	
14	Interaction between a synthetic dye bath and selenium in their toxicity to Dunaliella tertiolecta under two light intensities. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2007 , 78, 142-6	2.7	4	
13	Exposure of Dunaliella tertiolecta to lead and aluminum: toxicity and effects on ultrastructure. <i>Biological Trace Element Research</i> , 2007 , 120, 264-72	4.5	53	
12	QSPR analysis of the toxicity of aromatic compounds to the algae (Scenedesmus obliquus). <i>Chemosphere</i> , 2007 , 68, 695-702	8.4	11	
11	A case study on algal response to raw and treated effluents from an aluminum plating plant and a pharmaceutical plant. <i>Ecotoxicology and Environmental Safety</i> , 2006 , 64, 234-43	7	16	
10	Investigation on the aromaticity of 1,3,4-thiadiazole-2-thione and its oxygen analogs including their tautomeric forms. <i>Computational and Theoretical Chemistry</i> , 2005 , 726, 233-243		17	
9	Physico-chemical properties of PCDD/PCDFs and phthalate esters. <i>SAR and QSAR in Environmental Research</i> , 2005 , 16, 443-59	3.5	22	
8	QSPR study on the bioconcentration factors of nonionic organic compounds in fish by characteristic root index and semiempirical molecular descriptors. <i>Journal of Chemical Information and Computer Sciences</i> , 2004 , 44, 985-92		24	
7	Bioaccumulation of Aluminium in Dunaliella tertiolecta in Natural Seawater: Aluminium?Metal (Cu, Pb, Se) Interactions and Influence of pH. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2001 , 66, 214-221	2.7	7	
6	Bioaccumulation of aluminium in Dunaliella tertiolecta in natural seawater: aluminium-metal (Cu, Pb, Se) interactions and influence of pH. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2001 , 66, 214-21	2.7	6	
5	Homogenous and heterogenous advanced oxidation of two commercial reactive dyes. <i>Environmental Technology (United Kingdom)</i> , 2001 , 22, 813-22	2.6	49	

4	Laboratory bioaccumulation of copper, lead and selenium in the marine alga dunaliella tertiolecta metal pair situation. <i>Toxicological and Environmental Chemistry</i> , 2000 , 76, 17-27	1.4	3
3	Estimation of liquid vapor pressures for low-volatility environmental chemicals. <i>Chemosphere</i> , 1998 , 36, 451-460	8.4	4
2	Prediction of the soil sorption coefficient of organic pollutants by the characteristic root index model. <i>Chemosphere</i> , 1996 , 32, 1993-2001	8.4	13
1	Application of the characteristic root index model to the estimation of N-octanol/water partition coefficients. polychlorinated biphenyls. <i>Chemosphere</i> , 1995 , 30, 39-50	8.4	13