

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8576924/per-s-stal-publications-by-citations.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

81 papers	4,821 citations	27 h-index	69 g-index
86 ext. papers	6,133 ext. citations	5.3 avg, IF	5.63 L-index

#	Paper	IF	Citations
81	Fibrosis stage is the strongest predictor for disease-specific mortality in NAFLD after up to 33 years of follow-up. <i>Hepatology</i> , 2015 , 61, 1547-54	11.2	1158
80	Increased risk of mortality by fibrosis stage in nonalcoholic fatty liver disease: Systematic review and meta-analysis. <i>Hepatology</i> , 2017 , 65, 1557-1565	11.2	796
79	Decreased survival of subjects with elevated liver function tests during a 28-year follow-up. <i>Hepatology</i> , 2010 , 51, 595-602	11.2	537
78	Fibrosis stage but not NASH predicts mortality and time to development of severe liver disease in biopsy-proven NAFLD. <i>Journal of Hepatology</i> , 2017 , 67, 1265-1273	13.4	423
77	Cancer risk in patients with hereditary hemochromatosis and in their first-degree relatives. <i>Gastroenterology</i> , 2003 , 125, 1733-41	13.3	203
76	A risk for hepatocellular carcinoma persists long-term after sustained virologic response in patients with hepatitis C-associated liver cirrhosis. <i>Clinical Infectious Diseases</i> , 2013 , 57, 230-6	11.6	182
75	Risk for development of severe liver disease in lean patients with nonalcoholic fatty liver disease: A long-term follow-up study. <i>Hepatology Communications</i> , 2018 , 2, 48-57	6	106
74	Liver fibrosis in non-alcoholic fatty liver disease - diagnostic challenge with prognostic significance. <i>World Journal of Gastroenterology</i> , 2015 , 21, 11077-87	5.6	82
73	Iron increases ethanol toxicity in rat liver. <i>Journal of Hepatology</i> , 1993 , 17, 108-15	13.4	65
72	Overweight in late adolescence predicts development of severe liver disease later in life: A 39years follow-up study. <i>Journal of Hepatology</i> , 2016 , 65, 363-8	13.4	48
71	Liver transplantation for erythropoietic protoporphyria in Europe. <i>Liver Transplantation</i> , 2011 , 17, 1021-4	4.5	48
70	Hepatotoxicity induced by iron overload and alcohol. Studies on the role of chelatable iron, cytochrome P450 2E1 and lipid peroxidation. <i>Journal of Hepatology</i> , 1996 , 25, 538-46	13.4	46
69	Application of hepatocellular carcinoma surveillance in a European setting. What can we learn from clinical practice?. <i>Liver International</i> , 2015 , 35, 1862-71	7.9	44
68	Cardiovascular risk factors in non-alcoholic fatty liver disease. <i>Liver International</i> , 2019 , 39, 197-204	7.9	42
67	Quantitative determination of 8-hydroxy-2'-deoxyguanosine in human urine by isotope dilution mass spectrometry: normal levels in hemochromatosis. <i>Free Radical Biology and Medicine</i> , 1999 , 26, 129-35	7.8	39
66	Accuracy of Noninvasive Scoring Systems in Assessing Risk of Death and Liver-Related Endpoints in Patients With Nonalcoholic Fatty Liver Disease. <i>Clinical Gastroenterology and Hepatology</i> , 2019 , 17, 1148-1156.e4	6.9	38
65	Perfusion computed tomography for detection of hepatocellular carcinoma in patients with liver cirrhosis. <i>European Radiology</i> , 2015 , 25, 3123-32	8	38

64	Evaluation of genome-wide loci of iron metabolism in hereditary hemochromatosis identifies PCSK7 as a host risk factor of liver cirrhosis. <i>Human Molecular Genetics</i> , 2014 , 23, 3883-90	5.6	36
63	Characteristics and outcome of hepatocellular carcinoma in patients with NAFLD without cirrhosis. <i>Liver International</i> , 2019 , 39, 1098-1108	7.9	35
62	Low to moderate lifetime alcohol consumption is associated with less advanced stages of fibrosis in non-alcoholic fatty liver disease. <i>Scandinavian Journal of Gastroenterology</i> , 2017 , 52, 159-165	2.4	35
61	Liver cell damage and lysosomal iron storage in patients with idiopathic hemochromatosis. A light and electron microscopic study. <i>Journal of Hepatology</i> , 1990 , 11, 172-80	13.4	35
60	Iron as a hepatotoxin. <i>Digestive Diseases</i> , 1995 , 13, 205-22	3.2	34
59	The effects of dietary iron on initiation and promotion in chemical hepatocarcinogenesis. <i>Hepatology</i> , 1995 , 21, 521-8	11.2	34
58	Retained NK Cell Phenotype and Functionality in Non-alcoholic Fatty Liver Disease. <i>Frontiers in Immunology</i> , 2019 , 10, 1255	8.4	33
57	Expression of iron regulatory genes in a rat model of hepatocellular carcinoma. <i>Liver International</i> , 2006 , 26, 976-85	7.9	33
56	Elevated serum ferritin is associated with increased mortality in non-alcoholic fatty liver disease after 16 years of follow-up. <i>Liver International</i> , 2016 , 36, 1688-1695	7.9	31
55	Dietary iron overload inhibits carbon tetrachloride-induced promotion in chemical hepatocarcinogenesis: effects on cell proliferation, apoptosis, and antioxidation. <i>Journal of Hepatology</i> , 1999 , 30, 689-98	13.4	29
54	Effects on contralateral muscles after unilateral electrical muscle stimulation and exercise. <i>PLoS ONE</i> , 2012 , 7, e52230	3.7	25
53	A novel mutation in the biliverdin reductase-A gene combined with liver cirrhosis results in hyperbiliverdinaemia (green jaundice). <i>Liver International</i> , 2009 , 29, 1116-24	7.9	25
52	Structure and liver cell expression pattern of the HFE gene in the rat. <i>Journal of Hepatology</i> , 2003 , 39, 308-14	13.4	25
51	Histologic Scores for Fat and Fibrosis Associate With Development of Type 2 Diabetes in Patients With Nonalcoholic Fatty Liver Disease. <i>Clinical Gastroenterology and Hepatology</i> , 2017 , 15, 1461-1468	6.9	24
50	Nonalcoholic fatty liver disease is an increasing indication for liver transplantation in the Nordic countries. <i>Liver International</i> , 2018 , 38, 2082-2090	7.9	24
49	SAF score and mortality in NAFLD after up to 41 years of follow-up. <i>Scandinavian Journal of Gastroenterology</i> , 2017 , 52, 87-91	2.4	24
48	Microvesicular fat, inter cellular adhesion molecule-1 and regulatory T-lymphocytes are of importance for the inflammatory process in livers with non-alcoholic steatohepatitis. <i>Apmis</i> , 2011 , 119, 412-20	3.4	23
47	Expression of autoantibodies to specific cytochromes P450 in a case of disulfiram hepatitis. <i>Journal of Hepatology</i> , 1998 , 29, 819-25	13.4	23

46	Abnormal mitochondria organization and oxidative activity in the palate muscles of long-term snorers with obstructive sleep apnea. <i>Respiration</i> , 2012 , 83, 407-17	3.7	22
45	In Vivo Drug Delivery Performance of Lipiodol-Based Emulsion or Drug-Eluting Beads in Patients with Hepatocellular Carcinoma. <i>Molecular Pharmaceutics</i> , 2017 , 14, 448-458	5.6	20
44	IGFBP-1 and IGF-I as markers for advanced fibrosis in NAFLD - a pilot study. <i>Scandinavian Journal of Gastroenterology</i> , 2017 , 52, 1427-1434	2.4	18
43	Uncoordinated expression of myosin heavy chains and myosin-binding protein C isoforms in human extraocular muscles. <i>Investigative Ophthalmology and Visual Science</i> , 2006 , 47, 4188-93		17
42	Treatment of NAFLD with intermittent calorie restriction or low-carb high-fat diet - a randomised controlled trial. <i>JHEP Reports</i> , 2021 , 3, 100256	10.3	17
41	Increased mortality risk in patients with phenotypic hereditary hemochromatosis but not in their first-degree relatives. <i>Gastroenterology</i> , 2009 , 137, 1301-9	13.3	15
40	Quality of life as a prognostic factor for survival in hepatocellular carcinoma. <i>Liver International</i> , 2018 , 38, 885-894	7.9	14
39	Hepcidin levels correlate to liver iron content, but not steatohepatitis, in non-alcoholic fatty liver disease. <i>BMC Gastroenterology</i> , 2018 , 18, 78	3	14
38	FTY720 (Fingolimod) sensitizes hepatocellular carcinoma cells to sorafenib-mediated cytotoxicity. <i>Pharmacology Research and Perspectives</i> , 2015 , 3, e00171	3.1	14
37	Liver transplantation of patients with cryptogenic cirrhosis: clinical characteristics and outcome. <i>Scandinavian Journal of Gastroenterology</i> , 2010 , 45, 60-9	2.4	14
36	Regulatory effects of tumor necrosis factor-alpha and interleukin-6 on HAMP expression in iron loaded rat hepatocytes. <i>Journal of Hepatology</i> , 2006 , 44, 544-51	13.4	14
35	Alcohol consumption in patients with primary sclerosing cholangitis. <i>World Journal of Gastroenterology</i> , 2012 , 18, 3105-11	5.6	14
34	Steatohepatitis Is Not Associated with an Increased Risk for Fibrosis Progression in Nonalcoholic Fatty Liver Disease. <i>Gastroenterology Research and Practice</i> , 2018 , 2018, 1942648	2	13
33	Impact of sorafenib dosing on outcome from the European patient subset of the GIDEON study. <i>Future Oncology</i> , 2015 , 11, 2553-62	3.6	12
32	Reverse lectin ELISA for detecting fucosylated forms of N-acid glycoprotein associated with hepatocellular carcinoma. <i>PLoS ONE</i> , 2017 , 12, e0173897	3.7	12
31	Development of Serum Marker Models to Increase Diagnostic Accuracy of Advanced Fibrosis in Nonalcoholic Fatty Liver Disease: The New LINKI Algorithm Compared with Established Algorithms. <i>PLoS ONE</i> , 2016 , 11, e0167776	3.7	12
30	Transient liver elastography in normal pregnancy - a longitudinal cohort study. <i>Scandinavian Journal of Gastroenterology</i> , 2019 , 54, 761-765	2.4	11
29	DNA adducts in normal colonic mucosa from healthy controls and patients with colon polyps and colorectal carcinomas. <i>Mutagenesis</i> , 2010 , 25, 499-504	2.8	11

28	Health check-ups and family screening allow detection of hereditary hemochromatosis with less advanced liver fibrosis and survival comparable with the general population. <i>Scandinavian Journal of Gastroenterology</i> , 2011 , 46, 1118-26	2.4	11
27	Validation of a competitive ELISA assay for the quantification of human serum hepcidin. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2015 , 75, 652-8	2	11
26	Triple Arterial Phase CT of the Liver with Radiation Dose Equivalent to That of Single Arterial Phase CT: Initial Experience. <i>Radiology</i> , 2018 , 289, 111-118	20.5	10
25	Comparison of lipiodol infusion and drug-eluting beads transarterial chemoembolization of hepatocellular carcinoma in a real-life setting. <i>Scandinavian Journal of Gastroenterology</i> , 2019 , 54, 905-912	2.4	10
24	Attenuated liver fibrosis after bile duct ligation and defective hepatic stellate cell activation in neural cell adhesion molecule knockout mice. <i>Liver International</i> , 2011 , 31, 630-41	7.9	10
23	Studies on genotoxic effects of iron overload and alcohol in an animal model of hepatocarcinogenesis. <i>Journal of Hepatology</i> , 1997 , 27, 562-71	13.4	10
22	Statins and Angiotensin-Converting Enzyme Inhibitors are Associated with Reduced Mortality and Morbidity in Chronic Liver Disease. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2018 , 122, 104-110	3.1	9
21	Iron-regulatory gene expression during liver regeneration. <i>Scandinavian Journal of Gastroenterology</i> , 2012 , 47, 591-600	2.4	7
20	Sorafenib prolongs liver regeneration after hepatic resection in rats. <i>Journal of Surgical Research</i> , 2013 , 184, 847-54	2.5	6
19	Effects of dietary iron overload on progression in chemical hepatocarcinogenesis. <i>Liver International</i> , 1999 , 19, 326-34	7.9	5
18	Health Care Costs of Patients With Biopsy-Confirmed Nonalcoholic Fatty Liver Disease Are Nearly Twice Those of Matched Controls. <i>Clinical Gastroenterology and Hepatology</i> , 2020 , 18, 1592-1599.e8	6.9	5
17	A Dynamic Aspartate-to-Alanine Aminotransferase Ratio Provides Valid Predictions of Incident Severe Liver Disease. <i>Hepatology Communications</i> , 2021 , 5, 1021-1035	6	5
16	Bilateral muscle fiber and nerve influences by TNF-alpha in response to unilateral muscle overuse - studies on TNF receptor expressions. <i>BMC Musculoskeletal Disorders</i> , 2017 , 18, 498	2.8	4
15	Ubiquinone, Oxidative Stress, and Liver Carcinogenesis. <i>Modern Nutrition</i> , 2000 , 317-329		4
14	Morbidity, risk of cancer and mortality in 3645 HFE mutations carriers. <i>Liver International</i> , 2021 , 41, 545-553	5.3	4
13	Non-alcoholic fatty liver disease does not increase dementia risk although histology data might improve risk prediction. <i>JHEP Reports</i> , 2021 , 3, 100218	10.3	3
12	Pembrolizumab (pembro) monotherapy for previously untreated advanced hepatocellular carcinoma (HCC): Phase II KEYNOTE-224 study.. <i>Journal of Clinical Oncology</i> , 2021 , 39, 297-297	2.2	3
11	Neutrophil extracellular traps in patients with liver cirrhosis and hepatocellular carcinoma. <i>Scientific Reports</i> , 2021 , 11, 18025	4.9	3

10	Alcohol and drug use prior to liver transplantation: more common than expected in patients with non-alcoholic liver disease. <i>Scandinavian Journal of Gastroenterology</i> , 2019 , 54, 1146-1154	2.4	2
9	Liquid Biopsy in Hepatocellular Carcinoma: Opportunities and Challenges for Immunotherapy. <i>Cancers</i> , 2021 , 13,	6.6	2
8	A personalized treatment program in persons with type 2 diabetes is associated with a reduction in liver steatosis. <i>European Journal of Gastroenterology and Hepatology</i> , 2021 , 33, 1420-1426	2.2	1
7	Variation in textural parameters of hepatic lesions during contrast medium injection. <i>Acta Radiologica</i> , 2021 , 62, 1317-1323	2	1
6	Macrophage Markers Do Not Add to the Prediction of Liver Fibrosis by Transient Elastography in Patients With Metabolic Associated Fatty Liver Disease. <i>Frontiers in Medicine</i> , 2020 , 7, 616212	4.9	1
5	Serum levels of endotrophin are associated with nonalcoholic steatohepatitis. <i>Scandinavian Journal of Gastroenterology</i> , 2021 , 56, 437-442	2.4	1
4	Reply to "Comment on Tn Vivo Drug Delivery Performance of Lipiodol-Based Emulsion or Drug-Eluting Beads in Patients with Hepatocellular Carcinoma". <i>Molecular Pharmaceutics</i> , 2018 , 15, 336-340	5.6	1
3	Risk of hepatocellular carcinoma in hepatitis B and D virus co-infected patients: A systematic review and meta-analysis of longitudinal studies. <i>Journal of Viral Hepatitis</i> , 2021 , 28, 1431-1442	3.4	1
2	Reply. <i>Hepatology</i> , 2016 , 64, 310-1	11.2	
1	Reply to: "Reduced steatosis and weight as a result of specific diets or the dietitian themselves". <i>JHEP Reports</i> , 2021 , 3, 100366	10.3	