Luminita Pricop

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8578928/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Secukinumab, a human anti-interleukin-17A monoclonal antibody, in patients with psoriatic arthritis (FUTURE 2): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet, The, 2015, 386, 1137-1146.	13.7	722
2	Secukinumab Inhibition of Interleukin-17A in Patients with Psoriatic Arthritis. New England Journal of Medicine, 2015, 373, 1329-1339.	27.0	629
3	Secukinumab improves active psoriatic arthritis symptoms and inhibits radiographic progression: primary results from the randomised, double-blind, phase III FUTURE 5 study. Annals of the Rheumatic Diseases, 2018, 77, annrheumdis-2017-212687.	0.9	193
4	Efficacy and safety of secukinumab administration by autoinjector in patients with psoriatic arthritis: results from a randomized, placebo-controlled trial (FUTURE 3). Arthritis Research and Therapy, 2018, 20, 47.	3.5	117
5	Secukinumab for psoriatic arthritis: comparative effectiveness versus licensed biologics/apremilast: a network meta-analysis. Journal of Comparative Effectiveness Research, 2018, 7, 1107-1123.	1.4	35
6	Secukinumab Immunogenicity over 52 Weeks in Patients with Psoriatic Arthritis and Ankylosing Spondylitis. Journal of Rheumatology, 2020, 47, 539-547.	2.0	16
7	Matching-adjusted indirect comparison: secukinumab versus infliximab in biologic-naive patients with psoriatic arthritis. Journal of Comparative Effectiveness Research, 2019, 8, 497-510.	1.4	15
8	Clinically relevant patient clusters identified by machine learning from the clinical development programme of secukinumab in psoriatic arthritis. RMD Open, 2021, 7, e001845.	3.8	11
9	Comparative effectiveness of secukinumab and etanercept in biologic-naÃ ⁻ ve patients with psoriatic arthritis assessed by matching-adjusted indirect comparison. European Journal of Rheumatology, 2019, 6, 113-121.	0.6	5
10	256 Secukinumab provides sustained improvements in the signs and symptoms of active psoriatic arthritis: 4-year results from the Phase 3 FUTURE 2 study. Rheumatology, 2019, 58, .	1.9	3
11	257 Secukinumab demonstrates a consistent safety profile with up to 5 years' treatment in patients with psoriatic arthritis and moderate-to-severe plaque psoriasis: updated pooled safety analyses. Rheumatology, 2019, 58, .	1.9	1
12	263 Secukinumab, a fully human anti-interleukin-17A monoclonal antibody, exhibits low immunogenicity in patients with psoriatic arthritis and ankylosing spondylitis during a 52week treatment period. Rheumatology, 2019, 58, .	1.9	0
13	P250 Effect of secukinumab on radiographic progression through 2 years in patients with active PsA: end-of-study results from a Phase 3 study. Rheumatology, 2020, 59, .	1.9	0
14	OA37 Secukinumab treatment in children and adolescents with enthesitis-related arthritis and juvenile psoriatic arthritis: efficacy and safety results from a Phase 3 study. Rheumatology, 2022, 61, .	1.9	0