

Jan-Peter Sowa

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8590340/jan-peter-sowa-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

62
papers

2,107
citations

23
h-index

45
g-index

66
ext. papers

2,561
ext. citations

5.4
avg, IF

4.62
L-index

#	Paper	IF	Citations
62	Transcriptome-Wide Analysis of Human Liver Reveals Age-Related Differences in the Expression of Select Functional Gene Clusters and Evidence for a PPP1R10-Governed Aging CascadeS. <i>Pharmaceutics</i> , 2021 , 13,	6.4	1
61	Association of cell death mechanisms and fibrosis in visceral white adipose tissue with pathological alterations in the liver of morbidly obese patients with NAFLD. <i>Adipocyte</i> , 2021 , 10, 558-573	3.2	4
60	Lipoprotein and Metabolic Profiles Indicate Similar Cardiovascular Risk of Liver Steatosis and NASH. <i>Digestion</i> , 2021 , 102, 671-681	3.6	4
59	Healthcare resource utilization and costs among nonalcoholic fatty liver disease patients in Germany. <i>Annals of Translational Medicine</i> , 2021 , 9, 615	3.2	2
58	GALAD Score Detects Early-Stage Hepatocellular Carcinoma in a European Cohort of Chronic Hepatitis B and C Patients. <i>Pharmaceutics</i> , 2021 , 14,	5.2	7
57	Hepatokines and adipokines in NASH-related hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2021 , 74, 442-457	13.4	29
56	Transaminase concentrations cannot separate NAFL and NASH in morbidly obese patients irrespective of histological algorithm. <i>Digestive Diseases</i> , 2021 ,	3.2	1
55	GALAD Score Detects Early Hepatocellular Carcinoma in an International Cohort of Patients With Nonalcoholic Steatohepatitis. <i>Clinical Gastroenterology and Hepatology</i> , 2020 , 18, 728-735.e4	6.9	73
54	Significance of Simple Steatosis: An Update on the Clinical and Molecular Evidence. <i>Cells</i> , 2020 , 9,	7.9	12
53	Patterns and predictors of mortality and disease progression among patients with non-alcoholic fatty liver disease. <i>Alimentary Pharmacology and Therapeutics</i> , 2020 , 52, 1185-1194	6.1	14
52	Liver parameters as part of a non-invasive model for prediction of all-cause mortality after myocardial infarction. <i>Archives of Medical Science</i> , 2020 , 16, 71-80	2.9	5
51	L-Ornithine L-Aspartate (LOLA) as a Novel Approach for Therapy of Non-alcoholic Fatty Liver Disease. <i>Drugs</i> , 2019 , 79, 39-44	12.1	16
50	Non-invasive assessment of NAFLD as systemic disease-A machine learning perspective. <i>PLoS ONE</i> , 2019 , 14, e0214436	3.7	30
49	The virtual doctor: An interactive clinical-decision-support system based on deep learning for non-invasive prediction of diabetes. <i>Artificial Intelligence in Medicine</i> , 2019 , 100, 101706	7.4	35
48	Three Cases of Alcohol-Induced Acute-On-Chronic Liver Failure With Successful Support by Adipose-Derived Stem Cells. <i>Clinical and Translational Gastroenterology</i> , 2019 , 10, e00095	4.2	4
47	How to Define Acute Liver Failure Patients with Pre-Existing Liver Disease without Signs of Cirrhosis. <i>Digestive Diseases</i> , 2019 , 37, 147-154	3.2	7
46	Nonalcoholic-Fatty-Liver-Disease and Nonalcoholic Steatohepatitis: Successful Development of Pharmacological Treatment Will Depend on Translational Research. <i>Digestion</i> , 2019 , 100, 79-85	3.6	14

45	SPARC expression is associated with hepatic injury in rodents and humans with non-alcoholic fatty liver disease. <i>Scientific Reports</i> , 2018 , 8, 725	4.9	11
44	Low transferrin and high ferritin concentrations are associated with worse outcome in acute liver failure. <i>Liver International</i> , 2017 , 37, 1032-1041	7.9	18
43	Acid sphingomyelinase deficiency in Western diet-fed mice protects against adipocyte hypertrophy and diet-induced liver steatosis. <i>Molecular Metabolism</i> , 2017 , 6, 416-427	8.8	6
42	Krppel-like factor 6 is a transcriptional activator of autophagy in acute liver injury. <i>Scientific Reports</i> , 2017 , 7, 8119	4.9	19
41	Rotational thromboelastometry can detect factor XIII deficiency and bleeding diathesis in patients with cirrhosis. <i>Liver International</i> , 2017 , 37, 562-568	7.9	30
40	Management of acute-on-chronic liver failure: rotational thromboelastometry may reduce substitution of coagulation factors in liver cirrhosis. <i>Gut</i> , 2016 , 65, 357-8	19.2	44
39	NASH Cirrhosis - the New Burden in Liver Transplantation: How Should It Be Managed?. <i>Visceral Medicine</i> , 2016 , 32, 234-238	2.4	18
38	Cell death mechanisms in human chronic liver diseases: a far cry from clinical applicability. <i>Clinical Science</i> , 2016 , 130, 2121-2138	6.5	11
37	Macrophage Depletion Attenuates Extracellular Matrix Deposition and Ductular Reaction in a Mouse Model of Chronic Cholangiopathies. <i>PLoS ONE</i> , 2016 , 11, e0162286	3.7	19
36	Economic growth leads to increase of obesity and associated hepatocellular carcinoma in developing countries. <i>Annals of Hepatology</i> , 2016 , 15, 662-72	3.1	32
35	Epidemiology of nonalcoholic steatohepatitis and hepatocellular carcinoma. <i>Clinical Liver Disease</i> , 2016 , 8, 119-122	2.2	18
34	Compensation of feature selection biases accompanied with improved predictive performance for binary classification by using a novel ensemble feature selection approach. <i>BioData Mining</i> , 2016 , 9, 36	4.3	30
33	Human Ex-Vivo Liver Model for Acetaminophen-induced Liver Damage. <i>Scientific Reports</i> , 2016 , 6, 31916	4.9	9
32	In Acute Myocardial Infarction Liver Parameters Are Associated With Stenosis Diameter. <i>Medicine (United States)</i> , 2016 , 95, e2807	1.8	13
31	Acute Liver Failure - It's Just a Matter of Cell Death. <i>Digestive Diseases</i> , 2016 , 34, 423-8	3.2	12
30	Vitamin D counteracts fibrogenic TGF-β signalling in human hepatic stellate cells both receptor-dependently and independently. <i>Gut</i> , 2015 , 64, 791-9	19.2	89
29	Self-reports on symptoms of alcohol abuse: liver transplant patients versus rehabilitation therapy patients. <i>Progress in Transplantation</i> , 2015 , 25, 203-9	1.1	4
28	Normal liver enzymes are correlated with severity of metabolic syndrome in a large population based cohort. <i>Scientific Reports</i> , 2015 , 5, 13058	4.9	44

27	Higher Thyroid-Stimulating Hormone, Triiodothyronine and Thyroxine Values Are Associated with Better Outcome in Acute Liver Failure. <i>PLoS ONE</i> , 2015 , 10, e0132189	3.7	9
26	Liver Injury Indicating Fatty Liver but Not Serologic NASH Marker Improves under Metformin Treatment in Polycystic Ovary Syndrome. <i>International Journal of Endocrinology</i> , 2015 , 2015, 254169	2.7	7
25	Etiology, outcome and prognostic factors of childhood acute liver failure in a German Single Center. <i>Annals of Hepatology</i> , 2015 , 14, 722-8	3.1	8
24	Adipocyte cell size, free fatty acids and apolipoproteins are associated with non-alcoholic liver injury progression in severely obese patients. <i>Metabolism: Clinical and Experimental</i> , 2014 , 63, 1542-52	12.7	75
23	Mini-laparoscopy guided liver biopsy increases diagnostic accuracy in acute liver failure. <i>Digestion</i> , 2014 , 90, 240-7	3.6	15
22	From bedside to bench and back again-molecular mechanisms in acute liver failure. <i>Frontiers in Physiology</i> , 2014 , 5, 18	4.6	
21	Aflatoxin exposure may not play a role in liver cancer development in Mongolia. <i>Digestion</i> , 2014 , 89, 2683-7	3.7	11
20	Non-invasive separation of alcoholic and non-alcoholic liver disease with predictive modeling. <i>PLoS ONE</i> , 2014 , 9, e101444	3.7	32
19	Serum sodium based modification of the MELD does not improve prediction of outcome in acute liver failure. <i>BMC Gastroenterology</i> , 2013 , 13, 58	3	11
18	Vascular endothelial growth factor improves liver regeneration and survival after 90% hepatectomy in a rat model of diet-induced steatosis. <i>Digestion</i> , 2013 , 88, 235-42	3.6	8
17	Steatosis does not impair liver regeneration after partial hepatectomy. <i>Laboratory Investigation</i> , 2013 , 93, 20-30	5.9	38
16	Free fatty acids repress small heterodimer partner (SHP) activation and adiponectin counteracts bile acid-induced liver injury in superobese patients with nonalcoholic steatohepatitis. <i>Hepatology</i> , 2013 , 57, 1394-406	11.2	143
15	Fetuin-A mRNA expression is elevated in NASH compared with NAFL patients. <i>Clinical Science</i> , 2013 , 125, 391-400	6.5	44
14	Expression of apoptosis- and vitamin D pathway-related genes in hepatocellular carcinoma. <i>Digestion</i> , 2013 , 87, 176-81	3.6	15
13	Novel algorithm for non-invasive assessment of fibrosis in NAFLD. <i>PLoS ONE</i> , 2013 , 8, e62439	3.7	40
12	Circulating microRNAs: promising candidates serving as novel biomarkers of acute hepatitis. <i>Frontiers in Physiology</i> , 2012 , 3, 476	4.6	30
11	An ex vivo perfusion system emulating in vivo conditions in noncirrhotic and cirrhotic human liver. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2012 , 342, 730-41	4.7	3
10	Deficiency of the promyelocytic leukemia protein fosters hepatitis C-associated hepatocarcinogenesis in mice. <i>PLoS ONE</i> , 2012 , 7, e44474	3.7	8

9	Corticosteroids shift the Toll-like receptor response pattern of primary-isolated murine liver cells from an inflammatory to an anti-inflammatory state. <i>International Immunology</i> , 2011 , 23, 537-44	4.9	24
8	Onset of heart failure determines the hepatic cell death pattern. <i>Annals of Hepatology</i> , 2011 , 10, 174-179.1	9.1	13
7	Non-alcoholic fatty liver disease progresses to hepatocellular carcinoma in the absence of apparent cirrhosis. <i>International Journal of Cancer</i> , 2011 , 128, 2436-43	7.5	325
6	Cytokeratin 18-based modification of the MELD score improves prediction of spontaneous survival after acute liver injury. <i>Journal of Hepatology</i> , 2010 , 53, 639-47	13.4	130
5	Apoptosis is associated with CD36/fatty acid translocase upregulation in non-alcoholic steatohepatitis. <i>Liver International</i> , 2010 , 30, 850-9	7.9	65
4	Acute liver failure is associated with elevated liver stiffness and hepatic stellate cell activation. <i>Hepatology</i> , 2010 , 52, 1008-16	11.2	110
3	Hepatitis B virus suppresses toll-like receptor-mediated innate immune responses in murine parenchymal and nonparenchymal liver cells. <i>Hepatology</i> , 2009 , 49, 1132-40	11.2	247
2	Extent of liver resection modulates the activation of transcription factors and the production of cytokines involved in liver regeneration. <i>World Journal of Gastroenterology</i> , 2008 , 14, 7093-100	5.6	19
1	Creld2 function during unfolded protein response is essential for liver metabolism homeostasis		1