

Igal Ifergan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8591443/publications.pdf>

Version: 2024-02-01

41
papers

5,474
citations

257450

24
h-index

289244

40
g-index

51
all docs

51
docs citations

51
times ranked

8301
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | PLG nanoparticles target fibroblasts and MARCO+ monocytes to reverse multiorgan fibrosis. JCI Insight, 2022, 7, . | 5.0 | 8 |
| 2 | ZEB1 promotes pathogenic Th1 and Th17 cell differentiation in multiple sclerosis. Cell Reports, 2021, 36, 109602. | 6.4 | 22 |
| 3 | Tolerance Induced by Antigen-Loaded PLG Nanoparticles Affects the Phenotype and Trafficking of Transgenic CD4+ and CD8+ T Cells. Cells, 2021, 10, 3445. | 4.1 | 4 |
| 4 | Intravenous Immunomodulatory Nanoparticle Treatment for Traumatic Brain Injury. Annals of Neurology, 2020, 87, 442-455. | 5.3 | 29 |
| 5 | Monocytes prime autoreactive T cells after myocardial infarction. American Journal of Physiology - Heart and Circulatory Physiology, 2020, 318, H116-H123. | 3.2 | 15 |
| 6 | Potential for Targeting Myeloid Cells in Controlling CNS Inflammation. Frontiers in Immunology, 2020, 11, 571897. | 4.8 | 12 |
| 7 | Herpesvirus Entry Mediator Binding Partners Mediate Immunopathogenesis of Ocular Herpes Simplex Virus 1 Infection. MBio, 2020, 11, . | 4.1 | 7 |
| 8 | CRISPR screen in regulatory T cells reveals modulators of Foxp3. Nature, 2020, 582, 416-420. | 27.8 | 141 |
| 9 | Methodology for in vitro Assessment of Human T Cell Activation and Blockade. Bio-protocol, 2020, 10, e3644. | 0.4 | 0 |
| 10 | Peripherally derived T regulatory and Th17 T cells have opposing roles in the pathogenesis of intractable pediatric epilepsy. Journal of Experimental Medicine, 2018, 215, 1169-1186. | 8.5 | 80 |
| 11 | B7-H4 Modulates Regulatory CD4+ T Cell Induction and Function via Ligation of a Semaphorin 3a/Plexin A4/Neuropilin-1 Complex. Journal of Immunology, 2018, 201, 897-907. | 0.8 | 34 |
| 12 | Intravenous immune-modifying nanoparticles as a therapy for spinal cord injury in mice. Neurobiology of Disease, 2017, 108, 73-82. | 4.4 | 48 |
| 13 | Pre-metastatic cancer exosomes induce immune surveillance by patrolling monocytes at the metastatic niche. Nature Communications, 2017, 8, 1319. | 12.8 | 237 |
| 14 | Targeting the GM-CSF receptor for the treatment of CNS autoimmunity. Journal of Autoimmunity, 2017, 84, 1-11. | 6.5 | 53 |
| 15 | Murine Corneal Inflammation and Nerve Damage After Infection With HSV-1 Are Promoted by HVEM and Ameliorated by Immune-Modifying Nanoparticle Therapy. , 2017, 58, 282. | | 19 |
| 16 | Biodegradable antigen-associated PLG nanoparticles tolerize Th2-mediated allergic airway inflammation pre- and postsensitization. Proceedings of the National Academy of Sciences of the United States of America, 2016, 113, 5059-5064. | 7.1 | 78 |
| 17 | Cutting Edge: MicroRNA-223 Regulates Myeloid Dendritic Cell-Driven Th17 Responses in Experimental Autoimmune Encephalomyelitis. Journal of Immunology, 2016, 196, 1455-1459. | 0.8 | 45 |
| 18 | Functional analyses of transmigrated monocyte-derived CD123+ dendritic cells across the inflamed blood-brain barrier endothelium. Journal of Neuroimmunology, 2014, 275, 155. | 2.3 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 19 | Experimental Autoimmune Encephalomyelitis in Mice. <i>Methods in Molecular Biology</i> , 2014, 1304, 145-160. | 0.9 | 58 |
| 20 | Melanoma cell adhesion molecule identifies encephalitogenic T lymphocytes and promotes their recruitment to the central nervous system. <i>Brain</i> , 2012, 135, 2906-2924. | 7.6 | 128 |
| 21 | B Cell-Derived IL-15 Enhances CD8 T Cell Cytotoxicity and Is Increased in Multiple Sclerosis Patients. <i>Journal of Immunology</i> , 2011, 187, 4119-4128. | 0.8 | 59 |
| 22 | The Hedgehog Pathway Promotes Blood-Brain Barrier Integrity and CNS Immune Quiescence. <i>Science</i> , 2011, 334, 1727-1731. | 12.6 | 676 |
| 23 | Role of ninjurin-1 in the migration of myeloid cells to central nervous system inflammatory lesions. <i>Annals of Neurology</i> , 2011, 70, 751-763. | 5.3 | 126 |
| 24 | Central nervous system recruitment of effector memory CD8+ T lymphocytes during neuroinflammation is dependent on $\alpha 4$ integrin. <i>Brain</i> , 2011, 134, 3560-3577. | 7.6 | 112 |
| 25 | Isolation of Human Brain Endothelial Cells and Characterization of Lipid Raft-Associated Proteins by Mass Spectroscopy. <i>Methods in Molecular Biology</i> , 2011, 686, 275-295. | 0.9 | 18 |
| 26 | OR.17. Ninjurin-1 is a Novel Adhesion Molecule of the Blood-brain Barrier Involved in the Recruitment of Monocytes to the Central Nervous System. <i>Clinical Immunology</i> , 2009, 131, S10-S11. | 3.2 | 1 |
| 27 | OR.21. MCAM/CD146 is Expressed by Brain Endothelial Cells and Defines a Unique Effector Memory Lymphocyte Subset Involved in Neuroinflammation. <i>Clinical Immunology</i> , 2009, 131, S12. | 3.2 | 2 |
| 28 | OR.81. Astrocyte-secreted Sonic Hedgehog Supports CNS Anti-inflammatory Activity and Promotes Optimal Human Blood Brain Barrier Functioning. <i>Clinical Immunology</i> , 2009, 131, S34. | 3.2 | 2 |
| 29 | Preferential recruitment of interferon- γ -expressing T _H 17 cells in multiple sclerosis. <i>Annals of Neurology</i> , 2009, 66, 390-402. | 5.3 | 494 |
| 30 | Activation of kinin receptor B1 limits encephalitogenic T lymphocyte recruitment to the central nervous system. <i>Nature Medicine</i> , 2009, 15, 788-793. | 30.7 | 118 |
| 31 | IFN- γ regulates CD73 and adenosine expression at the blood-brain barrier. <i>European Journal of Immunology</i> , 2008, 38, 2718-2726. | 2.9 | 72 |
| 32 | Activated leukocyte cell adhesion molecule promotes leukocyte trafficking into the central nervous system. <i>Nature Immunology</i> , 2008, 9, 137-145. | 14.5 | 358 |
| 33 | The blood-brain barrier induces differentiation of migrating monocytes into Th17-polarizing dendritic cells. <i>Brain</i> , 2008, 131, 785-799. | 7.6 | 169 |
| 34 | Human TH17 lymphocytes promote blood-brain barrier disruption and central nervous system inflammation. <i>Nature Medicine</i> , 2007, 13, 1173-1175. | 30.7 | 1,442 |
| 35 | Human Blood-brain Barrier-associated DCs Originate from Blood Monocytes and Polarize CD4+ Lymphocytes into Th17 or Th1. <i>Clinical Immunology</i> , 2007, 123, S151-S152. | 3.2 | 0 |
| 36 | Cytotoxic Human IL-22-expressing Th17 Lymphocytes Promote Immune Cell Migration Into the Central Nervous System. <i>Clinical Immunology</i> , 2007, 123, S60. | 3.2 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Statins reduce human blood-brain barrier permeability and restrict leukocyte migration: Relevance to multiple sclerosis. <i>Annals of Neurology</i> , 2006, 60, 45-55. | 5.3 | 144 |
| 38 | Microglial Expression of the B7 Family Member B7 Homolog 1 Confers Strong Immune Inhibition: Implications for Immune Responses and Autoimmunity in the CNS. <i>Journal of Neuroscience</i> , 2005, 25, 2537-2546. | 3.6 | 150 |
| 39 | Type 2 Monocyte and Microglia Differentiation Mediated by Glatiramer Acetate Therapy in Patients with Multiple Sclerosis. <i>Journal of Immunology</i> , 2004, 172, 7144-7153. | 0.8 | 187 |
| 40 | Analyses of all matrix metalloproteinase members in leukocytes emphasize monocytes as major inflammatory mediators in multiple sclerosis. <i>Brain</i> , 2003, 126, 2738-2749. | 7.6 | 300 |
| 41 | Allele frequency of three functionally active polymorphisms of the MDR-1 gene in high-risk HIV-negative and HIV-positive Caucasians. <i>Aids</i> , 2002, 16, 2340-2342. | 2.2 | 19 |