

Fangbin Cao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8596033/publications.pdf>

Version: 2024-02-01

41
papers

2,065
citations

279798

23
h-index

276875

41
g-index

41
all docs

41
docs citations

41
times ranked

2369
citing authors

#	ARTICLE	IF	CITATIONS
1	Identification of low grain cadmium accumulation genotypes and its physiological mechanism in maize (<i>Zea mays</i> L.). <i>Environmental Science and Pollution Research</i> , 2022, 29, 20721-20730.	5.3	8
2	Genome-wide association study reveals a genomic region on 5AL for salinity tolerance in wheat. <i>Theoretical and Applied Genetics</i> , 2022, 135, 709-721.	3.6	10
3	An ATP binding cassette transporter HvABCB25 confers aluminum detoxification in wild barley. <i>Journal of Hazardous Materials</i> , 2021, 401, 123371.	12.4	33
4	Metabolome Analysis Revealed the Mechanism of Exogenous Glutathione to Alleviate Cadmium Stress in Maize (<i>Zea mays</i> L.) Seedlings. <i>Plants</i> , 2021, 10, 105.	3.5	23
5	Mechanistic Insights into Potassium-Conferred Drought Stress Tolerance in Cultivated and Tibetan Wild Barley: Differential Osmoregulation, Nutrient Retention, Secondary Metabolism and Antioxidative Defense Capacity. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13100.	4.1	7
6	Genome-Wide Discovery of miRNAs with Differential Expression Patterns in Responses to Salinity in the Two Contrasting Wheat Cultivars. <i>International Journal of Molecular Sciences</i> , 2021, 22, 12556.	4.1	10
7	Silicon regulates the expression of vacuolar H ⁺ -pyrophosphatase 1 and decreases cadmium accumulation in rice (<i>Oryza sativa</i> L.). <i>Chemosphere</i> , 2020, 240, 124907.	8.2	40
8	Overexpression of HvAKT1 improves drought tolerance in barley by regulating root ion homeostasis and ROS and NO signaling. <i>Journal of Experimental Botany</i> , 2020, 71, 6587-6600.	4.8	31
9	The Barley S-Adenosylmethionine Synthetase 3 Gene HvSAMS3 Positively Regulates the Tolerance to Combined Drought and Salinity Stress in Tibetan Wild Barley. <i>Cells</i> , 2020, 9, 1530.	4.1	20
10	Genome-Wide Identification and Characterization of Drought Stress Responsive microRNAs in Tibetan Wild Barley. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2795.	4.1	29
11	Resemblance and Difference of Seedling Metabolic and Transporter Gene Expression in High Tolerance Wheat and Barley Cultivars in Response to Salinity Stress. <i>Plants</i> , 2020, 9, 519.	3.5	18
12	Comparative physiological analysis in the tolerance to salinity and drought individual and combination in two cotton genotypes with contrasting salt tolerance. <i>Physiologia Plantarum</i> , 2019, 165, 155-168.	5.2	46
13	Foliar application of betaine improves water-deficit stress tolerance in barley (<i>Hordeum vulgare</i> L.). <i>Plant Growth Regulation</i> , 2019, 89, 109-118.	3.4	22
14	HvPAA1 Encodes a P-Type ATPase, a Novel Gene for Cadmium Accumulation and Tolerance in Barley (<i>Hordeum vulgare</i> L.). <i>International Journal of Molecular Sciences</i> , 2019, 20, 1732.	4.1	20
15	Response of Tibetan Wild Barley Genotypes to Drought Stress and Identification of Quantitative Trait Loci by Genome-Wide Association Analysis. <i>International Journal of Molecular Sciences</i> , 2019, 20, 791.	4.1	15
16	Differences in physiological and biochemical characteristics in response to single and combined drought and salinity stresses between wheat genotypes differing in salt tolerance. <i>Physiologia Plantarum</i> , 2019, 165, 134-143.	5.2	66
17	Tolerance to Drought, Low pH and Al Combined Stress in Tibetan Wild Barley Is Associated with Improvement of ATPase and Modulation of Antioxidant Defense System. <i>International Journal of Molecular Sciences</i> , 2018, 19, 3553.	4.1	12
18	Effect of combined application of lead, cadmium, chromium and copper on grain, leaf and stem heavy metal contents at different growth stages in rice. <i>Ecotoxicology and Environmental Safety</i> , 2018, 162, 71-76.	6.0	57

#	ARTICLE	IF	CITATIONS
19	Genotypic differences in cadmium transport in developing barley grains. <i>Environmental Science and Pollution Research</i> , 2017, 24, 7009-7015.	5.3	2
20	Alleviation of cadmium toxicity by potassium supplementation involves various physiological and biochemical features in <i>Nicotiana tabacum</i> L.. <i>Acta Physiologiae Plantarum</i> , 2017, 39, 1.	2.1	17
21	Genotypic-dependent effects of N fertilizer, glutathione, silicon, zinc, and selenium on proteomic profiles, amino acid contents, and quality of rice genotypes with contrasting grain Cd accumulation. <i>Functional and Integrative Genomics</i> , 2017, 17, 387-397.	3.5	9
22	Glutathione-induced alleviation of cadmium toxicity in <i>Zea mays</i> . <i>Plant Physiology and Biochemistry</i> , 2017, 119, 240-249.	5.8	34
23	Foliar application of betaine alleviates cadmium toxicity in maize seedlings. <i>Acta Physiologiae Plantarum</i> , 2016, 38, 1.	2.1	15
24	Genotype-dependent effect of exogenous 24-epibrassinolide on chromium-induced changes in ultrastructure and physicochemical traits in tobacco seedlings. <i>Environmental Science and Pollution Research</i> , 2016, 23, 18229-18238.	5.3	54
25	Identification and comparative analysis of the microRNA transcriptome in roots of two contrasting tobacco genotypes in response to cadmium stress. <i>Scientific Reports</i> , 2016, 6, 32805.	3.3	37
26	Physiological and molecular analysis on root growth associated with the tolerance to aluminum and drought individual and combined in Tibetan wild and cultivated barley. <i>Planta</i> , 2016, 243, 973-985.	3.2	22
27	<i>HvEXPB7</i> , a novel β -expansin gene revealed by the root hair transcriptome of Tibetan wild barley, improves root hair growth under drought stress. <i>Journal of Experimental Botany</i> , 2015, 66, 7405-7419.	4.8	94
28	Genotypic differences in photosynthetic performance, antioxidant capacity, ultrastructure and nutrients in response to combined stress of salinity and Cd in cotton. <i>BioMetals</i> , 2015, 28, 1063-1078.	4.1	29
29	Differences in photosynthesis, yield and grain cadmium accumulation as affected by exogenous cadmium and glutathione in the two rice genotypes. <i>Plant Growth Regulation</i> , 2015, 75, 715-723.	3.4	84
30	Secondary metabolism and antioxidants are involved in the tolerance to drought and salinity, separately and combined, in Tibetan wild barley. <i>Environmental and Experimental Botany</i> , 2015, 111, 1-12.	4.2	129
31	N-acetyl-cysteine alleviates Cd toxicity and reduces Cd uptake in the two barley genotypes differing in Cd tolerance. <i>Plant Growth Regulation</i> , 2014, 74, 93-105.	3.4	18
32	Genotypic and environmental variation in cadmium, chromium, lead and copper in rice and approaches for reducing the accumulation. <i>Science of the Total Environment</i> , 2014, 496, 275-281.	8.0	81
33	Genome-wide transcriptome and functional analysis of two contrasting genotypes reveals key genes for cadmium tolerance in barley. <i>BMC Genomics</i> , 2014, 15, 611.	2.8	101
34	Differences in physiological features associated with aluminum tolerance in Tibetan wild and cultivated barleys. <i>Plant Physiology and Biochemistry</i> , 2014, 75, 36-44.	5.8	14
35	Genotypic differences in physiological characteristics in the tolerance to drought and salinity combined stress between Tibetan wild and cultivated barley. <i>Plant Physiology and Biochemistry</i> , 2013, 63, 49-60.	5.8	219
36	Differential changes in grain ultrastructure, amylase, protein and amino acid profiles between Tibetan wild and cultivated barleys under drought and salinity alone and combined stress. <i>Food Chemistry</i> , 2013, 141, 2743-2750.	8.2	66

#	ARTICLE	IF	CITATIONS
37	Comparative study of alleviating effects of GSH, Se and Zn under combined contamination of cadmium and chromium in rice (<i>Oryza sativa</i>). <i>BioMetals</i> , 2013, 26, 297-308.	4.1	50
38	Difference in Yield and Physiological Features in Response to Drought and Salinity Combined Stress during Anthesis in Tibetan Wild and Cultivated Barleys. <i>PLoS ONE</i> , 2013, 8, e77869.	2.5	116
39	Selenium reduces cadmium uptake and mitigates cadmium toxicity in rice. <i>Journal of Hazardous Materials</i> , 2012, 235-236, 343-351.	12.4	259
40	Genotypic dependent effect of exogenous glutathione on Cd-induced changes in proteins, ultrastructure and antioxidant defense enzymes in rice seedlings. <i>Journal of Hazardous Materials</i> , 2011, 192, 1056-1066.	12.4	72
41	Modulation of Exogenous Glutathione in Phytochelatins and Photosynthetic Performance Against Cd Stress in the Two Rice Genotypes Differing in Cd Tolerance. <i>Biological Trace Element Research</i> , 2011, 143, 1159-1173.	3.5	76