

# Kale Oluwafemi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8600818/publications.pdf>

Version: 2024-02-01

15  
papers

123  
citations

1306789

7  
h-index

1281420

11  
g-index

15  
all docs

15  
docs citations

15  
times ranked

183  
citing authors

#	ARTICLE	IF	CITATIONS
1	Ethyl acetate fraction of <i>Spathodea campanulata</i> ( <i>Bignoniaceae</i> ) attenuates lead acetate induced testicular toxicity in male Wistar rats. <i>Journal of Complementary and Integrative Medicine</i> , 2022, 19, 19-26.	0.4	4
2	M3, a 1,4-Dihydropyridine Derivative and Mixed L-/T-Type Calcium Channel Blocker, Attenuates Isoproterenol-Induced Toxicity in Male Wistar Rats. <i>Cardiovascular Toxicology</i> , 2020, 20, 627-640.	1.1	3
3	Protective Effects of <i>Acridocarpus smeathmannii</i> (DC.) Guill. & Perr. Root Extract against Phenylhydrazine-Induced Haematotoxicity, Biochemical Changes, and Oxidative Stress in Rats. <i>Biochemistry Insights</i> , 2019, 12, 117862641988324.	3.3	3
4	<i>Acridocarpus Smeathmannii</i> (DC.) Guill. & Perr. Root enhanced reproductive behavior and sexual function in male wistar rats: Biochemical and pharmacological mechanisms. <i>Journal of Ethnopharmacology</i> , 2019, 230, 95-108.	2.0	7
5	Chemoprotective effects of butanol fraction of <i>Buchholzia coriacea</i> (Capparidaceae) against type 2 diabetes and oxidative stress in male Wistar rats. <i>Bioscience Reports</i> , 2019, 39, .	1.1	8
6	Celecoxib, a cyclooxygenase-2 inhibitor, offers chemoprevention against reproductive and neurobehavioural abnormalities induced by atrazine in male Wistar rats. <i>Environmental Toxicology and Pharmacology</i> , 2018, 58, 84-97.	2.0	10
7	Five traditional Nigerian Polyherbal remedies protect against high fructose fed, Streptozotocin-induced type 2 diabetes in male Wistar rats. <i>BMC Complementary and Alternative Medicine</i> , 2018, 18, 160.	3.7	8
8	Amlodipine, an L-type calcium channel blocker, protects against chlorpromazine-induced neurobehavioural deficits in mice. <i>Fundamental and Clinical Pharmacology</i> , 2017, 31, 329-339.	1.0	5
9	Safety evaluation of Bon-sant® cleanser polyherbal in male Wistar rats. <i>BMC Complementary and Alternative Medicine</i> , 2016, 16, 188.	3.7	14
10	The combined fixed-dose antituberculous drugs alter some reproductive functions with oxidative stress involvement in wistar rats. <i>Toxicology Reports</i> , 2016, 3, 620-627.	1.6	6
11	Protective Effect of Curcumin against the Liver Toxicity Caused by Propanil in Rats. <i>International Scholarly Research Notices</i> , 2014, 2014, 1-8.	0.9	13
12	Modulation of paracetamol-induced hepatotoxicity by phosphodiesterase isozyme inhibition in rats: a preliminary study. <i>Journal of Basic and Clinical Physiology and Pharmacology</i> , 2013, 24, 73-9.	0.7	7
13	Celecoxib, a selective cyclooxygenase-2 inhibitor, lowers plasma cholesterol and attenuates hepatic lipid peroxidation during carbon-tetrachloride-associated hepatotoxicity in rats. <i>Drug and Chemical Toxicology</i> , 2013, 36, 1-8.	1.2	22
14	Subacute Therapeutic Dosing of Artemether-Lumefantrine and Artesunate-Amodiaquine Combination Preserves Plasma Cholesterol, Renal Antioxidant Status, and Organ Weights in Rats. <i>Malaria Research and Treatment</i> , 2012, 2012, 1-5.	2.0	9
15	Pharmacologic inhibition of the renin-angiotensin system did not attenuate hepatic toxicity induced by carbon tetrachloride in rats. <i>Human and Experimental Toxicology</i> , 2011, 30, 1840-1848.	1.1	4