

Janine Stein

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8609183/publications.pdf>

Version: 2024-02-01

65
papers

1,745
citations

279701

23
h-index

345118

36
g-index

84
all docs

84
docs citations

84
times ranked

2651
citing authors

#	ARTICLE	IF	CITATIONS
1	What Comes after the Trial? An Observational Study of the Real-World Uptake of an E-Mental Health Intervention by General Practitioners to Reduce Depressive Symptoms in Their Patients. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 6203.	1.2	3
2	Associations and correlates of general versus specific successful ageing components. <i>European Journal of Ageing</i> , 2021, 18, 549-563.	1.2	9
3	Valuing end-of-life care: translation and content validation of the ICECAP-SCM measure. <i>BMC Palliative Care</i> , 2021, 20, 29.	0.8	4
4	Unmet needs of the oldest old primary care patients with common somatic and psychiatric disordersâ€”A psychometric evaluation. <i>International Journal of Methods in Psychiatric Research</i> , 2021, 30, e1872.	1.1	2
5	Using a brief web-based 5A intervention to improve weight management in primary care: results of a cluster-randomized controlled trial. <i>BMC Family Practice</i> , 2021, 22, 61.	2.9	9
6	Correlates of institutionalization among the oldest oldâ€”Evidence from the multicenter AgeCoDeâ€”AgeQualiDe study. <i>International Journal of Geriatric Psychiatry</i> , 2021, 36, 1095-1102.	1.3	13
7	Needs and Healthcare Costs in Old Age. , 2021, , 91-96.		0
8	Needs of Older Primary Care Patients. , 2021, , 20-33.		0
9	Loss and bereavement in late life (60+): Study protocol for a randomized controlled trial regarding an internet-based self-help intervention. <i>Internet Interventions</i> , 2021, 26, 100451.	1.4	3
10	Driving status and health-related quality of life among the oldest old: a population-based examination using data from the AgeCoDeâ€”AgeQualiDe prospective cohort study. <i>Aging Clinical and Experimental Research</i> , 2021, 33, 3109-3115.	1.4	8
11	Effectiveness and Feasibility of Internet-Based Interventions for Grief After Bereavement: Systematic Review and Meta-analysis. <i>JMIR Mental Health</i> , 2021, 8, e29661.	1.7	17
12	Late-Life Depressive Symptoms Are Associated With Functional Impairment Cross-sectionally and Over Time: Results of the AgeMooDe Study. <i>Journals of Gerontology - Series B Psychological Sciences and Social Sciences</i> , 2020, 75, 811-820.	2.4	6
13	Needs and health care costs in old age: a longitudinal perspective: results from the AgeMooDe study. <i>Aging and Mental Health</i> , 2020, 24, 1763-1768.	1.5	8
14	Correlates of hospitalization among the oldest old: results of the AgeCoDeâ€”AgeQualiDe prospective cohort study. <i>Aging Clinical and Experimental Research</i> , 2020, 32, 1295-1301.	1.4	14
15	Effectiveness and feasibility of internet-based and mobile-based interventions for individuals experiencing bereavement: a systematic review protocol. <i>BMJ Open</i> , 2020, 10, e036034.	0.8	2
16	Determinants of Willingness to Pay for Health Insurance in Germanyâ€”Results of the Population-Based Health Study of the Leipzig Research Centre for Civilization Diseases (LIFE-Adult-Study). <i>Frontiers in Public Health</i> , 2020, 8, 456.	1.3	7
17	Internet-Based Cognitive Behavior Therapy Only for the Young? A Secondary Analysis of a Randomized Controlled Trial of Depression Treatment. <i>Frontiers in Psychiatry</i> , 2020, 11, 735.	1.3	8
18	Unmet care needs in the oldest old with social loss experiences: results of a representative survey. <i>BMC Geriatrics</i> , 2020, 20, 416.	1.1	8

#	ARTICLE	IF	CITATIONS
19	Internet Use in Old Age: Results of a German Population-Representative Survey. <i>Journal of Medical Internet Research</i> , 2020, 22, e15543.	2.1	50
20	Are older men more vulnerable to depression than women after losing their spouse? Evidence from three German old-age cohorts (AgeDifferent.de platform). <i>Journal of Affective Disorders</i> , 2019, 256, 650-657.	2.0	23
21	Does transpersonal trust moderate the association between chronic conditions and general practitioner visits in the oldest old? Results of the AgeCoDe and AgeQualiDe study. <i>Geriatrics and Gerontology International</i> , 2019, 19, 705-710.	0.7	5
22	Psychometric properties of the Patient Assessment of Chronic Illness Care measure (PACIC-5A) among patients with obesity. <i>BMC Health Services Research</i> , 2019, 19, 61.	0.9	4
23	Prevalence of Anxiety Symptoms and Their Association With Loss Experience in a Large Cohort Sample of the Oldest-Old. Results of the AgeCoDe AgeQualiDe Study. <i>Frontiers in Psychiatry</i> , 2019, 10, 285.	1.3	22
24	Determinants of health-care costs in the oldest-old in Germany. <i>Journal of the Economics of Ageing</i> , 2019, 14, 100200.	0.6	7
25	Prevalence and determinants of driving habits in the oldest old: Results of the multicenter prospective AgeCoDe-AgeQualiDe study. <i>Archives of Gerontology and Geriatrics</i> , 2019, 82, 245-250.	1.4	11
26	Association of anxiety symptoms with health care use and costs in people aged 85 and over. <i>International Journal of Geriatric Psychiatry</i> , 2019, 34, 765-776.	1.3	4
27	Unmet health care needs in old age and their association with depression – results of a population-representative survey. <i>Journal of Affective Disorders</i> , 2019, 245, 998-1006.	2.0	28
28	Unmet needs in the depressed primary care elderly and their relation to severity of depression: results from the AgeMooDe study. <i>Aging and Mental Health</i> , 2018, 22, 1038-1045.	1.5	6
29	Categorical and dimensional perspectives on depression in elderly primary care patients – Results of the AgeMooDe study. <i>International Journal of Methods in Psychiatric Research</i> , 2018, 27, .	1.1	5
30	Loss experiences in old age and their impact on the social network and depression – results of the Leipzig Longitudinal Study of the Aged (LEILA 75+). <i>Journal of Affective Disorders</i> , 2018, 241, 94-102.	2.0	24
31	Preferences for Depression Treatment Including Internet-Based Interventions: Results From a Large Sample of Primary Care Patients. <i>Frontiers in Psychiatry</i> , 2018, 9, 181.	1.3	53
32	Impact of falls on depressive symptoms among the oldest old: Results from the AgeQualiDe study. <i>International Journal of Geriatric Psychiatry</i> , 2018, 33, 1383-1388.	1.3	13
33	Five A – counseling in weight management of obese patients in primary care: a cluster-randomized controlled trial (INTERACT). <i>BMC Family Practice</i> , 2018, 19, 97.	2.9	13
34	A qualitative study on older primary care patients – perspectives on depression and its treatments - potential barriers to and opportunities for managing depression. <i>BMC Family Practice</i> , 2018, 19, 2.	2.9	26
35	Computerized cognitive behavior therapy for patients with mild to moderately severe depression in primary care: A pragmatic cluster randomized controlled trial (@ktiv). <i>Journal of Affective Disorders</i> , 2018, 238, 317-326.	2.0	53
36	Disentangling the complex relation of disability and depressive symptoms in old age – findings of a multicenter prospective cohort study in Germany. <i>International Psychogeriatrics</i> , 2017, 29, 885-895.	0.6	30

#	ARTICLE	IF	CITATIONS
37	Treatment preferences for depression in the elderly. <i>International Psychogeriatrics</i> , 2017, 29, 389-398.	0.6	34
38	How does social support affect functional impairment in late life? Findings of a multicenter prospective cohort study in Germany. <i>Age and Ageing</i> , 2017, 46, 813-820.	0.7	23
39	Unmet Care Needs in the Oldest Old Primary Care Patients with Cognitive Disorders: Results of the AgeCoDe and AgeQualiDe Study. <i>Dementia and Geriatric Cognitive Disorders</i> , 2017, 44, 71-83.	0.7	19
40	The Impact of Depressive Symptoms on Healthcare Costs in Late Life: Longitudinal Findings From the AgeMooDe Study. <i>American Journal of Geriatric Psychiatry</i> , 2017, 25, 131-141.	0.6	22
41	Which types of mental work demands may be associated with reduced risk of dementia?. <i>Alzheimer's and Dementia</i> , 2017, 13, 431-440.	0.4	33
42	Choosing the right rehabilitation setting after herniated disc surgery: Motives, motivations and expectations from the patients' perspective. <i>PLoS ONE</i> , 2017, 12, e0183698.	1.1	1
43	Frequent attenders in late life in primary care: a systematic review of European studies. <i>BMC Family Practice</i> , 2017, 18, 104.	2.9	37
44	The impact of social engagement on health-related quality of life and depressive symptoms in old age - evidence from a multicenter prospective cohort study in Germany. <i>Health and Quality of Life Outcomes</i> , 2017, 15, 140.	1.0	48
45	Risk Factors for Postoperative Pain Intensity in Patients Undergoing Lumbar Disc Surgery: A Systematic Review. <i>PLoS ONE</i> , 2017, 12, e0170303.	1.1	35
46	User Acceptance of Computerized Cognitive Behavioral Therapy for Depression: Systematic Review. <i>Journal of Medical Internet Research</i> , 2017, 19, e309.	2.1	81
47	Unmet care needs of the oldest old with late-life depression: A comparison of patient, caring relative and general practitioner perceptions – Results of the AgeMooDe study. <i>Journal of Affective Disorders</i> , 2016, 205, 182-189.	2.0	10
48	Late-Life Depressive Symptoms and Lifetime History of Major Depression: Cognitive Deficits are Largely Due to Incipient Dementia rather than Depression. <i>Journal of Alzheimer's Disease</i> , 2016, 54, 185-199.	1.2	23
49	Differential Risk of Incident Alzheimer's Disease Dementia in Stable Versus Unstable Patterns of Subjective Cognitive Decline. <i>Journal of Alzheimer's Disease</i> , 2016, 54, 1135-1146.	1.2	70
50	Excess health care costs of late-life depression – Results of the AgeMooDe study. <i>Journal of Affective Disorders</i> , 2016, 199, 139-147.	2.0	40
51	The impact of depressive symptoms on utilization of home care by the elderly: Longitudinal results from the AgeMooDe study. <i>Journal of Affective Disorders</i> , 2016, 204, 247-254.	2.0	6
52	Longitudinal predictors of informal and formal caregiving time in community-dwelling dementia patients. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016, 51, 607-616.	1.6	46
53	Gender differences in the effect of social support on health-related quality of life: results of a population-based prospective cohort study in old age in Germany. <i>Quality of Life Research</i> , 2016, 25, 1159-1168.	1.5	61
54	The assessment of met and unmet care needs in the oldest old with and without depression using the Camberwell Assessment of Need for the Elderly (CANE): Results of the AgeMooDe study. <i>Journal of Affective Disorders</i> , 2016, 193, 309-317.	2.0	49

#	ARTICLE	IF	CITATIONS
55	The Course of Pain Intensity in Patients Undergoing Herniated Disc Surgery: A 5-Year Longitudinal Observational Study. PLoS ONE, 2016, 11, e0156647.	1.1	8
56	The German version of the Camberwell Assessment of Need for the Elderly (CANE): evaluation of content validity and adaptation to the German-speaking context. International Psychogeriatrics, 2015, 27, 1919-1926.	0.6	19
57	Is the Short Form of the Mini-Mental State Examination (MMSE) a better screening instrument for dementia in older primary care patients than the original MMSE? Results of the German study on ageing, cognition, and dementia in primary care patients (AgeCoDe).. Psychological Assessment, 2015, 27, 895-904.	1.2	9
58	Complex coevolution of depression and health-related quality of life in old age. Quality of Life Research, 2015, 24, 2713-2722.	1.5	20
59	Longitudinal Predictors of Institutionalization in Old Age. PLoS ONE, 2015, 10, e0144203.	1.1	128
60	Assessing met and unmet needs in the oldest-old and psychometric properties of the German version of the Camberwell Assessment of Need for the Elderly (CANE) – a pilot study. International Psychogeriatrics, 2014, 26, 285-295.	0.6	25
61	The CERAD Neuropsychological Assessment Battery Total Score Detects and Predicts Alzheimer Disease Dementia with High Diagnostic Accuracy. American Journal of Geriatric Psychiatry, 2014, 22, 1017-1028.	0.6	56
62	Measuring Negative Attitudes towards Overweight and Obesity in the German Population – Psychometric Properties and Reference Values for the German Short Version of the Fat Phobia Scale (FPS). PLoS ONE, 2014, 9, e114641.	1.1	21
63	The Assessment of Changes in Cognitive Functioning in the Elderly: Age- and Education-Specific Reliable Change Indices for the SIDAM. Dementia and Geriatric Cognitive Disorders, 2012, 33, 73-83.	0.7	11
64	The Assessment of Changes in Cognitive Functioning: Age-, Education-, and Gender-Specific Reliable Change Indices for Older Adults Tested on the CERAD-NP Battery: Results of the German Study on Ageing, Cognition, and Dementia in Primary Care Patients (AgeCoDe). American Journal of Geriatric Psychiatry, 2012, 20, 84-97.	0.6	29
65	The Assessment of Changes in Cognitive Functioning: Reliable Change Indices for Neuropsychological Instruments in the Elderly – A Systematic Review. Dementia and Geriatric Cognitive Disorders, 2010, 29, 275-286.	0.7	53