

# Jun Fan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8611009/publications.pdf>

Version: 2024-02-01

115  
papers

2,859  
citations

201674

27  
h-index

206112

48  
g-index

118  
all docs

118  
docs citations

118  
times ranked

3089  
citing authors

#	ARTICLE	IF	CITATIONS
1	Interactions of diclazuril enantiomers with serum albumins: Multi-spectroscopic and molecular docking approaches. <i>Journal of Molecular Recognition</i> , 2022, 35, e2948.	2.1	2
2	Risk Assessment of the Chiral Fungicide Triconazole: Enantioselective Effects, Toxicity, and Fate. <i>Journal of Agricultural and Food Chemistry</i> , 2022, 70, 2712-2721.	5.2	12
3	Enantioselective acute toxicity, oxidative stress effects, neurotoxicity, and thyroid disruption of uniconazole in zebrafish ( <i>Danio rerio</i> ). <i>Environmental Science and Pollution Research</i> , 2022, 29, 40157-40168.	5.3	4
4	Enantioselective bioaccumulation, oxidative stress, and thyroid disruption assessment of cis-metconazole enantiomers in zebrafish ( <i>Danio rerio</i> ). <i>Aquatic Toxicology</i> , 2022, 248, 106205.	4.0	7
5	Fabrication of cellulose derivative coated spherical covalent organic frameworks as chiral stationary phases for high-performance liquid chromatographic enantioseparation. <i>Journal of Chromatography A</i> , 2022, 1675, 463155.	3.7	11
6	Triconazole enantiomers induced enantioselective metabolic phenotypes in <i>Fusarium graminearum</i> and HepG2 cells. <i>Environmental Science and Pollution Research</i> , 2022, 29, 75978-75988.	5.3	3
7	Enantioselective neurotoxicity and oxidative stress effects of paclobutrazol in zebrafish ( <i>Danio rerio</i> ). <i>Environmental Science and Pollution Research</i> , 2022, 29, 75978-75988.	3.6	3
8	Transformation of a Hydrazone-Linked Covalent Organic Framework into a Highly Stable Hydrazide-Linked One. <i>ACS Applied Polymer Materials</i> , 2022, 4, 4624-4631.	4.4	13
9	Stereoselective in vitro metabolism of cyproconazole in rat liver microsomes and identification of major metabolites. <i>Chemosphere</i> , 2021, 264, 128495.	8.2	13
10	Covalent Cross-Linking of Metal-Organic Cages: Formation of an Amorphous Cationic Porous Extended Framework for the Uptake of Oxo-Anions from Water. <i>ChemPlusChem</i> , 2021, 86, 709-715.	2.8	8
11	A hydrolytically stable cage-based metal-organic framework containing two types of building blocks for the adsorption of iodine and dyes. <i>Inorganic Chemistry Frontiers</i> , 2021, 8, 1083-1092.	6.0	55
12	Development of a sensitive and stable GC-MS/MS method for simultaneous determination of four N-nitrosamine genotoxic impurities in sartan substances. <i>Journal of Analytical Science and Technology</i> , 2021, 12, .	2.1	12
13	Metal-Free Polymer-Based Affinity Medium for Selective Purification of His6-Tagged Proteins. <i>Biomacromolecules</i> , 2021, 22, 1695-1705.	5.4	8
14	Covalent Cross-Linking of Metal-Organic Cages: Formation of an Amorphous Cationic Porous Extended Framework for the Uptake of Oxo-Anions from Water. <i>ChemPlusChem</i> , 2021, 86, 699-699.	2.8	1
15	Systematic investigation of stereochemistry, stereoselective bioactivity, and antifungal mechanism of chiral triazole fungicide metconazole. <i>Science of the Total Environment</i> , 2021, 784, 147194.	8.0	12
16	Stereochemistry of chiral pesticide uniconazole and enantioselective metabolism in rat liver microsomes. <i>Pesticide Biochemistry and Physiology</i> , 2021, 179, 104964.	3.6	6
17	A hydrolytically stable hydrogen-bonded inorganic-organic network as a luminescence turn-on sensor for the detection of Bi <sup>3+</sup> and Fe <sup>3+</sup> cations in water. <i>Polyhedron</i> , 2021, 205, 115284.	2.2	9
18	A recyclable bipyridine-containing covalent organic framework-based QCM sensor for detection of Hg(II) ion in aqueous solution. <i>Journal of Solid State Chemistry</i> , 2021, 302, 122421.	2.9	19

#	ARTICLE	IF	CITATIONS
19	Facile synthesis of nitrogen-doped carbon dots from pork liver and its sensing of 6-thioguanine based on the inner filter effect. <i>New Journal of Chemistry</i> , 2021, 45, 5114-5120.	2.8	10
20	Stable hydrazone-linked chiral covalent organic frameworks: Synthesis, modification, and chiral signal inversion from monomers. <i>Chinese Chemical Letters</i> , 2021, 32, 107-112.	9.0	15
21	Enantioselective analyses of chloroquine and hydroxychloroquine in rat liver microsomes through chiral liquid chromatography-tandem mass spectrometry. <i>Chirality</i> , 2021, 34, 126.	2.6	2
22	Facile and Site-Selective Synthesis of an Amine-Functionalized Covalent Organic Framework. <i>ACS Macro Letters</i> , 2021, 10, 1590-1596.	4.8	32
23	A new QCM signal enhancement strategy based on streptavidin@metal-organic framework complex for miRNA detection. <i>Analytica Chimica Acta</i> , 2020, 1095, 212-218.	5.4	13
24	HPLC semi-preparative separation of diclazuril enantiomers and racemization in solution. <i>Journal of Separation Science</i> , 2020, 43, 1240-1247.	2.5	9
25	A new amplification strategy for a quartz crystal microbalance miRNA sensor based on selective interactions between a metal-organic framework and miRNA. <i>New Journal of Chemistry</i> , 2020, 44, 1684-1688.	2.8	4
26	A Mn-MOF with inherent missing metal-ion defects based on an imidazole-tetrazole tripodal ligand and its application in supercapacitors. <i>Dalton Transactions</i> , 2020, 49, 12150-12155.	3.3	11
27	Four cypermethrin isomers induced stereoselective metabolism in H295R cells. <i>Chirality</i> , 2020, 32, 1107-1118.	2.6	5
28	Pharmacokinetics, Activity, and Residue Elimination of R- and S-Diclazuril in Broiler Chickens. <i>Journal of Agricultural and Food Chemistry</i> , 2020, 68, 8987-8995.	5.2	5
29	Reversible Interlayer Sliding and Conductivity Changes in Adaptive Tetrathiafulvalene-Based Covalent Organic Frameworks. <i>ACS Applied Materials &amp; Interfaces</i> , 2020, 12, 19054-19061.	8.0	40
30	Protein A-mesoporous silica composites for chromatographic purification of immunoglobulin G. <i>New Journal of Chemistry</i> , 2020, 44, 7884-7890.	2.8	4
31	Cu-MOF derived Cu-C nanocomposites towards high performance electrochemical supercapacitors. <i>RSC Advances</i> , 2020, 10, 4621-4629.	3.6	17
32	Fast enantioselective determination of triadimefon in different matrices by supercritical fluid chromatography. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2019, 1126-1127, 121740.	2.3	10
33	Cationic Amorphous Metal-Organic Cage-Based Materials for the Removal of Oxo-Anions from Water. <i>ACS Applied Nano Materials</i> , 2019, 2, 5824-5832.	5.0	28
34	Semi-preparative separation of dihydromyricetin enantiomers by supercritical fluid chromatography and determination of anti-inflammatory activities. <i>Journal of Chromatography A</i> , 2019, 1606, 460386.	3.7	17
35	Identification, Quantification, and Stereoselective Degradation of Triazole Fungicide Cyproconazole in Two Matrixes through Chiral Liquid Chromatography-Tandem Mass Spectrometry. <i>Journal of Agricultural and Food Chemistry</i> , 2019, 67, 10782-10790.	5.2	20
36	An Anionic Nanotubular Metal-Organic Framework for High-Capacity Dye Adsorption and Dye Degradation in Darkness. <i>Inorganic Chemistry</i> , 2019, 58, 13979-13987.	4.0	75

#	ARTICLE	IF	CITATIONS
37	Hydrolytically Stable Nanotubular Cationic Metal-Organic Framework for Rapid and Efficient Removal of Toxic Oxo-Anions and Dyes from Water. <i>Inorganic Chemistry</i> , 2019, 58, 2899-2909.	4.0	106
38	A Benzimidazole-Containing Covalent Organic Framework-Based QCM Sensor for Exceptional Detection of CEES. <i>Crystal Growth and Design</i> , 2019, 19, 3543-3550.	3.0	26
39	Determination of residual enantiomers of diclazuril in chicken edible tissues by high performance liquid chromatography. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2019, 1118-1119, 203-209.	2.3	12
40	Stable Hydrazone-Linked Covalent Organic Frameworks Containing O,N-Chelating Sites for Fe(III) Detection in Water. <i>ACS Applied Materials &amp; Interfaces</i> , 2019, 11, 12830-12837.	8.0	152
41	Fabrication of a hydrazone-linked covalent organic framework-bound capillary column for gas chromatography separation. <i>Separation Science Plus</i> , 2019, 2, 120-128.	0.6	14
42	An unprecedented 2D covalent organic framework with an htb net topology. <i>Chemical Communications</i> , 2019, 55, 13454-13457.	4.1	26
43	Enantioselectivity in endocrine disrupting effects of four cypermethrin enantiomers based on in vitro models. <i>Chemosphere</i> , 2019, 220, 766-773.	8.2	14
44	Assembly of a miRNA-modified QCM sensor for miRNA recognition through response patterns. <i>Journal of Molecular Recognition</i> , 2019, 32, e2772.	2.1	3
45	Efficient preparative separation of $\beta$ -cypermethrin stereoisomers by supercritical fluid chromatography with a two-step combined strategy. <i>Journal of Separation Science</i> , 2018, 41, 1442-1449.	2.5	11
46	High-fast enantioselective determination of prothioconazole in different matrices by supercritical fluid chromatography and vibrational circular dichroism spectroscopic study. <i>Talanta</i> , 2018, 187, 40-46.	5.5	31
47	Enantioselective determination of metconazole in multi matrices by high-performance liquid chromatography. <i>Talanta</i> , 2018, 178, 980-986.	5.5	32
48	A new single-urea-bound 3,5-dimethylphenylcarbamoylated $\beta$ -cyclodextrin chiral stationary phase and its enhanced separation performance in normal-phase liquid chromatography. <i>Electrophoresis</i> , 2018, 39, 348-355.	2.4	21
49	HPLC Enantioseparation of Menthol with Non-ultraviolet Detectors and Effect of Chromatographic Conditions. <i>Chromatographia</i> , 2018, 81, 871-879.	1.3	1
50	Lenalidomide, a blockbuster drug for the treatment of multiple myeloma: Semipreparative separation through supercritical fluid chromatography and vibrational circular dichroism spectroscopy. <i>Journal of Separation Science</i> , 2018, 41, 3840-3847.	2.5	8
51	A series of alkaline earth metal coordination polymers constructed from two newly designed imidazole-based dicarboxylate ligands containing pyridinylmethyl groups. <i>CrystEngComm</i> , 2017, 19, 3003-3016.	2.6	16
52	Construction of Four Coordination Polymers based on 2-[4-(Pyridine-4-yl)phenyl]-1H-imidazole-4,5-dicarboxylic Acid. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , 2017, 643, 593-600.	1.2	8
53	Anion and pH-regulated assembly of three Cd(II) coordination polymers based on 3,5-di(1H-benzo[d]imidazol-1-yl)benzoate. <i>Journal of Coordination Chemistry</i> , 2017, 70, 135-144.	2.2	4
54	Construction of a hydrazone-linked chiral covalent organic framework-silica composite as the stationary phase for high performance liquid chromatography. <i>Journal of Chromatography A</i> , 2017, 1519, 100-109.	3.7	110

#	ARTICLE	IF	CITATIONS
55	Reliable HPLC separation, vibrational circular dichroism spectra, and absolute configurations of isoborneol enantiomers. <i>Chirality</i> , 2017, 29, 550-557.	2.6	10
56	Lanthanide contraction effect on the crystal structures of 2D lanthanide coordination polymers based on 2-(trifluoromethyl)-1H-imidazole-4,5-dicarboxylic acid. <i>Structural Chemistry</i> , 2017, 28, 577-586.	2.0	9
57	Stereoselective quantification of triticonazole in vegetables by supercritical fluid chromatography. <i>Talanta</i> , 2017, 164, 362-367.	5.5	50
58	Anion- and temperature-dependent assembly, crystal structures and luminescence properties of six new Cd( $\text{II}$ ) coordination polymers based on 2,3,5,6-tetrakis(2-pyridyl)pyrazine. <i>CrystEngComm</i> , 2016, 18, 5164-5176.	2.6	24
59	Construction of six new luminescent Ln( $\text{III}$ )-Zn( $\text{II}$ ) heterometallic coordination polymers based on heterometallic secondary building units. <i>CrystEngComm</i> , 2016, 18, 8672-8682.	2.6	16
60	Triticonazole enantiomers: Separation by supercritical fluid chromatography and the effect of the chromatographic conditions. <i>Journal of Separation Science</i> , 2016, 39, 4251-4257.	2.5	23
61	Rationally Designed 2D Covalent Organic Framework with a Brick-Wall Topology. <i>ACS Macro Letters</i> , 2016, 5, 1348-1352.	4.8	59
62	Analysis of metalaxyl racemate using high performance liquid chromatography coupled with four kinds of detectors. <i>Journal of Chromatography A</i> , 2016, 1467, 246-254.	3.7	11
63	Two new three-dimensional metal-organic frameworks with 4-connected diamondoid and unusual (6,16)-connected net topologies based on planar tetranuclear squares as secondary building units. <i>CrystEngComm</i> , 2016, 18, 1174-1183.	2.6	15
64	Comparative separation of chiral compounds by supercritical fluid chromatography and high performance liquid chromatography. <i>Chinese Journal of Chromatography (Se Pu)</i> , 2016, 34, 321.	0.8	3
65	Syntheses, structures, and properties of nine d10 or p-block coordination polymers based on a ligand containing both terpyridyl and sulfo groups. <i>CrystEngComm</i> , 2015, 17, 5538-5550.	2.6	12
66	Synthesis of a New Cyclosporine-based Stationary Phase and Separation Behaviors toward Aromatic Positional Isomers by High-Performance Liquid Chromatography. <i>Journal of Chromatographic Science</i> , 2015, 53, 548-553.	1.4	3
67	Assembly of Cd( $\text{II}$ ) coordination polymers: structural variation, supramolecular isomers, and temperature/anion-induced solvent-mediated structural transformations. <i>CrystEngComm</i> , 2015, 17, 947-959.	2.6	36
68	Two Coordination Polymers Constructed from 5-(4-pyridyl)-1H-tetrazole Ligands with Different Organic Carboxylates: Structure and Luminescence Properties. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , 2014, 640, 2057-2061.	1.2	6
69	Synthesis and enantioseparation behaviors of novel immobilized 3,5-dimethylphenylcarbamoylated polysaccharide chiral stationary phases. <i>Journal of Separation Science</i> , 2014, 37, 488-494.	2.5	16
70	Construction of terpyridine-Ln( $\text{III}$ ) coordination polymers: structural diversity, visible and NIR luminescence properties and response to nerve-agent mimics. <i>CrystEngComm</i> , 2014, 16, 2898.	2.6	39
71	Tunable electrical conductivity in oriented thin films of tetrathiafulvalene-based covalent organic framework. <i>Chemical Science</i> , 2014, 5, 4693-4700.	7.4	295
72	Construction of several new s/p-block complexes containing binuclear metal-terpyridine building blocks: dependence of structural diversity on the number of coordinated water molecules. <i>CrystEngComm</i> , 2014, 16, 4029.	2.6	16

#	ARTICLE	IF	CITATIONS
73	Comparative HPLC enantioseparation on substituted phenylcarbamoylated cyclodextrin chiral stationary phases and mobile phase effects. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 98, 221-227.	2.8	22
74	Construction of Ag(I)-Ln(III) Heterometallic Coordination Polymers Based on Binuclear Ag <sub>2</sub> (DSPT) <sub>2</sub> (H <sub>2</sub> DSPT = 4-(2,4-Disulfophenyl)-2,2,6,6-tetra-2-terpyridine) Rings and Ln(III) Dimeric Molecular Building Blocks. <i>Crystal Growth and Design</i> , 2013, 13, 4428-4434.		43
75	Tubular metal-organic framework-based capillary gas chromatography column for separation of alkanes and aromatic positional isomers. <i>Journal of Chromatography A</i> , 2013, 1285, 132-138.	3.7	64
76	Synthesis of a novel cyclodextrin-derived chiral stationary phase with multiple urea linkages and enantioseparation toward chiral osmabenzene complex. <i>Journal of Chromatography A</i> , 2013, 1283, 68-74.	3.7	21
77	The construction of Cu(i)/Cu(ii) coordination polymers based on pyrazine-carboxylate: Structural diversity tuned by in situ hydrolysis reaction. <i>CrystEngComm</i> , 2013, 15, 5359.	2.6	26
78	Reversal of elution order of N-(2,4-dinitrophenyl)proline and N-(2,4-dinitrophenyl)serine in HPLC by BSA chiral stationary phase. <i>Journal of Separation Science</i> , 2013, 36, 1343-1348.	2.5	5
79	Construction of luminescent three-dimensional Ln(iii)-Zn(ii) heterometallic coordination polymers based on 2-pyridyl imidazole dicarboxylate. <i>CrystEngComm</i> , 2012, 14, 8236.	2.6	29
80	Two Types of New Three-Dimensional f Heterometallic Coordination Polymers Based on 2-(Pyridin-3-yl)-1H-imidazole-4,5-Dicarboxylate and Oxalate Ligands: Syntheses, Structures, Luminescence, and Magnetic Properties. <i>Crystal Growth and Design</i> , 2012, 12, 4441-4449.	3.0	63
81	Construction of Ba(II) Coordination Polymers Based on Imidazole-Based Dicarboxylate Ligands: Structural Diversity Tuned by Alcohol Solvents. <i>Crystal Growth and Design</i> , 2012, 12, 3575-3582.	3.0	59
82	A Series of New Three-Dimensional f Heterometallic Coordination Polymers with Rare 10-Connected Net Topology Based on Planar Hexanuclear Heterometallic Second Building Units. <i>Crystal Growth and Design</i> , 2012, 12, 5737-5745.	3.0	67
83	Solvent-regulated assembly of two new Cd(II) coordination polymers based on 3-(1H-benzimidazol-2-yl) propanoic acid. <i>Inorganic Chemistry Communication</i> , 2012, 21, 100-103.	3.9	9
84	Assembly of two new Mn(II) coordination polymers based on 5-aminoisophthalate: Structural diversity and properties. <i>Inorganic Chemistry Communication</i> , 2012, 22, 93-97.	3.9	2
85	A New Biosensor for Chiral Recognition Using Goat and Rabbit Serum Albumin Self-Assembled Quartz Crystal Microbalance. <i>Chirality</i> , 2012, 24, 804-809.	2.6	5
86	An unprecedented (3,4,14)-connected 3D metal-organic framework based on planar octanuclear lead(ii) clusters as a secondary building unit. <i>CrystEngComm</i> , 2012, 14, 1193-1196.	2.6	36
87	Anion-dependent assembly and solvent-mediated structural transformations of three Cd(ii) coordination polymers based on 1H-imidazole-4-carboxylic acid. <i>CrystEngComm</i> , 2012, 14, 2308.	2.6	36
88	New Poly(N,N-Dimethylaminoethyl Methacrylate)/Polyvinyl Alcohol Copolymer Coated QCM Sensor for Interaction with CWA Simulants. <i>ACS Applied Materials &amp; Interfaces</i> , 2012, 4, 944-949.	8.0	19
89	Assembly of Chiral/Achiral Coordination Polymers Based on 2-(Pyridine-3-yl)-1H-4,5-imidazolecarboxylic Acid: Chirality Transfer between Chiral Two-Dimensional Networks Containing Helical Chains. <i>Crystal Growth and Design</i> , 2012, 12, 2355-2361.	3.0	57
90	Spontaneous resolution of a coordination polymer containing stereogenic five-coordinate Zn(ii) centers and achiral ligands with axially chiral conformation. <i>CrystEngComm</i> , 2012, 14, 6241.	2.6	13

#	ARTICLE	IF	CITATIONS
91	Effect of Ionic-Strength and pH Value in Mobile Phase on Enantio- Separation of BSA High Performance Liquid Chromatography Column. Chinese Journal of Analytical Chemistry, 2012, 40, 89.	1.7	2
92	An unprecedented supramolecular network with channels filled by 1D coordination polymer chains: Cocrystallization of Ag(i)-4,4'-bipyridine and Ag(i)-benzimidazole complexes. CrystEngComm, 2011, 13, 6345.	2.6	17
93	The construction of coordination networks based on imidazole-based dicarboxylate ligand containing hydroxymethyl group. CrystEngComm, 2011, 13, 883-888.	2.6	68
94	Synthesis, Crystal Structures and Thermal Stabilities of Lanthanide Coordination Polymers with 5-Nitroisophthalate. Journal of Inorganic and Organometallic Polymers and Materials, 2011, 21, 723-729.	3.7	2
95	Synthesis, crystal structure, supramolecular assembly, and thermal stability of two new lanthanide coordination polymers based on 1-naphthoxyacetate. Structural Chemistry, 2011, 22, 943-949.	2.0	2
96	Luminescent lanthanide complexes with 4-acetamidobenzoate: Synthesis, supramolecular assembly via hydrogen bonds, crystal structures and photoluminescence. Journal of Solid State Chemistry, 2011, 184, 1850-1857.	2.9	8
97	Synthesis, Crystal Structures, and Photoluminescence of Lanthanide Coordination Polymers with 4-Acetamidobenzoate. Zeitschrift Fur Anorganische Und Allgemeine Chemie, 2011, 637, 773-777.	1.2	7
98	Two mononuclear octahedral complexes with benzimidazole-2-carboxylate: supramolecular networks constructed by hydrogen bonds. Acta Crystallographica Section C: Crystal Structure Communications, 2011, 67, m346-m350.	0.4	5
99	Two new 2D coordination polymers containing bowl-shaped voids assembled from the bis(chelating) bridging ligand. Inorganic Chemistry Communication, 2011, 14, 818-821.	3.9	11
100	Two new coordination frameworks based on rod-shaped secondary building blocks with five-nodal (3, 6) topology. Inorganic Chemistry Communication, 2011, 14, 1156-1160.	3.9	7
101	Synthesis, characterization, structures and magnetic property of chiral oxalate-bridged dicopper(II) complexes. Science China Chemistry, 2010, 53, 1255-1260.	8.2	4
102	A two-dimensional lanthanide coordination framework with a new amide-type tripodal ligand, 2,2',2''-nitrotris[[(2-benzylaminoformyl)phenoxy]ethyl]amine. Chinese Journal of Chemistry, 2010, 22, 508-511.	4.9	2
103	Phenylcarbamoylated $\beta$ -CD: Acidic and basic chiral selectors for HPLC. Journal of Separation Science, 2010, 33, 1558-1562.	2.5	14
104	The impact of silica gel pore and particle sizes on HPLC column efficiency and resolution for an immobilized, cyclodextrin-based, chiral stationary phase. Journal of Separation Science, 2010, 33, 2582-2589.	2.5	19
105	Synthesis, crystal structures and photoluminescent properties of lanthanide supramolecular complexes with 4-oxo-1(4H)-quinolineacetate. Journal of Solid State Chemistry, 2010, 183, 575-583.	2.9	8
106	$\{[Eu(ox)(H_2O)_2]_4[CuBr(2-pz)_2]_4 \cdot 4H_2O\}$ : Hydrogen Bonding Directed Assembly to Supramolecular Network. Zeitschrift Fur Anorganische Und Allgemeine Chemie, 2010, 636, 1111-1114.	1.2	2
107	Controlled synthesis, structures and properties of one-, two-, and three-dimensional lanthanide coordination polymers based on (8-quinolyloxy)acetate. CrystEngComm, 2010, 12, 216-225.	2.6	18
108	Synthesis and Crystal Structure of a New Binuclear Samarium Complex with Salicylate. Journal of Chemical Crystallography, 2009, 39, 585-588.	1.1	1

#	ARTICLE	IF	CITATIONS
109	Studies of Radii-Dependent Lanthanide Coordination Behavior with 4-Acetamidobenzoate and 1,10-Phenanthroline. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , 2009, 635, 2333-2339.	1.2	19
110	A novel strategy for rapid real-time chiral discrimination of enantiomers using serum albumin functionalized QCM biosensor. <i>Biosensors and Bioelectronics</i> , 2009, 25, 488-492.	10.1	44
111	Preparation and enantioseparation characteristics of a novel chiral stationary phase based on mono (6A-azido-6A-deoxy)-per(p-chlorophenylcarbamoylated) $\beta$ -cyclodextrin. <i>Journal of Chromatography A</i> , 2008, 1213, 162-168.	3.7	36
112	Crystallographic report: Bis[bis(N,N-dibenzylthiocarbamato)cadmium(II)]. <i>Applied Organometallic Chemistry</i> , 2004, 18, 139-140.	3.5	10
113	Crystallographic report: Bis[bis(N,N-dibenzylthiocarbamato)zinc(II)](4,4'-bipyridine). <i>Applied Organometallic Chemistry</i> , 2003, 17, 889-890.	3.5	9
114	Self-Assembly of Porphyrin Arrays via Coordination to Transition Metal Bisphosphine Complexes and the Unique Spectral Properties of the Product Metallacyclic Ensembles. <i>Journal of the American Chemical Society</i> , 1999, 121, 2741-2752.	13.7	203
115	A Ni metal-organic framework with helical channels for the capture of iodine via guest exchange induced amorphization. <i>New Journal of Chemistry</i> , 0, , .	2.8	7