

# Eloisa Colin-Ramirez

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8616385/publications.pdf>

Version: 2024-02-01

54  
papers

995  
citations

394421

19  
h-index

434195

31  
g-index

56  
all docs

56  
docs citations

56  
times ranked

1678  
citing authors

#	ARTICLE	IF	CITATIONS
1	Editorial: Nutritional Assessment Tools for Identification and Monitoring of Malnutrition in Patients With Chronic Disease. <i>Frontiers in Nutrition</i> , 2022, 9, 870514.	3.7	1
2	Reduction of dietary sodium to less than 100 mmol in heart failure (SODIUM-HF): an international, open-label, randomised, controlled trial. <i>Lancet, The</i> , 2022, 399, 1391-1400.	13.7	67
3	Serum Asprosin Concentrations in Children with Prader-Willi Syndrome: Correlations with Metabolic Parameters. <i>Journal of Clinical Medicine</i> , 2022, 11, 2268.	2.4	2
4	Comparison of nutritional and hydration status in patients undergoing twice and thrice-weekly hemodialysis: a silent drama in developing countries. <i>International Urology and Nephrology</i> , 2021, 53, 571-581.	1.4	2
5	Efficacy of metformin and fermentable fiber combination therapy in adolescents with severe obesity and insulin resistance: study protocol for a double-blind randomized controlled trial. <i>Trials</i> , 2021, 22, 148.	1.6	4
6	Effect of High-Protein Diet on Postprandial Energy Expenditure in Children with Prader-Willi Syndrome: A Pilot and Feasibility Study. <i>Current Developments in Nutrition</i> , 2021, 5, nzab016.	0.3	1
7	Frequency of Intradialytic Hypotension Events Do Not Increase with Oral Nutritional Supplementation during Hemodialysis Treatment: A Randomized Controlled Trial. , 2021, 31, 669-678.		4
8	A two-component pictured-based appetite assessment tool is capable of detecting appetite sensations in younger children: A pilot study. <i>Nutrition Research</i> , 2021, 89, 45-55.	2.9	5
9	Diet Quality Is Associated with a High Newborn Size and Reduction in the Risk of Low Birth Weight and Small for Gestational Age in a Group of Mexican Pregnant Women: An Observational Study. <i>Nutrients</i> , 2021, 13, 1853.	4.1	5
10	La adherencia al tratamiento no farmacol3gico se asocia con metas de control cardiovascular y mejores h4bitos diet3ticos en pacientes mexicanos con diabetes mellitus tipo 2. <i>Cl3nica E Investigaci3n En Arteriosclerosis</i> , 2021, , .	0.8	1
11	A dietary pattern high in full-fat dairy and sweetened beverages is associated with glycated hemoglobin and weight in Mexican patients with type-2 diabetes. <i>Nutricion Hospitalaria</i> , 2021, , .	0.3	1
12	Adherence to the DASH dietary pattern is associated with blood pressure and anthropometric indicators in Mexican adults. <i>Nutricion Hospitalaria</i> , 2021, , .	0.3	0
13	Design and Region-Specific Adaptation of the Dietary Intervention Used in the SODIUM-HF Trial: A Multicentre Study. <i>CJC Open</i> , 2020, 2, 8-14.	1.5	5
14	Adipose Tissue Development and Expansion from the Womb to Adolescence: An Overview. <i>Nutrients</i> , 2020, 12, 2735.	4.1	44
15	Alteraciones card3acas subcl3nicas detectadas por ecocardiograf3a en escolares Mexicanos con sobrepeso y obesidad. <i>Archivos De Cardiologia De Mexico</i> , 2020, 89, 222-232.	0.2	1
16	Subclinical cardiac alterations detected by echocardiography in Mexican schoolchildren with overweight and obesity. <i>Archivos De Cardiolog3a De M3xico (English Ed Internet)</i> , 2020, 89, 202-211.	0.0	0
17	Grip strength predicts cardiac adverse events in patients with cardiac disorders: an individual patient pooled meta-analysis. <i>Heart</i> , 2019, 105, 834-841.	2.9	61
18	Development of an online tool for sodium intake assessment in Mexico. <i>Revista Panamericana De Salud Publica/Pan American Journal of Public Health</i> , 2019, 43, 1.	1.1	1

#	ARTICLE	IF	CITATIONS
19	Indicators of accumulated fat are stronger associated with prehypertension compared with indicators of circulating fat. <i>Medicine (United States)</i> , 2018, 97, e11869.	1.0	11
20	Performance of Waist-To-Height Ratio, Waist Circumference, and Body Mass Index in Discriminating Cardio-Metabolic Risk Factors in a Sample of School-Aged Mexican Children. <i>Nutrients</i> , 2018, 10, 1850.	4.1	30
21	Activity of Antioxidant Enzymes and Their Association with Lipid Profile in Mexican People without Cardiovascular Disease: An Analysis of Interactions. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 2687.	2.6	10
22	Rationale and design of the Study of Dietary Intervention Under 100 MMOL in Heart Failure (SODIUM-HF). <i>American Heart Journal</i> , 2018, 205, 87-96.	2.7	11
23	Bread Consumption Is Associated with Elevated Blood Pressure among Adults Living in Mexico City—A Sub-Analysis of the Tlalpan 2020 Study. <i>Nutrients</i> , 2018, 10, 1969.	4.1	3
24	Assessment of Sodium and Potassium Intake by 24h Urinary Excretion in a Healthy Mexican Cohort. <i>Archives of Medical Research</i> , 2017, 48, 195-202.	3.3	21
25	Dietary Self-management in Heart Failure: High Tech or High Touch?. <i>Current Treatment Options in Cardiovascular Medicine</i> , 2017, 19, 19.	0.9	5
26	Protocol for a prospective longitudinal study of risk factors for hypertension incidence in a Mexico City population: the Tlalpan 2020 cohort. <i>BMJ Open</i> , 2017, 7, e016773.	1.9	13
27	Food Sources of Sodium Intake in an Adult Mexican Population: A Sub-Analysis of the SALMEX Study. <i>Nutrients</i> , 2017, 9, 810.	4.1	21
28	Salt in the diet in patients with heart failure. <i>Current Opinion in Cardiology</i> , 2016, 31, 196-203.	1.8	13
29	Changes in dietary intake and nutritional status associated with a significant reduction in sodium intake in patients with heart failure. A sub-analysis of the SODIUM-HF pilot study. <i>Clinical Nutrition ESPEN</i> , 2016, 11, e26-e32.	1.2	24
30	Maternal Underestimation of Child's Weight Status and Health Behaviors as Risk Factors for Overweight in Children. <i>Journal of Pediatric Nursing</i> , 2015, 30, e29-e33.	1.5	12
31	Association Between Self-reported Adherence to a Low-Sodium Diet and Dietary Habits Related to Sodium Intake in Heart Failure Patients. <i>Journal of Cardiovascular Nursing</i> , 2015, 30, 58-65.	1.1	25
32	Estimates of Dietary Sodium Consumption in Patients With Chronic Heart Failure. <i>Journal of Cardiac Failure</i> , 2015, 21, 981-988.	1.7	16
33	Re. "Dietary fatty acids intake and mortality in patients with heart failure" Authors' response. <i>Nutrition</i> , 2015, 31, 1185-1186.	2.4	0
34	The long-term effects of dietary sodium restriction on clinical outcomes in patients with heart failure. The SODIUM-HF (Study of Dietary Intervention Under 100 mmol in Heart Failure): A pilot study. <i>American Heart Journal</i> , 2015, 169, 274-281.e1.	2.7	53
35	Dietary fatty acids intake and mortality in patients with heart failure. <i>Nutrition</i> , 2014, 30, 1366-1371.	2.4	22
36	Dietary Fatty Acids Intake Associated With Mortality in Patients With Heart Failure. <i>Canadian Journal of Cardiology</i> , 2013, 29, S382.	1.7	0

#	ARTICLE	IF	CITATIONS
37	Cachexia assessed by bioimpedance vector analysis as a prognostic indicator in chronic stable heart failure patients. <i>Nutrition</i> , 2012, 28, 886-891.	2.4	51
38	Bioelectrical impedance phase angle as a prognostic marker in chronic heart failure. <i>Nutrition</i> , 2012, 28, 901-905.	2.4	78
39	Response to the letter to the Editor "Malnutrition syndrome, but not body mass index, is associated to worse prognosis in heart failure patients". <i>Clinical Nutrition</i> , 2012, 31, 291.	5.0	1
40	PP133-SUN WEIGHT LOSS AND ITS ASSOCIATION WITH BODY COMPOSITION AND WORSENING FUNCTIONAL CLASS IN PATIENTS WITH COMPENSATED HEART FAILURE. <i>Clinical Nutrition Supplements</i> , 2011, 6, 74-75.	0.0	0
41	Malnutrition syndrome, but not body mass index, is associated to worse prognosis in heart failure patients. <i>Clinical Nutrition</i> , 2011, 30, 753-758.	5.0	19
42	Overweight, obesity, high blood pressure and lifestyle factors among Mexican children and their parents. <i>Environmental Health and Preventive Medicine</i> , 2010, 15, 358-366.	3.4	4
43	Outcomes of a school-based intervention (RESCATE) to improve physical activity patterns in Mexican children aged 8-10 years. <i>Health Education Research</i> , 2010, 25, 1042-1049.	1.9	32
44	Prognostic value of cardiac troponin T elevation is independent of renal function and clinical findings in heart failure patients. <i>Cardiology Journal</i> , 2010, 17, 42-8.	1.2	6
45	Waist Circumference and Fat Intake Are Associated with High Blood Pressure in Mexican Children Aged 8 to 10 Years. <i>Journal of the American Dietetic Association</i> , 2009, 109, 996-1003.	1.1	48
46	No Evidence of Increased Risk for Certain Highly Atherogenic Lipoprotein Phenotypes in HIV-infected Patients. <i>Archives of Medical Research</i> , 2008, 39, 84-91.	3.3	2
47	Prevalence of right ventricular dysfunction in left-sided heart failure. <i>European Journal of Heart Failure</i> , Supplement, 2008, 7, 134-134.	0.0	0
48	Microalbuminuria in systolic and diastolic chronic heart failure patients. <i>Cardiology Journal</i> , 2008, 15, 143-9.	1.2	10
49	Human immunodeficiency virus-infected subjects have no altered myocardial perfusion. <i>International Journal of Cardiology</i> , 2007, 122, 90-92.	1.7	10
50	Bioelectrical impedance and strength measurements in patients with heart failure: comparison with functional class. <i>Nutrition</i> , 2007, 23, 412-418.	2.4	69
51	Aldosterone receptor antagonists induce favorable cardiac remodeling in diastolic heart failure patients. <i>Revista De Investigacion Clinica</i> , 2007, 59, 103-7.	0.4	27
52	Body composition and echocardiographic abnormalities associated to anemia and volume overload in heart failure patients. <i>Clinical Nutrition</i> , 2006, 25, 746-757.	5.0	27
53	Anthropometric variables and physical activity as predictors of cardiac cachexia. <i>International Journal of Cardiology</i> , 2005, 99, 239-245.	1.7	20
54	Effects of a nutritional intervention on body composition, clinical status, and quality of life in patients with heart failure. <i>Nutrition</i> , 2004, 20, 890-895.	2.4	95