
## **Toralf Reimer**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8624069/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Clinical and molecular characteristics of HER2-low-positive breast cancer: pooled analysis of individual patient data from four prospective, neoadjuvant clinical trials. Lancet Oncology, The, 2021, 22, 1151-1161.	10.7	248
2	Palbociclib for Residual High-Risk Invasive HR-Positive and HER2-Negative Early Breast Cancer—The Penelope-B Trial. Journal of Clinical Oncology, 2021, 39, 1518-1530.	1.6	153
3	Prognostic impact of androgen receptor (AR) and forkhead box A1 (FOXA1) in early HER2-negative primary breast cancer: A translational substudy of the prospective phase III WSG-PlanB-trial Journal of Clinical Oncology, 2016, 34, 557-557.	1.6	91
4	Wire localization of clip-marked axillary lymph nodes in breast cancer patients treated with primary systemic therapy. European Journal of Surgical Oncology, 2018, 44, 1307-1311.	1.0	81
5	Local Treatment of the Axilla in Early Breast Cancer: Concepts from the National Surgical Adjuvant Breast and Bowel Project B-04 to the Planned Intergroup Sentinel Mamma Trial. Breast Care, 2014, 9, 87-87.	1.4	68
6	Influence of Lifestyle Factors on Breast Cancer Risk. Breast Care, 2014, 9, 407-414.	1.4	51
7	AGO Recommendations for the Diagnosis and Treatment of Patients with Locally Advanced and Metastatic Breast Cancer: Update 2020. Breast Care, 2020, 15, 294-309.	1.4	47
8	Quality-of-Life Considerations in the Treatment of Early-Stage Breast Cancer in the Elderly. Drugs and Aging, 2010, 27, 791-800.	2.7	43
9	Diagnosing Pathologic Complete Response in the Breast After Neoadjuvant Systemic Treatment of Breast Cancer Patients by Minimal Invasive Biopsy. Annals of Surgery, 2022, 275, 576-581.	4.2	38
10	Avoiding Axillary Sentinel Lymph Node Biopsy after Neoadjuvant Systemic Therapy in Breast Cancer: Rationale for the Prospective, Multicentric EUBREAST-01 Trial. Cancers, 2020, 12, 3698.	3.7	33
11	RESPONDER – diagnosis of pathological complete response by vacuum-assisted biopsy after neoadjuvant chemotherapy in breast Cancer - a multicenter, confirmative, one-armed, intra-individually-controlled, open, diagnostic trial. BMC Cancer, 2018, 18, 851.	2.6	32
12	Angiogenic factors and acute-phase proteins in serum samples of preeclampsia and HELLP patients: a matched-pair analysis. Journal of Maternal-Fetal and Neonatal Medicine, 2013, 26, 263-269.	1.5	31
13	Intelligent Vacuum-Assisted Biopsy to Identify Breast Cancer Patients With Pathologic Complete Response (ypT0 and ypN0) After Neoadjuvant Systemic Treatment for Omission of Breast and Axillary Surgery. Journal of Clinical Oncology, 2022, 40, 1903-1915.	1.6	31
14	Is Axillary Sentinel Lymph Node Biopsy Required in Patients Who Undergo Primary Breast Surgery. Breast Care, 2018, 13, 324-330.	1.4	22
15	Central Review of Radiation Therapy Planning Among Patients with Breast-Conserving Surgery: Results from a Quality Assurance Process Integrated into the INSEMA Trial. International Journal of Radiation Oncology Biology Physics, 2020, 107, 683-693.	0.8	20
16	SLCO1B1*5 polymorphism (rs4149056) is associated with chemotherapy-induced amenorrhea in premenopausal women with breast cancer: a prospective cohort study. BMC Cancer, 2016, 16, 337.	2.6	17
17	Primary Metastatic Breast Cancer: The Impact of Locoregional Therapy. Breast Care, 2014, 9, 23-28.	1.4	16
18	Immunohistochemically Detected Lymph-Node Micrometastases in Breast Cancer and their Correlation with Prognostic Factors. Breast Journal, 1997, 3, 106-111.	1.0	15

TORALF REIMER

#	Article	IF	CITATIONS
19	Induction of apoptosis in breast cancer cells in vitro by Fas ligand reverse signaling. Journal of Cancer Research and Clinical Oncology, 2018, 144, 249-256.	2.5	15
20	The impact of anthracyclines in intermediate and high-risk HER2-negative early breast cancer—a pooled analysis of the randomised clinical trials PlanB and SUCCESS C. British Journal of Cancer, 2022, 126, 1715-1724.	6.4	14
21	False-negative rate in the extended prospective TATTOO trial evaluating targeted axillary dissection by carbon tattooing in clinically node-positive breast cancer patients receiving neoadjuvant systemic therapy. Breast Cancer Research and Treatment, 2022, 193, 589-595.	2.5	14
22	What Is the Best Management of cN0pN1(sn) Breast Cancer Patients. Breast Care, 2018, 13, 331-336.	1.4	13
23	Acute-phase proteins in prediction of preeclampsia in patients with abnormal midtrimester uterine Doppler velocimetry. Archives of Gynecology and Obstetrics, 2016, 294, 1151-1160.	1.7	11
24	How Important is the Axillary Nodal Status for Adjuvant Treatment Decisions at a Breast Cancer Multidisciplinary Tumor Board? A Survival Analysis. Annals of Surgical Oncology, 2008, 15, 472-477.	1.5	10
25	Diagnosis of Ruptured Breast Implants Through High-Resolution Ultrasound Combined With Real-Time Elastography. Aesthetic Surgery Journal, 2015, 35, 410-418.	1.6	10
26	Prospective WSG phase III PlanB trial: Final analysis of adjuvant 4xEC→4x doc vs. 6x docetaxel/cyclophosphamide in patients with high clinical risk and intermediate-to-high genomic risk HER2-negative, early breast cancer Journal of Clinical Oncology, 2017, 35, 504-504.	1.6	10
27	Distinct Ezrin Truncations Differentiate Metastases in Sentinel Lymph Nodes from Unaffected Lymph Node Tissues, from Primary Breast Tumors, and from Healthy Clandular Breast Tissues. Translational Oncology, 2018, 11, 1-10.	3.7	7
28	Radiation Sensitivity of Adipose-Derived Stem Cells Isolated from Breast Tissue. International Journal of Molecular Sciences, 2018, 19, 1988.	4.1	6
29	Lost clips after targeted lymph node biopsy in breast cancer patients: Follow-up of the CLIP-study. European Journal of Surgical Oncology, 2021, 47, 1907-1912.	1.0	5
30	A matched pair analysis of intra- and postoperative catumaxomab in patients with ovarian cancer from a multicenter, single-arm phase II trial versusÂa consecutive single-center collective of ovarian cancer patients without immunotherapy Journal of Clinical Oncology, 2012, 30, 5080-5080.	1.6	5
31	Prognostic impact of 21 Gene Recurrence Score, IHC4, and central grade in high-risk HR+/HER2- early breast cancer (EBC): 5-year results of the prospective Phase III WSG PlanB trial Journal of Clinical Oncology, 2016, 34, 556-556.	1.6	4
32	Abstract GS4-03: Patient-reported outcomes (PROs) for the intergroup sentinel mamma study (INSEMA,) Tj ETQ cancer. Cancer Research, 2022, 82, GS4-03-GS4-03.	00 0 0 rgl 0.9	BT /Overlock 1 4
33	Oncologic Outcome of Patients with (Low-Risk) Endometrial Carcinoma Undergoing Laparotomy versus Minimally Invasive Hysterectomy: A Retrospective Analysis. Oncology Research and Treatment, 2019, 42, 636-649.	1.2	2
34	Palbociclib combined with endocrine treatment in breast cancer patients with high relapse risk after neoadjuvant chemotherapy: Subgroup analyses of premenopausal patients in PENELOPE-B Journal of Clinical Oncology, 2021, 39, 518-518.	1.6	2
35	The Clinical Relevance of Target Lymph Node Biopsy after Primary Systemic Therapy in Initially Node-Positive Breast Cancer Patients. Cancers, 2021, 13, 2620.	3.7	2
36	Radiological Underestimation of Tumor Size as a Relevant Risk Factor for Positive Margin Rate in Breast-Conserving Therapy of Pure Ductal Carcinoma In Situ (DCIS). Cancers, 2022, 14, 2367.	3.7	2

TORALF REIMER

#	Article	IF	CITATIONS
37	A gap analysis of opportunities and priorities for breast surgical research. Lancet Oncology, The, 2019, 20, e1.	10.7	1
38	Prevalence, Clinical Significance and Risk Factors for Developing Scar Pain and Sensibility Disorders in Breast Cancer Patients after Breast-Conserving Therapy and Mastectomy. Breast Care, 2021, 16, 1-9.	1.4	1
39	Value of Axillary Surgery for Postoperative Therapy in Breast Cancer. Annals of Surgical Oncology, 2008, 15, 2980-2980.	1.5	0
40	Combination chemotherapy with nintedanib/placebo for patients with advanced or recurrent endometrial cancer: The NSGO ENGOT-EN1/FANDANGO trial Journal of Clinical Oncology, 2017, 35, TPS5611-TPS5611.	1.6	0
41	Abstract P5-13-36: Germline <i>BRCA</i> 1/2 and other predisposition genes in high-risk early-stage HR+/HER2- breast cancer (BC) patients treated with endocrine therapy (ET) with or without palbociclib: A secondary analysis from the PENELOPE-B study. Cancer Research, 2022, 82, P5-13-36-P5-13-36.	0.9	0
42	Abstract PD2-04: Molecular plasticity of luminal breast cancer and response to CDK 4/6 inhibition - The biomarker program of the PENELOPE-B trial investigating post-neoadjuvant palbociclib. Cancer Research, 2022, 82, PD2-04-PD2-04.	0.9	0
43	Abstract OT1-02-01: Phase III postneoadjuvant study evaluating sacituzumab govitecan (SG), an antibody drug conjugate in primary HER2-negative breast cancer patients with high relapse risk after standard neoadiuvant treatment - SASCIA. Cancer Research. 2022. 82. OT1-02-01-OT1-02-01.	0.9	0