

Amelia Brunani

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8645602/amelia-brunani-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

37
papers

761
citations

15
h-index

27
g-index

37
ext. papers

860
ext. citations

3.9
avg, IF

3.08
L-index

#	Paper	IF	Citations
37	Body composition assessment using bioelectrical impedance analysis (BIA) in a wide cohort of patients affected with mild to severe obesity. <i>Clinical Nutrition</i> , 2021 , 40, 3973-3981	5.9	1
36	Disability assessment in an Italian cohort of patients with obesity using an International Classification of Functioning, Disability and Health (ICF)-derived questionnaire. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2021 , 57, 630-638	4.4	1
35	ISPRM/ESPRM guidelines on Physical and Rehabilitation Medicine professional practice for adults with obesity and related comorbidities. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2020 , 56, 496-507	4.4	7
34	Edmonton Obesity Staging System (EOSS) and Work Ability in the Evaluation of Workers Affected by Obesity: A Preliminary Report. <i>Journal of Occupational and Environmental Medicine</i> , 2018 , 60, 732-736 ²		4
33	Defining the appropriate setting for treating obese patients: do we have the right tools?. <i>Eating and Weight Disorders</i> , 2018 , 23, 871-876	3.6	3
32	ICF-OB: a multidisciplinary questionnaire based on the International Classification of Functioning, Disability and Health to address disability in obesity. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2018 , 54, 119-121	4.4	1
31	Psychological Treatments and Psychotherapies in the Neurorehabilitation of Pain: Evidences and Recommendations from the Italian Consensus Conference on Pain in Neurorehabilitation. <i>Frontiers in Psychology</i> , 2016 , 7, 115	3.4	51
30	Psychological Considerations in the Assessment and Treatment of Pain in Neurorehabilitation and Psychological Factors Predictive of Therapeutic Response: Evidence and Recommendations from the Italian Consensus Conference on Pain in Neurorehabilitation. <i>Frontiers in Psychology</i> , 2016 , 7, 468	3.4	29
29	Patients with obesity-related comorbidities have higher disability compared with those without obesity-related comorbidities: results from a cross-sectional study. <i>International Journal of Rehabilitation Research</i> , 2016 , 39, 63-9	1.8	10
28	Short-term HIIT and Fat max training increase aerobic and metabolic fitness in men with class II and III obesity. <i>Obesity</i> , 2015 , 23, 1987-94	8	40
27	An ICF-Based Model for Implementing and Standardizing Multidisciplinary Obesity Rehabilitation Programs within the Healthcare System. <i>International Journal of Environmental Research and Public Health</i> , 2015 , 12, 6084-91	4.6	5
26	Long maximal incremental tests accurately assess aerobic fitness in class II and III obese men. <i>PLoS ONE</i> , 2015 , 10, e0124180	3.7	6
25	Rehabilitation in obesity with comorbidities: a consensus document from experts of the Italian Society of Physical and Rehabilitation Medicine (SIMFER), the Italian Society of Obesity (SIO) and the Italian Society of Eating Disorders (SISDCA). <i>Eating and Weight Disorders</i> , 2014 , 19, 383-6	3.6	3
24	Short bouts of anaerobic exercise increase non-esterified fatty acids release in obesity. <i>European Journal of Nutrition</i> , 2014 , 53, 243-9	5.2	9
23	Inherent insulin sensitivity is a major determinant of multimeric adiponectin responsiveness to short-term weight loss in extreme obesity. <i>Scientific Reports</i> , 2014 , 4, 5803	4.9	8
22	Fat oxidation, hormonal and plasma metabolite kinetics during a submaximal incremental test in lean and obese adults. <i>PLoS ONE</i> , 2014 , 9, e88707	3.7	26
21	Reference values for the 6-Min Walking Test in obese subjects. <i>Disability and Rehabilitation</i> , 2013 , 35, 1199-203	2.4	33

20	Disability affects the 6-minute walking distance in obese subjects (BMI>40 kg/m(2)). <i>PLoS ONE</i> , 2013 , 8, e75491	3.7	32
19	The Obesity-Related Disability 2013 , 55-76		3
18	Critical Aspects in Nursing 2013 , 77-106		
17	Postprandial triglyceride profile after a standardized oral fat load is altered in growth hormone (GH)-deficient adult patients and is not improved after short-term GH replacement therapy. <i>Clinical Endocrinology</i> , 2012 , 77, 721-7	3.4	1
16	Obesity is a marker of reduction in QoL and disability. <i>Scientific World Journal, The</i> , 2012 , 2012, 167520	2.2	23
15	Assessing disability in morbidly obese individuals: the Italian Society of Obesity test for obesity-related disabilities. <i>Disability and Rehabilitation</i> , 2011 , 33, 2509-18	2.4	38
14	Quality of life, disability, and body mass index are related in obese patients. <i>International Journal of Rehabilitation Research</i> , 2011 , 34, 270-2	1.8	6
13	Dynamics of GH secretion during incremental exercise in obesity, before and after a short period of training at different work-loads. <i>Clinical Endocrinology</i> , 2010 , 73, 491-6	3.4	10
12	Obesity-related disability: key factors identified by the International Classification of Functioning, Disability and Health. <i>Disability and Rehabilitation</i> , 2010 , 32, 2028-34	2.4	26
11	Mapping an obesity clinical evaluation protocol to the International Classification of Functioning, Disability and Health. <i>Disability and Rehabilitation</i> , 2010 , 32, 417-23	2.4	10
10	Use of the ICF to describe functioning and disability in obese patients. <i>Disability and Rehabilitation</i> , 2009 , 31 Suppl 1, S153-8	2.4	15
9	Sporadic mutations in melanocortin receptor 3 in morbid obese individuals. <i>European Journal of Human Genetics</i> , 2008 , 16, 581-6	5.3	54
8	Rosiglitazone is more effective than metformin in improving fasting indexes of glucose metabolism in severely obese, non-diabetic patients. <i>Diabetes, Obesity and Metabolism</i> , 2008 , 10, 460-7	6.7	5
7	Acylated ghrelin decreases during acute exercise in the lean and obese state. <i>Clinical Endocrinology</i> , 2008 , 69, 970-1	3.4	32
6	The Italian National Survey for Prader-Willi syndrome: an epidemiologic study. <i>American Journal of Medical Genetics, Part A</i> , 2008 , 146A, 861-72	2.5	65
5	Evaluation of rosiglitazone administration on cardiovascular function in severe obesity. <i>Clinical Cardiology</i> , 2008 , 31, 602-6	3.3	3
4	New insights on the simultaneous assessment of insulin sensitivity and beta-cell function with the HOMA2 method. <i>Diabetes Care</i> , 2006 , 29, 2733-4	14.6	47
3	Obesity and peripheral neuropathy risk: a dangerous liaison. <i>Journal of the Peripheral Nervous System</i> , 2005 , 10, 354-8	4.7	108

2	Short- and long-term mortality in a prevalent cohort of morbidly obese patients in Italy. <i>European Journal of Nutrition</i> , 2002 , 41, 183-5	5.2	5
1	Serum leptin levels in post-hepatitis liver cirrhosis. <i>Journal of Hepatology</i> , 2000 , 33, 38-42	13.4	41