

Michael S Engel

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8649219/publications.pdf>

Version: 2024-02-01

568
papers

11,112
citations

50244
46
h-index

69214
77
g-index

578
all docs

578
docs citations

578
times ranked

4627
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Fossiliferous Cretaceous Amber from Myanmar (Burma): Its Rediscovery, Biotic Diversity, and Paleontological Significance. American Museum Novitates, 2002, 3361, 1-71. | 0.2 | 645 |
| 2 | A MONOGRAPH OF THE BALTIC AMBER BEES AND EVOLUTION OF THE APOIDEA (HYMENOPTERA). Bulletin of the American Museum of Natural History, 2001, 259, 1-192. | 1.2 | 364 |
| 3 | New light shed on the oldest insect. Nature, 2004, 427, 627-630. | 13.7 | 252 |
| 4 | Treatise on the Isoptera of the World. Bulletin of the American Museum of Natural History, 2013, 377, 1-200. | 1.2 | 228 |
| 5 | Biogeographic and evolutionary implications of a diverse paleobiota in amber from the early Eocene of India. Proceedings of the National Academy of Sciences of the United States of America, 2010, 107, 18360-18365. | 3.3 | 184 |
| 6 | Order Hymenoptera. In: Zhang, Z.-Q. (Ed.) Animal Biodiversity: An Outline of Higher-level Classification and Survey of Taxonomic Richness (Addenda) Tj ETQq0 0 0 rgBT/Overlock 10 Tf 500 | 0.2 | 184 |
| 7 | Termites (Isoptera): Their Phylogeny, Classification, and Rise to Ecological Dominance. American Museum Novitates, 2009, 2009, 1. | 0.2 | 183 |
| 8 | The earliest known holometabolous insects. Nature, 2013, 503, 257-261. | 13.7 | 165 |
| 9 | The taxonomic impediment: a shortage of taxonomists, not the lack of technical approaches. Zoological Journal of the Linnean Society, 2021, 193, 381-387. | 1.0 | 135 |
| 10 | Evolution of lacewings and allied orders using anchored phylogenomics (<scp>N</scp>europtera,) Tj ETQq0 0 0 rgBT/Overlock 10 Tf 500 | 1.7 | 133 |
| 11 | Integrated phylogenomics and fossil data illuminate the evolution of beetles. Royal Society Open Science, 2022, 9, 211771. | 1.1 | 117 |
| 12 | A New Interpretation of the Oldest Fossil Bee (Hymenoptera: Apidae). American Museum Novitates, 2000, 3296, 1-11. | 0.2 | 113 |
| 13 | Population structure and classification of <i>Apis cerana</i>. Apidologie, 2010, 41, 589-601. | 0.9 | 110 |
| 14 | Why Descriptive Science Still Matters. BioScience, 2007, 57, 646-647. | 2.2 | 103 |
| 15 | CLASSIFICATION OF THE BEE TRIBE AUGOCHLORINI (HYMENOPTERA: HALICTIDAE). Bulletin of the American Museum of Natural History, 2000, 250, 1. | 1.2 | 101 |
| 16 | Wing stridulation in a Jurassic katydid (Insecta, Orthoptera) produced low-pitched musical calls to attract females. Proceedings of the National Academy of Sciences of the United States of America, 2012, 109, 3868-3873. | 3.3 | 100 |
| 17 | Early evolution and ecology of camouflage in insects. Proceedings of the National Academy of Sciences of the United States of America, 2012, 109, 21414-21419. | 3.3 | 93 |
| 18 | Phylogeny and Evolution of Neuropterida: Where Have Wings of Lace Taken Us?. Annual Review of Entomology, 2018, 63, 531-551. | 5.7 | 93 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 19 | Wing Tracheation in Chrysopidae and Other Neuropterida (Insecta): A Resolution of the Confusion about Vein Fusion. <i>American Museum Novitates</i> , 2017, 3890, 1-44. | 0.2 | 90 |
| 20 | Debris-carrying camouflage among diverse lineages of Cretaceous insects. <i>Science Advances</i> , 2016, 2, e1501918. | 4.7 | 87 |
| 21 | Ancient pinnate leaf mimesis among lacewings. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 16212-16215. | 3.3 | 84 |
| 22 | Family-Group Names for Bees (Hymenoptera: Apoidea). <i>American Museum Novitates</i> , 2005, 3476, 1. | 0.2 | 77 |
| 23 | Morphologically Specialized Termite Castes and Advanced Sociality in the Early Cretaceous. <i>Current Biology</i> , 2016, 26, 522-530. | 1.8 | 76 |
| 24 | Primitive New Ants in Cretaceous Amber from Myanmar, New Jersey, and Canada (Hymenoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 !). 0.2 | 72 | |
| 25 | The Neuropterid Fauna of Dominican and Mexican Amber (Neuropterida: Megaloptera, Neuroptera). <i>American Museum Novitates</i> , 2007, 3587, 1-58. | 0.2 | 69 |
| 26 | The effects of fossil placement and calibration on divergence times and rates: An example from the termites (Insecta: Isoptera). <i>Arthropod Structure and Development</i> , 2010, 39, 204-219. | 0.8 | 69 |
| 27 | Diverse transitional giant fleas from the Mesozoic era of China. <i>Nature</i> , 2012, 483, 201-204. | 13.7 | 69 |
| 28 | Diverse Cretaceous larvae reveal the evolutionary and behavioural history of antlions and lacewings. <i>Nature Communications</i> , 2018, 9, 3257. | 5.8 | 67 |
| 29 | Save Isoptera: A comment on Inward <i>et al</i> .. <i>Biology Letters</i> , 2007, 3, 562-563. | 1.0 | 65 |
| 30 | The First Mesozoic Zoraptera (Insecta). <i>American Museum Novitates</i> , 2002, 3362, 1-20. | 0.2 | 64 |
| 31 | Beetle Pollination of Cycads in the Mesozoic. <i>Current Biology</i> , 2018, 28, 2806-2812.e1. | 1.8 | 64 |
| 32 | A complete insect from the Late Devonian period. <i>Nature</i> , 2012, 488, 82-85. | 13.7 | 63 |
| 33 | Photography-based taxonomy is inadequate, unnecessary, and potentially harmful for biological sciences. <i>Zootaxa</i> , 2016, 4196, zootaxa.4196.3.9. | 0.2 | 63 |
| 34 | Termite evolution: mutualistic associations, key innovations, and the rise of Termitidae. <i>Cellular and Molecular Life Sciences</i> , 2021, 78, 2749-2769. | 2.4 | 63 |
| 35 | Phylogeny and Behavior in Honey Bees (Hymenoptera: Apidae). <i>Annals of the Entomological Society of America</i> , 1997, 90, 43-53. | 1.3 | 62 |
| 36 | Paleozoic Nymphal Wing Pads Support Dual Model of Insect Wing Origins. <i>Current Biology</i> , 2017, 27, 263-269. | 1.8 | 62 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Mexican Stingless Bees (Hymenoptera: Apidae): Diversity, Distribution, and Indigenous Knowledge. , 2013, , 135-152. | 59 | |
| 38 | Family-Group Names for Termites (Isoptera). American Museum Novitates, 2004, 3432, 1. | 0.2 | 58 |
| 39 | Insect evolution. Current Biology, 2015, 25, R868-R872. | 1.8 | 57 |
| 40 | Obtaining a better taxonomic understanding of native bees: where do we start?. Systematic Entomology, 2013, 38, 645-653. | 1.7 | 56 |
| 41 | Systematic melittology: where to from here?. Systematic Entomology, 2011, 36, 2-15. | 1.7 | 55 |
| 42 | The indigenous honey bees of Saudi Arabia (Hymenoptera, Apidae, <i>Apis mellifera jemenitica</i> Ruttner): Their natural history and role in beekeeping. ZooKeys, 2011, 134, 83-98. | 0.5 | 55 |
| 43 | Liverwort Mimesis in a Cretaceous Lacewing Larva. Current Biology, 2018, 28, 1475-1481.e1. | 1.8 | 53 |
| 44 | The Smallest Snakefly (Raphidioptera: Mesoraphidiidae): A New Species in Cretaceous Amber from Myanmar, with a Catalog of Fossil Snakeflies. American Museum Novitates, 2002, 3363, 1-22. | 0.2 | 52 |
| 45 | Family-group Names for Earwigs (Dermaptera). American Museum Novitates, 2007, 3567, 1. | 0.2 | 51 |
| 46 | The mid-Miocene Zhangpu biota reveals an outstandingly rich rainforest biome in East Asia. Science Advances, 2021, 7, . | 4.7 | 51 |
| 47 | A Diverse Paleobiota in Early Eocene Fushun Amber from China. Current Biology, 2014, 24, 1606-1610. | 1.8 | 50 |
| 48 | New fossil insect order Permopsocida elucidates major radiation and evolution of suction feeding in hemimetabolous insects (Hexapoda: Acercaria). Scientific Reports, 2016, 6, 23004. | 1.6 | 47 |
| 49 | Early Morphological Specialization for Insect-Spider Associations in Mesozoic Lacewings. Current Biology, 2016, 26, 1590-1594. | 1.8 | 47 |
| 50 | Early origin of parental care in Mesozoic carrion beetles. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, 14170-14174. | 3.3 | 45 |
| 51 | Extreme Morphogenesis and Ecological Specialization among Cretaceous Basal Ants. Current Biology, 2016, 26, 1468-1472. | 1.8 | 45 |
| 52 | <i>Leehermania prorova</i>, the Earliest Staphyliniform Beetle, from the Late Triassic of Virginia (Coleoptera: Staphylinidae). American Museum Novitates, 2012, 3761, 1-28. | 0.2 | 44 |
| 53 | Fossil Liposcelididae and the lice ages (Insecta: Psocodea). Proceedings of the Royal Society B: Biological Sciences, 2006, 273, 625-633. | 1.2 | 43 |
| 54 | Fossil honey bees and evolution in the genus <i>Apis</i> (Hymenoptera: Apidae). Apidologie, 1998, 29, 265-281. | 0.9 | 42 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 55 | Direct and indirect fossil records of megachilid bees from the Paleogene of Central Europe (Hymenoptera: Megachilidae). <i>Die Naturwissenschaften</i> , 2009, 96, 703-712. | 0.6 | 42 |
| 56 | A thorny, “anareolate” stick-insect (Phasmatidae s.l.) in Upper Cretaceous amber from Myanmar, with remarks on diversification times among Phasmatodea. <i>Cretaceous Research</i> , 2016, 63, 45-53. | 0.6 | 42 |
| 57 | Hymenoptera in Canadian Cretaceous amber (Insecta). <i>Cretaceous Research</i> , 2012, 35, 258-279. | 0.6 | 41 |
| 58 | Early Evolution of Specialized Termitophily in Cretaceous Rove Beetles. <i>Current Biology</i> , 2017, 27, 1229-1235. | 1.8 | 41 |
| 59 | Insect outbreaks produce distinctive carbon isotope signatures in defensive resins and fossiliferous ambers. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2011, 278, 3219-3224. | 1.2 | 40 |
| 60 | Geometric morphometric analysis of a new Miocene bumble bee from the Randeck Maar of southwestern Germany (Hymenoptera: Apidae). <i>Systematic Entomology</i> , 2012, 37, 784-792. | 1.7 | 40 |
| 61 | Greater past disparity and diversity hints at ancient migrations of European honey bee lineages into Africa and Asia. <i>Journal of Biogeography</i> , 2013, 40, 1832-1838. | 1.4 | 40 |
| 62 | Phase contrast X-ray synchrotron microtomography and the oldest damselflies in amber (Odonata: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5 | 1.0 | 39 |
| 63 | Blood-Feeding True Bugs in the Early Cretaceous. <i>Current Biology</i> , 2014, 24, 1786-1792. | 1.8 | 39 |
| 64 | The evolution of insect biodiversity. <i>Current Biology</i> , 2021, 31, R1299-R1311. | 1.8 | 39 |
| 65 | The Species of Isoptera (Insecta) from The Early Cretaceous Crato Formation: A Revision. <i>American Museum Novitates</i> , 2008, 3626, 1. | 0.2 | 38 |
| 66 | False fairy wasps in Early Cretaceous amber from Spain (Hymenoptera: Mymarommatoidae). <i>Palaeontology</i> , 2011, 54, 511-523. | 1.0 | 38 |
| 67 | THE MIDDLE EOCENE BEE FAUNAS OF ECKFELD AND MESSEL, GERMANY (HYMENOPTERA: APOIDEA). <i>Journal of Paleontology</i> , 2003, 77, 908-921. | 0.5 | 37 |
| 68 | Rock Crawlers in Baltic Amber (Notoptera: Mantophasmatodea). <i>American Museum Novitates</i> , 2006, 3539, 1. | 0.2 | 36 |
| 69 | Remarkable stasis in a phloeocarine rove beetle from the Late Cretaceous of New Jersey (Coleoptera, Tj ETQq1 1 0.784314 rgBT /Overlock 0.5 Tf 36 | 1.0 | 36 |
| 70 | Specialized and Generalized Pollen-Collection Strategies in an Ancient Bee Lineage. <i>Current Biology</i> , 2015, 25, 3092-3098. | 1.8 | 36 |
| 71 | Fossil record of stem groups employed in evaluating the chronogram of insects (Arthropoda: Tj ETQq1 1 0.784314 rgBT /Overlock 1.6 Tf 36 | 1.0 | 36 |
| 72 | Amphibious flies and paedomorphism in the Jurassic period. <i>Nature</i> , 2013, 495, 94-97. | 13.7 | 35 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | A New Genus of Eastern Hemisphere Stingless Bees (Hymenoptera: Apidae), with a Key to the Supraspecific Groups of Indomalayan and Australasian Meliponini. <i>American Museum Novitates</i> , 2017, 3888, 1-33. | 0.2 | 35 |
| 74 | Monophyly and extensive extinction of advanced eusocial bees: Insights from an unexpected Eocene diversity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2001, 98, 1661-1664. | 3.3 | 35 |
| 75 | The Middle Eocene bee faunas of Eckfeld and Messel, Germany (Hymenoptera: Apoidea). <i>Journal of Paleontology</i> , 2003, 77, 908-921. | 0.5 | 34 |
| 76 | The termites of Early Eocene Cambay amber, with the earliest record of the Termitidae (Isoptera). <i>ZooKeys</i> , 2011, 148, 105-123. | 0.5 | 34 |
| 77 | A defensive behavior and plant-insect interaction in Early Cretaceous amber – The case of the immature lacewing <i>Hallucinochrysa diogenesi</i> . <i>Arthropod Structure and Development</i> , 2016, 45, 133-139. | 0.8 | 34 |
| 78 | A soil-carrying lacewing larva in Early Cretaceous Lebanese amber. <i>Scientific Reports</i> , 2018, 8, 16663. | 1.6 | 34 |
| 79 | Early specializations for mimicry and defense in a Jurassic stick insect. <i>National Science Review</i> , 2021, 8, nwaa056. | 4.6 | 34 |
| 80 | Thorny lacewings (neuroptera: Rhachiberothidae) in cretaceous Amber from Myanmar. <i>Journal of Systematic Palaeontology</i> , 2004, 2, 137-140. | 0.6 | 33 |
| 81 | Basal polyphagan beetles in mid-Cretaceous amber from Myanmar: biogeographic implications and long-term morphological stasis. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2019, 286, 20182175. | 1.2 | 33 |
| 82 | A Unique Box in 28S rRNA Is Shared by the Enigmatic Insect Order Zoraptera and Dictyoptera. <i>PLoS ONE</i> , 2013, 8, e53679. | 1.1 | 32 |
| 83 | The earwigs of Kansas, with a key to genera north of Mexico (Insecta: Dermaptera). <i>Transactions of the Kansas Academy of Science</i> , 2003, 106, 115-123. | 0.0 | 31 |
| 84 | A Giant Honey Bee from the Middle Miocene of Japan (Hymenoptera: Apidae). <i>American Museum Novitates</i> , 2006, 3504, 1. | 0.2 | 31 |
| 85 | The Earliest Webspinners (Insecta: Embioidea). <i>American Museum Novitates</i> , 2006, 3514, 1. | 0.2 | 31 |
| 86 | Can higher-level phylogenies of weevils explain their evolutionary success? A critical review. <i>Systematic Entomology</i> , 2010, 35, 597-606. | 1.7 | 31 |
| 87 | Amber inclusions from New Zealand. <i>Gondwana Research</i> , 2018, 56, 135-146. | 3.0 | 31 |
| 88 | Phylogeny and Geological History of the Cynipoid Wasps (Hymenoptera: Cynipoidea). <i>American Museum Novitates</i> , 2007, 3583, 1-48. | 0.2 | 30 |
| 89 | New ant-like stone beetles in mid-Cretaceous amber from Myanmar (Coleoptera: Staphylinidae): Tj ETQql 1 0.784314 rgBT /Overlock 100.6 | | |
| 90 | A new trap-jawed ant (Hymenoptera: Formicidae: Haidomyrmecini) from Canadian Late Cretaceous amber. <i>Canadian Entomologist</i> , 2013, 145, 454-465. | 0.4 | 30 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | A NEW RECORD OF MASTOTERMES FROM THE EOCENE OF GERMANY (ISOPTERA: MASTOTERMITIDAE). Journal of Paleontology, 2006, 80, 380-385. | 0.5 | 29 |
| 92 | New Snakeflies from the Jiulongshan Formation of Inner Mongolia, China (Raphidioptera). Journal of the Kansas Entomological Society, 2008, 81, 188-193. | 0.1 | 29 |
| 93 | Primitive Termites in Cretaceous Amber from Spain and Canada (Isoptera). Journal of the Kansas Entomological Society, 2010, 83, 111-128. | 0.1 | 29 |
| 94 | Family-group names for termites (Isoptera), redux. ZooKeys, 2011, 148, 171-184. | 0.5 | 29 |
| 95 | <p>Zorotypus in Peninsular Malaysia (Zoraptera: Zorotypidae), with the description of three new species</p>. Zootaxa, 2013, 3717, 498. | 0.2 | 29 |
| 96 | New mantises (Insecta: Mantodea) in Cretaceous ambers from Lebanon, Spain, and Myanmar. Cretaceous Research, 2016, 60, 91-108. | 0.6 | 29 |
| 97 | The hatching mechanism of 130â€¢millionâ€¢yearâ€¢old insects: an association of neonates, egg shells and egg bursters in Lebanese amber. Palaeontology, 2019, 62, 547-559. | 1.0 | 29 |
| 98 | New orchid and leaf-cutter bee gynandromorphs, with an updated review (Hymenoptera, Apoidea). Zoosystematics and Evolution, 2012, 88, 205-214. | 0.4 | 28 |
| 99 | A revised definition for copal and its significance for palaeontological and Anthropocene biodiversity-loss studies. Scientific Reports, 2020, 10, 19904. | 1.6 | 28 |
| 100 | A Primitive Aphidiine Wasp in Albian Amber from Spain and a Northern Hemisphere Origin for the Subfamily (Hymenoptera: Braconidae: Aphidiinae). Journal of the Kansas Entomological Society, 2009, 82, 273-282. | 0.1 | 27 |
| 101 | Revision of the bee genus Chlerogella (Hymenoptera, Halictidae), Part I: Central American species. ZooKeys, 0, 23, 47-75. | 0.5 | 27 |
| 102 | A New Rock Crawler in Baltic Amber, with Comments on the Order(Mantophasmatodea:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 302 Td (| 0.2 | 26 |
| 103 | A New Species of Zorotypus from Central Amazonia, Brazil (Zoraptera: Zorotypidae). American Museum Novitates, 2006, 3528, 1. | 0.2 | 26 |
| 104 | Direct evidence for eudicot pollen-feeding in a Cretaceous stinging wasp (Angiospermae; Hymenoptera,) Tj ETQq0 0 0 rgBT /Overlock 10 | 2.0 | 26 |
| 105 | Straight-jawed lacewing larvae (Neuroptera) from Lower Cretaceous Spanish amber, with an account on the known amber diversity of neuropterid immatures. Cretaceous Research, 2020, 106, 104200. | 0.6 | 26 |
| 106 | Gregarious behaviour in Cretaceous earwig nymphs (Insecta, Dermaptera) from southwestern France. Geodiversitas, 2009, 31, 129-135. | 0.2 | 25 |
| 107 | Diverse stigmaphroniid wasps in Early Cretaceous amber from Spain (Hymenoptera: Ceraphronoidea:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 | 0.6 | 25 |
| 108 | A New Lineage of Enigmatic Diaprioid Wasps in Cretaceous Amber (Hymenoptera: Diapriidae). American Museum Novitates, 2013, 3771, 1-23. | 0.2 | 25 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | A remarkable evanioid wasp in mid-Cretaceous amber from northern Myanmar (Hymenoptera: Tj ETQql 1 0.784314 rgBT /Overlock 10 | 0.6 | 25 |
| 110 | Insect Wings: The Evolutionary Development of Natureâ€™s First Flyers. , 2013, , 269-298. | | 25 |
| 111 | A PRIMITIVE EARWIG IN CRETACEOUS AMBER FROM MYANMAR (DERMAPTERA: PYGIDICRANIDAE). Journal of Paleontology, 2004, 78, 1018-1023. | 0.5 | 24 |
| 112 | Fig Wasps in Dominican Amber (Hymenoptera: Agaonidae). American Museum Novitates, 2006, 3541, 1. | 0.2 | 24 |
| 113 | Potential distribution of orchid bees outside their native range: The cases of <i>Eulaema polychroma</i> (MocsÃ¡ry) and <i>Euglossa viridissima</i> Friese in the USA (Hymenoptera: Apidae). Diversity and Distributions, 2009, 15, 421-428. | 1.9 | 24 |
| 114 | Description of an ancient social bee trapped in amber using diagnostic radioentomology. Insectes Sociaux, 2011, 58, 487-494. | 0.7 | 24 |
| 115 | Three new cryptic species of Euglossa from Brazil (Hymenoptera, Apidae). ZooKeys, 2012, 222, 47-68. | 0.5 | 24 |
| 116 | Weevils of the Yixian Formation, China (Coleoptera: Curculionoidea): phylogenetic considerations and comparison with other Mesozoic faunas. Journal of Systematic Palaeontology, 2013, 11, 399-429. | 0.6 | 24 |
| 117 | An evolutionary history embedded in amber: reflection of the Mesozoic shift in weevil-dominated (Coleoptera: Curculionoidea) faunas. Zoological Journal of the Linnean Society, 2014, 171, 534-553. | 1.0 | 24 |
| 118 | Fire-prone Rhamnaceae with South African affinities in Cretaceous Myanmar amber. Nature Plants, 2022, 8, 125-135. | 4.7 | 24 |
| 119 | A new xeromelissine bee in Tertiary amber of the Dominican Republic (Hymenoptera: Colletidae). Insect Systematics and Evolution, 1999, 30, 453-458. | 0.2 | 23 |
| 120 | Cretaceous Scolebythidae and Phylogeny of the Family (Hymenoptera: Chryridoidea). American Museum Novitates, 2007, 3568, 1. | 0.2 | 23 |
| 121 | Serphitid wasps in Early Cretaceous amber from Spain (Hymenoptera: Serphitidae). Cretaceous Research, 2011, 32, 143-154. | 0.6 | 23 |
| 122 | Miocene honey bees from the Randeck Maar of southwestern Germany (Hymenoptera, Apidae). ZooKeys, 2011, 96, 11-37. | 0.5 | 23 |
| 123 | Mesozoic giant fleas from northeastern China (Siphonaptera): Taxonomy and implications for palaeodiversity. Science Bulletin, 2013, 58, 1682-1690. | 1.7 | 23 |
| 124 | Morphological phylogeny of Megachilini and the evolution of leaf-cutter behavior in bees (Hymenoptera: Megachilidae). Journal of Melittology, 2019, , 1-123. | 0.2 | 23 |
| 125 | Wing Shape of Four New Bee Fossils (Hymenoptera: Anthophila) Provides Insights to Bee Evolution. PLoS ONE, 2014, 9, e108865. | 1.1 | 23 |
| 126 | Morphometric analysis of fossil bumble bees (Hymenoptera, Apidae, Bombini) reveals their taxonomic affinities. ZooKeys, 2019, 891, 71-118. | 0.5 | 23 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 127 | Late Carboniferous paleoichnology reveals the oldest full-body impression of a flying insect. Proceedings of the National Academy of Sciences of the United States of America, 2011, 108, 6515-6519. | 3.3 | 22 |
| 128 | A History of Entomological Classification. Annual Review of Entomology, 2013, 58, 585-607. | 5.7 | 22 |
| 129 | The first Mesozoic microwhip scorpion (Palpigradi): a new genus and species in mid-Cretaceous amber from Myanmar. Die Naturwissenschaften, 2016, 103, 19. | 0.6 | 22 |
| 130 | A remarkable diversity of parasitoid beetles (Ripiphoridae) in Cretaceous amber, with a summary of the Mesozoic record of Tenebrionoidea. Cretaceous Research, 2018, 90, 296-310. | 0.6 | 22 |
| 131 | Myanmar: palaeontologists must stop buying conflict amber. Nature, 2020, 584, 525-525. | 13.7 | 22 |
| 132 | A remarkable kalligrammatid lacewing from the Upper Jurassic of Kazakhstan (Neuroptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 542 Tc 0.0 21 | | |
| 133 | <I>Permocoleus</I>, New Genus, the First Permian Beetle (Coleoptera) from North America. Annals of the Entomological Society of America, 2005, 98, 73-76. | 1.3 | 21 |
| 134 | The earliest earwigs in amber (Dermaptera): A new genus and species from the Early Cretaceous of Lebanon. Insect Systematics and Evolution, 2011, 42, 139-148. | 0.2 | 21 |
| 135 | Snakefly diversity in Early Cretaceous amber from Spain (Neuropterida, Raphidioptera). ZooKeys, 2012, 204, 1-40. | 0.5 | 21 |
| 136 | Ichneumonidae (Insecta: Hymenoptera) in Canadian Late Cretaceous amber. Fossil Record, 2013, 16, 217-227. | 0.4 | 21 |
| 137 | A primer of host-plant specialization in bees. Emerging Topics in Life Sciences, 2020, 4, 7-17. | 1.1 | 21 |
| 138 | A Diminutive Pelecinid Wasp in Cretaceous Amber from New Jersey (Hymenoptera: Pelecinidae). Northeastern Naturalist, 2006, 13, 291-297. | 0.1 | 20 |
| 139 | New false fairy wasps in Cretaceous amber from New Jersey and Myanmar (Hymenoptera: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 0.0 20 | | |
| 140 | The enigmatic Mesozoic insect taxon Chresmodidae (Polyneoptera): New palaeobiological and phylogenetic data, with the description of a new species from the Lower Cretaceous of Brazil. Neues Jahrbuch Fur Geologie Und Palaontologie - Abhandlungen, 2008, 247, 353-381. | 0.2 | 20 |
| 141 | A Protorhyssaline Wasp in Early Cretaceous Amber from Spain (Hymenoptera: Braconidae). Journal of the Kansas Entomological Society, 2011, 84, 51-57. | 0.1 | 20 |
| 142 | Social Bees and the Current Status of Beekeeping in Indonesia. , 2018, , 287-306. | | 20 |
| 143 | Male sleeping aggregations of solitary oil-collecting bees in Brazil (Centridini, Tapinotaspidini, and) Tj ETQq1 1 0.784314 rgBT /Overlock 0.3 20 | | |
| 144 | An Eocene Bee in Rovno Amber, Ukraine (Hymenoptera: Megachilidae). American Museum Novitates, 2006, 3506, 1. | 0.2 | 19 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | The First Cretaceous Spider Wasp (Hymenoptera: Pompilidae). <i>Journal of the Kansas Entomological Society</i> , 2006, 79, 359-368. | 0.1 | 19 |
| 146 | A New Fossil Orchid Bee in Colombian Copal (Hymenoptera: Apidae). <i>American Museum Novitates</i> , 2007, 3589, 1-7. | 0.2 | 19 |
| 147 | Aetheogrammatidae, A New Family of Lacewings from the Mesozoic of China (Neuroptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 38 | 0.1 | 19 |
| 148 | A giant termite from the Late Miocene of Styria, Austria (Isoptera). <i>Die Naturwissenschaften</i> , 2009, 96, 289-295. | 0.6 | 19 |
| 149 | A New Thorny Lacewing (Neuroptera: Rhachiberothidae) from Canadian Cretaceous Amber. <i>Journal of the Kansas Entomological Society</i> , 2009, 82, 114-121. | 0.1 | 19 |
| 150 | Two new species of Euglossa from South America, with notes on their taxonomic affinities (Hymenoptera, Apidae). <i>ZooKeys</i> , 2012, 221, 63-79. | 0.5 | 19 |
| 151 | Long-term stasis in a diverse fauna of Early Cretaceous springtails (Collembola: Symphyleona). <i>Journal of Systematic Palaeontology</i> , 2017, 15, 513-537. | 0.6 | 19 |
| 152 | A key to the genera and subgenera of stingless bees in Indonesia (Hymenoptera: Apidae). <i>Treubia</i> , 0, 45, 65-84. | 0.1 | 19 |
| 153 | Extreme adaptations for aquatic ectoparasitism in a Jurassic fly larva. <i>ELife</i> , 2014, 3, e02844. | 2.8 | 19 |
| 154 | A new augochlorine bee species in Tertiary amber from the Dominican Republic (Hymenoptera:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 38 | 0.9 | 18 |
| 155 | The First Cretaceous Sclerogibbid Wasp (Hymenoptera: Sclerogibbidae). <i>American Museum Novitates</i> , 2006, 3515, 1. | 0.2 | 18 |
| 156 | Dryophthorine weevils in Dominican amber (Coleoptera: Curculionidae). <i>Transactions of the Kansas Academy of Science</i> , 2006, 109, 191-198. | 0.0 | 18 |
| 157 | A Lateral Gynandromorph in the Bee Genus Thyreus and the Sting Mechanism in the Melectini (Hymenoptera: Apidae). <i>American Museum Novitates</i> , 2007, 3553, 1. | 0.2 | 18 |
| 158 | Two New Termites in Baltic Amber (Isoptera). <i>Journal of the Kansas Entomological Society</i> , 2008, 81, 194-203. | 0.1 | 18 |
| 159 | New Data on Homocladus grandis, A Permian Stem-Mantodean (Polyneoptera: Dictyoptera). <i>Journal of Paleontology</i> , 2010, 84, 746-753. | 0.5 | 18 |
| 160 | Fossil bees and their plant associates. , 2011, , 103-164. | | 18 |
| 161 | New and revised maimetshid wasps from Cretaceous ambers (Hymenoptera, Maimetshidae). <i>ZooKeys</i> , 2011, 130, 421-453. | 0.5 | 18 |
| 162 | The large carpenter bees of central Saudi Arabia, with notes on the biology of Xylocopa sulcatipes Maa (Hymenoptera, Apidae, Xylocopinae). <i>ZooKeys</i> , 2012, 201, 1-14. | 0.5 | 18 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | Springtails from the Early Cretaceous Amber of Spain (Collembola: Entomobryomorpha), with an Annotated Checklist of Fossil Collembola. American Museum Novitates, 2016, 3862, 1-47. | 0.2 | 18 |
| 164 | Olfactory associative behavioral differences in three honey bee <i>Apis mellifera</i> L. races under the arid zone ecosystem of central Saudi Arabia. Saudi Journal of Biological Sciences, 2019, 26, 563-568. | 1.8 | 18 |
| 165 | Balance scientific and ethical concerns to achieve a nuanced perspective on “blood amber”. Nature Ecology and Evolution, 2021, 5, 705-706. | 3.4 | 18 |
| 166 | The Asian Species of <i>Apis</i> . , 2011, , 1-22. | | 18 |
| 167 | A new fossil bee from the Oligo-Miocene Dominican amber (Hymenoptera:Halictidae). Apidologie, 1997, 28, 97-102. | 0.9 | 18 |
| 168 | The Dustywings in cretaceous Burmese amber (Insecta: Neuroptera: Coniopterygidae). Journal of Systematic Palaeontology, 2004, 2, 133-136. | 0.6 | 17 |
| 169 | <i>Euglossa obrima</i> , a new species of orchid bee from Mesoamerica, with notes on the subgenus <i>Dasystilbe</i> Dressler (Hymenoptera, Apidae). ZooKeys, 2011, 97, 11-29. | 0.5 | 17 |
| 170 | New scolebythid wasps in Cretaceous amber from Spain and Canada, with implications for the phylogeny of the family (Hymenoptera: Scolebythidae). Cretaceous Research, 2013, 46, 31-42. | 0.6 | 17 |
| 171 | Lower Cretaceous origin of long-distance mate finding behaviour in Hymenoptera (Insecta). Journal of Systematic Palaeontology, 2013, 11, 83-89. | 0.6 | 17 |
| 172 | The bees of Early Eocene Cambay amber (Hymenoptera: Apidae). Journal of Melittology, 2013, , 1-12. | 0.2 | 17 |
| 173 | A new interpretation of the bee fossil <i>Melitta willardi</i> Cockerell (Hymenoptera, Melittidae) based on geometric morphometrics of the wing. ZooKeys, 2014, 389, 35-48. | 0.5 | 17 |
| 174 | A new twisted-wing parasitoid from mid-Cretaceous amber of Myanmar (Strepsiptera). Cretaceous Research, 2016, 58, 160-167. | 0.6 | 17 |
| 175 | Palaeozoic giant dragonflies were hawker predators. Scientific Reports, 2018, 8, 12141. | 1.6 | 17 |
| 176 | Evolution of green lacewings (Neuroptera: Chrysopidae): an anchored phylogenomics approach. Systematic Entomology, 2019, 44, 514-526. | 1.7 | 17 |
| 177 | A review of the Indo-Malayan meliponine genus <i>Lisotrigona</i> , with two new species (Hymenoptera: Tj ETQq1 0.1 0.784314 rgBT /Ov | | |
| 178 | Extralimital Fossils of the “Gondwanan” Family Sphaeropsocidae (Insecta: Psocodea). American Museum Novitates, 2006, 3523, 1. | 0.2 | 16 |
| 179 | Revision of the Bee Genus <i>Chlerogella</i> (Hymenoptera, Halictidae), Part II: South American Species and Generic Diagnosis. ZooKeys, 2010, 47, 1-100. | 0.5 | 16 |
| 180 | A New Species of <i>Megachile</i> from Pakistan, with Taxonomic Notes on the Subgenus <i>Eutricharaea</i> (Hymenoptera: Megachilidae). Journal of the Kansas Entomological Society, 2010, 83, 58-67. | 0.1 | 16 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 181 | New earwig nymphs (Dermaptera: Pygidicranidae) in mid-Cretaceous amber from France. <i>Cretaceous Research</i> , 2011, 32, 325-330. | 0.6 | 16 |
| 182 | New earwigs in mid-Cretaceous amber from Myanmar (Dermaptera, Neodermaptera). <i>ZooKeys</i> , 2011, 130, 137-152. | 0.5 | 16 |
| 183 | Revision of the orchid bee subgenus <i>Euglossella</i> (Hymenoptera, Apidae), Part I, The <i>decorata</i> species group. <i>ZooKeys</i> , 2011, 140, 27-69. | 0.5 | 16 |
| 184 | Coniopterygidae (Neuroptera: Aleuropteryginae) in Amber from the Eocene of India and the Miocene of Hispaniola. <i>American Museum Novitates</i> , 2013, 3770, 20-39. | 0.2 | 16 |
| 185 | An unusual new lineage of sawflies (Hymenoptera) in Upper Cretaceous amber from northern Myanmar. <i>Cretaceous Research</i> , 2016, 60, 281-286. | 0.6 | 16 |
| 186 | Haplotype diversity and genetic similarity among populations of the Eastern honey bee from Himalaya-Southwest China and Nepal (Hymenoptera: Apidae). <i>Apidologie</i> , 2016, 47, 197-205. | 0.9 | 16 |
| 187 | A new genus and species of pygidicranid earwigs from the Upper Cretaceous of southern Asia (Dermaptera: Pygidicranidae). <i>Cretaceous Research</i> , 2017, 69, 178-183. | 0.6 | 16 |
| 188 | Two new ripidiine species in Dominican amber with evidence of aggregative behaviour of males "frozen" in the fossil record (Coleoptera: Ripiphoridae). <i>European Journal of Entomology</i> , 2011, 108, 275-286. | 1.2 | 16 |
| 189 | A New Moustache Wasp in Dominican Amber, with an Account of Apoid Wasp Evolution Emphasizing Crabroninae (Hymenoptera: Crabronidae). <i>American Museum Novitates</i> , 2006, 3529, 1. | 0.2 | 15 |
| 190 | Two Wasp Families Rare in the Fossil Record (Hymenoptera): Perilampidae and Megaspilidae from the Miocene of Spain. <i>American Museum Novitates</i> , 2006, 3540, 1. | 0.2 | 15 |
| 191 | Hennigian Phylogenetic Systematics and the "Groundplan" vs. "Post-Groundplan" Approaches: A Reply to KukalovÁji-Peck. <i>Evolutionary Biology</i> , 2008, 35, 317-323. | 0.5 | 15 |
| 192 | Webspinners in Early Eocene amber from western India (Insecta, Embiodea). <i>ZooKeys</i> , 2011, 148, 197-208. | 0.5 | 15 |
| 193 | New Stigmaphronidae and Megaspilidae (Hymenoptera: Ceraphronoidea) from Canadian Cretaceous amber. <i>Cretaceous Research</i> , 2011, 32, 794-805. | 0.6 | 15 |
| 194 | New bethylid and chrysidid wasps (Hymenoptera: Chrysididae) from Canadian Late Cretaceous amber. <i>Palaontologische Zeitschrift</i> , 2014, 88, 433-451. | 0.8 | 15 |
| 195 | An earwig (Insecta: Dermaptera) in Early Cretaceous amber from Spain. <i>Insect Systematics and Evolution</i> , 2015, 46, 291-300. | 0.2 | 15 |
| 196 | Taxonomic description of <i>< i>in situ</i></i> bee pollen from the middle Eocene of Germany. <i>Grana</i> , 2017, 56, 37-70. | 0.4 | 15 |
| 197 | Images are not and should not ever be type specimens: a rebuttal to Garraffoni & Freitas. <i>Zootaxa</i> , 2017, 4269, 455-459. | 0.2 | 15 |
| 198 | Large expert-curated database for benchmarking document similarity detection in biomedical literature search. Database: the Journal of Biological Databases and Curation, 2019, 2019, . | 1.4 | 15 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 199 | A Note on the Taxonomy of Some Fossil Bees from Germany (Hymenoptera: Apidae). <i>Journal of the Kansas Entomological Society</i> , 2005, 78, 82. | 0.1 | 15 |
| 200 | An Early Eocene bee (Hymenoptera: Halictidae) from Quilchena, British Columbia. <i>Canadian Entomologist</i> , 2003, 135, 63-69. | 0.4 | 14 |
| 201 | The Wasp Family Rhopalosomatidae in Mid-Cretaceous Amber from Myanmar (Hymenoptera: Vespoidea). <i>Journal of the Kansas Entomological Society</i> , 2008, 81, 168-174. | 0.1 | 14 |
| 202 | Eocene tortoise beetles from the Green River Formation in Colorado, U.S.A. (Coleoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 622 Td (1.7 | 1.7 | 14 |
| 203 | The Wasp Family Embolemidae in Early Cretaceous Amber from Spain (Hymenoptera: Chrysididae). <i>Journal of the Kansas Entomological Society</i> , 2011, 84, 36-42. | 0.1 | 14 |
| 204 | The serphitid wasps (Hymenoptera: Proctotrupo- morpha Serphitoidea) of Canadian Cretaceous amber. <i>Systematic Entomology</i> , 2011, 36, 192-208. | 1.7 | 14 |
| 205 | A new species of Chalicodoma from Saudi Arabia with modified facial setae (Hymenoptera, Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 622 Td (0.5 | 0.5 | 14 |
| 206 | The Relict Scorpionfly Family Meropeidae (Mecoptera) in Cretaceous Amber. <i>Journal of the Kansas Entomological Society</i> , 2013, 86, 253-263. | 0.1 | 14 |
| 207 | The Fauna of Staphylininae in Dominican Amber (Coleoptera: Staphylinidae). <i>Annals of Carnegie Museum</i> , 2013, 81, 281-294. | 0.1 | 14 |
| 208 | Bugs in the Biogeography: Leptosaldinae (Heteroptera: Leptopodidae) in Amber from the Miocene of Hispaniola and Eocene of India. <i>Journal of the Kansas Entomological Society</i> , 2013, 86, 226-243. | 0.1 | 14 |
| 209 | The Mesozoic family Archegocimicidae and phylogeny of the infraorder Leptopodomorpha (Hemiptera). <i>Journal of Systematic Palaeontology</i> , 2014, 12, 93-111. | 0.6 | 14 |
| 210 | New Carboniferous fossils of Spilapteridae enlighten postembryonic wing development in Palaeodictyoptera. <i>Systematic Entomology</i> , 2016, 41, 178-190. | 1.7 | 14 |
| 211 | A new subgenus of <i>Heterotrigona</i> from New Guinea (Hymenoptera: Apidae). <i>Journal of Melittology</i> , 2017, , 1-16. | 0.2 | 14 |
| 212 | A New Subgenus of Megachile from Borneo with Arolia (Hymenoptera: Megachilidae). <i>American Museum Novitates</i> , 2006, 3505, 1. | 0.2 | 14 |
| 213 | Drywood termites in Dominican amber (Isoptera: Kalotermitidae).. <i>Contributions To Entomology</i> , 2007, 57, 263-275. | 0.1 | 14 |
| 214 | A New Fossil Snake-Fly Species from Baltic Amber (Raphidioptera: Inocelliidae). <i>Psyche: Journal of Entomology</i> , 1995, 102, 187-193. | 0.4 | 13 |
| 215 | THE FIRST MESOZOIC STEPHANID WASP (HYMENOPTERA: STEPHANIDAE). <i>Journal of Paleontology</i> , 2004, 78, 1192-1197. | 0.5 | 13 |
| 216 | An Eocene ectoparasite of bees: The oldest definitive record of phoretic meloid triungulins (Coleoptera: Meloidae; Hymenoptera: Megachilidae). <i>Acta Zoologica Cracoviensia</i> , 2005, 48, 43-48. | 0.3 | 13 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 217 | A note on the relic silverfish <i>Tricholepidion gertschi</i> (Zygentoma). <i>Transactions of the Kansas Academy of Science</i> , 2006, 109, 236-238. | 0.0 | 13 |
| 218 | A New Lower Permian Bristletail from the Wellington Formation in Kansas (Archaeognatha: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 702 T | 0.0 | 13 |
| 219 | A new species of <i>Diochus</i> from Baltic amber (Coleoptera, Staphylinidae, Diochini). <i>ZooKeys</i> , 2011, 138, 65-73. | 0.5 | 13 |
| 220 | An Exomalopsine Bee in Early Miocene Amber from the Dominican Republic (Hymenoptera: Apidae). <i>American Museum Novitates</i> , 2012, 3758, 1-16. | 0.2 | 13 |
| 221 | A preliminary list of bee genera in the Kingdom of Saudi Arabia (Hymenoptera: Apoidea). <i>Journal of the Saudi Society of Agricultural Sciences</i> , 2013, 12, 85-89. | 1.0 | 13 |
| 222 | A new genus of nemonychid weevil from Burmese Amber (Coleoptera, Curculionoidea). <i>ZooKeys</i> , 2014, 405, 127-138. | 0.5 | 13 |
| 223 | A Revised Key to the Living and Fossil Families of Strepsiptera, with the Description of a New Family, Cretostylopidae. <i>Journal of the Kansas Entomological Society</i> , 2014, 87, 385-388. | 0.1 | 13 |
| 224 | Before the â€˜Big Chillâ€™: A preliminary overview of arthropods from the middle Miocene of Iceland (Insecta, Crustacea). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2014, 401, 1-12. | 1.0 | 13 |
| 225 | The first mastotermitid termite from Africa (Isoptera: Mastotermitidae): a new species of <i>Mastotermes</i> from the early Miocene of Ethiopia. <i>Journal of Paleontology</i> , 2015, 89, 1038-1042. | 0.5 | 13 |
| 226 | A New Bethylid Wasp in Lebanese Early Cretaceous Amber (Hymenoptera: Chrysidoidea), with Comments on other Mesozoic Taxa. <i>American Museum Novitates</i> , 2016, 3855, 1-14. | 0.2 | 13 |
| 227 | The first arripeneurine antlion in Burmese amber (Neuroptera: Myrmeleontidae). <i>Cretaceous Research</i> , 2016, 63, 1-6. | 0.6 | 13 |
| 228 | Antiquity of cleptoparasitism among bees revealed by morphometric and phylogenetic analysis of a <scp>P</scp>aleocene fossil nomadine (<scp>H</scp>ymenoptera: <scp>A</scp>pidae). <i>Systematic Entomology</i> , 2017, 42, 543-554. | 1.7 | 13 |
| 229 | Colonizing the east and the west: distribution and niche properties of a dwarf Asian honey bee invading Africa, the Middle East, the Malay Peninsula, and Taiwan. <i>Apidologie</i> , 2020, 51, 75-87. | 0.9 | 13 |
| 230 | Early Cretaceous Snakefly Larvae in Amber from Lebanon, Myanmar, and France (Raphidioptera). <i>American Museum Novitates</i> , 2007, 3598, 1. | 0.2 | 13 |
| 231 | Diverse scelionid wasps in Early Cretaceous amber from Spain (Hymenoptera: Platygastroidea). <i>Bulletin of Geosciences</i> , 2014, , 553-571. | 0.5 | 13 |
| 232 | Two new orchid bees of the subgenus <i>Euglossella</i> from Peru (Hymenoptera: Apidae).. <i>Contributions To Entomology</i> , 2007, 57, 93-104. | 0.1 | 13 |
| 233 | <i>Ischnomelissa</i> , a New Genus of Augochlorine Bees (Halictidae) from Colombia. <i>Studies on Neotropical Fauna and Environment</i> , 1997, 32, 41-46. | 0.5 | 12 |
| 234 | Fideliine Phylogeny and Classification Revisited (Hymenoptera: Megachilidae). <i>Journal of the Kansas Entomological Society</i> , 2004, 77, 821-836. | 0.1 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 235 | A primitive sapygid wasp in Burmese amber (Hymenoptera: Sapygidae). <i>Acta Zoologica Cracoviensia</i> , 2005, 48, 1-9. | 0.3 | 12 |
| 236 | An Unusual, Primitive Piesmatidae (Insecta: Heteroptera) in Cretaceous Amber from Myanmar (Burma). <i>American Museum Novitates</i> , 2008, 3611, 1. | 0.2 | 12 |
| 237 | Notes on the Nesting Biology of the Small Carpenter Bee< i>Ceratina smaragdula</i> (Hymenoptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 0.2 12 | | |
| 238 | The first Oriental protorhyssaline wasp (Hymenoptera: Braconidae): A new genus and species in Upper Cretaceous amber from Myanmar. <i>Cretaceous Research</i> , 2016, 63, 28-32. | 0.6 | 12 |
| 239 | Two new species of mid-Cretaceous webspinners in amber from northern Myanmar (Embiodea:) Tj ETQq1 1 0.784314 rgBT /Overlock 0.6 12 | | |
| 240 | Discovery of the Bee Tribe Tarsaliini in Arabia (Hymenoptera: Apidae), with the Description of a New Species. <i>American Museum Novitates</i> , 2017, 3877, 1-28. | 0.2 | 12 |
| 241 | Notes on Southeast Asian Stingless Bees of the Genus< i>Tetragonula</i> (Hymenoptera: Apidae), with the Description of a New Species from Thailand. <i>American Museum Novitates</i> , 2017, 3886, 1-20. | 0.2 | 12 |
| 242 | Palaeodictyopterida. <i>Current Biology</i> , 2019, 29, R306-R309. | 1.8 | 12 |
| 243 | Widespread mimicry and camouflage among mid-Cretaceous insects. <i>Gondwana Research</i> , 2022, 101, 94-102. | 3.0 | 12 |
| 244 | Fossil snakeflies from the Early Cretaceous of southern Korea (Raphidioptera: Mesoraphidiidae). <i>Neues Jahrbuch FÄl4r Geologie Und PalÄontologie</i> , 2006, 2006, 249-256. | 0.3 | 12 |
| 245 | Halictine bees from the Eocene-Oligocene boundary of Florissant, Colorado (Hymenoptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 5 12 | | |
| 246 | Molecular Phylogeny Reveals the Past Transoceanic Voyages of Drywood Termites (Isoptera,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 302 | | |
| 247 | A new genus and species of Acaridae (Acar) phoretic onThectochlora alaris(Hymenoptera: Halictidae:) Tj ETQq1 1 0.784314 rgBT /Overlock 0.3 11 | | |
| 248 | The alderflies of Kansas (Megaloptera: Sialidae). <i>Transactions of the Kansas Academy of Science</i> , 2004, 107, 119-125. | 0.0 | 11 |
| 249 | New Dolichorhinotermes from Ecuador and in Mexican Amber (Isoptera: Rhinotermitidae). <i>American Museum Novitates</i> , 2007, 3592, 1-8. | 0.2 | 11 |
| 250 | A new species of Zorotypus from eastern Amazonia, Brazil (Zoraptera: Zorotypidae). <i>Transactions of the Kansas Academy of Science</i> , 2008, 111, 193-202. | 0.0 | 11 |
| 251 | A Second Specimen of <i>Permocoleus</i> (Coleoptera) from the Lower Permian Wellington Formation of Noble County, Oklahoma. <i>Journal of the Kansas Entomological Society</i> , 2008, 81, 4-7. | 0.1 | 11 |
| 252 | Maternally inherited architecture in tertiary leaf beetles: paleoichnology of cryptocephaline fecal cases in Dominican and Baltic amber. <i>Die Naturwissenschaften</i> , 2009, 96, 1121-1126. | 0.6 | 11 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 253 | An Aquatic Water Scavenger Beetle in Early Miocene Amber from the Dominican Republic (Coleoptera: Tj ETQq1 10.1784314rgBT /Ove | 0.1 | 11 |
| 254 | Euglossa williamsi, a new species of orchid bee from the Amazon Basin of Ecuador and Peru, with notes on its taxonomic association and biogeography (Hymenoptera, Apidae). ZooKeys, 2011, 159, 49-63. | 0.5 | 11 |
| 255 | A Stephanid Wasp in Mid-Cretaceous Burmese Amber (Hymenoptera: Stephanidae), with Comments on the Antiquity of the Hymenopteran Radiation. Journal of the Kansas Entomological Society, 2013, 86, 244-252. | 0.1 | 11 |
| 256 | Bethylidae from Early Cretaceous Spanish Amber (Hymenoptera: Chrysididae). Journal of the Kansas Entomological Society, 2013, 86, 264-276. | 0.1 | 11 |
| 257 | A minute stingless bee in Eocene Fushan amber from northeastern China (Hymenoptera: Apidae). Journal of Melittology, 2013, , 1-10. | 0.2 | 11 |
| 258 | Nesting Biology of the Leafcutting Bee <i>Megachile minutissima</i> (Hymenoptera: Megachilidae) in Central Saudi Arabia. Annals of the Entomological Society of America, 2014, 107, 635-640. | 1.3 | 11 |
| 259 | New ripiphorid beetles in mid-Cretaceous amber from Myanmar (Coleoptera: Ripiphoridae): First Pelecotominae and possible Mesozoic aggregative behaviour in male Ripidiinae. Cretaceous Research, 2016, 68, 70-78. | 0.6 | 11 |
| 260 | Diverse, primitive termites (Isoptera: Kalotermitidae, incertae sedis) from the early Miocene of New Zealand. Austral Entomology, 2017, 56, 94-103. | 0.8 | 11 |
| 261 | Ecomorphological diversification of the Late Palaeozoic Palaeodictyopterida reveals different larval strategies and amphibious lifestyle in adults. Royal Society Open Science, 2019, 6, 190460. | 1.1 | 11 |
| 262 | Evolution of green lacewings (Neuroptera: Chrysopidae): a molecular supermatrix approach. Systematic Entomology, 2019, 44, 499-513. | 1.7 | 11 |
| 263 | Cretaceous diversity and disparity in a lacewing lineage of predators (Neuroptera: Mantispidae). Proceedings of the Royal Society B: Biological Sciences, 2020, 287, 20200629. | 1.2 | 11 |
| 264 | Fossil Social Insects. , 2020, , 1-21. | | 11 |
| 265 | An Early Miocene bumble bee from northern Bohemia (Hymenoptera, Apidae). ZooKeys, 2017, 710, 43-63. | 0.5 | 11 |
| 266 | The wasp larva's last supper: 100 million years of evolutionary stasis in the larval development of rhopalosomatid wasps (Hymenoptera: Rhopalosomatidae). Fossil Record, 2017, 20, 239-244. | 0.5 | 11 |
| 267 | Phylogeny, biogeography and classification of Teletiosoptera (Blattaria: Isoptera). Systematic Entomology, 2022, 47, 581-590. | 1.7 | 11 |
| 268 | A new genus of cleptoparasitic bees from the West Indies (Hymenoptera: Halictidae). Acta Zoologica Cracoviensis - Series B: Invertebrata, 2006, 49, 1-8. | 0.2 | 10 |
| 269 | A Termite Bug in Early Miocene Amber of the Dominican Republic (Hemiptera: Termitaphididae). American Museum Novitates, 2008, 3619, 1. | 0.2 | 10 |
| 270 | AN OXYTELINE ROVE BEETLE IN DOMINICAN AMBER WITH POSSIBLE AFRICAN AFFINITIES (COLEOPTERA: Tj ETQq0.00rgBT /Overlock 18 | 0.1 | |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 271 | A second species of <i>Oculogryphus</i> (Coleoptera, Lampyridae), with notes on the phylogenetic affinities of the genus. <i>ZooKeys</i> , 2011, 97, 31-38. | 0.5 | 10 |
| 272 | Andinopanurgus, a new Andean subgenus of <i>Protandrena</i> (Hymenoptera, Andrenidae). <i>ZooKeys</i> , 2011, 126, 57-76. | 0.5 | 10 |
| 273 | Serphitid wasps in Cretaceous amber from New Jersey (Hymenoptera: Serphitidae). <i>Insect Systematics and Evolution</i> , 2011, 42, 197-204. | 0.2 | 10 |
| 274 | A new species of <i>Geotrigona</i> Moure from the Caribbean coast of Colombia (Hymenoptera, Apidae). <i>ZooKeys</i> , 2012, 172, 77-87. | 0.5 | 10 |
| 275 | A new wild, pollinating bee species of the genus <i>Tetraloniella</i> from the Arabian Peninsula (Hymenoptera, Apidae). <i>ZooKeys</i> , 2012, 172, 89-96. | 0.5 | 10 |
| 276 | A new genus of Ptilodactylidae (Coleoptera: Byrrhoidea) in mid-Cretaceous amber from Myanmar (Burma). <i>Geodiversitas</i> , 2012, 34, 569-574. | 0.2 | 10 |
| 277 | Treatise on the Isoptera of the World. <i>Bulletin of the American Museum of Natural History</i> , 2013, 377, 2433-2705. | 1.2 | 10 |
| 278 | The first diplatyid earwig in Tertiary amber (Dermaptera: Diplatyidae): A new species from Miocene Mexican amber. <i>Insect Systematics and Evolution</i> , 2013, 44, 157-166. | 0.2 | 10 |
| 279 | Garrouste et al. reply. <i>Nature</i> , 2013, 494, E4-E5. | 13.7 | 10 |
| 280 | Recent Findings of <i>Olixon banksii</i> in the Nearctic with Notes on Its Biology (Hymenoptera: Encyrtidae). <i>Tropical Entomologist</i> , 2013, 40, 382-387. | 0.1 | 10 |
| 281 | A new crown wasp in Cretaceous amber from Myanmar (Hymenoptera: Stephanidae). <i>Cretaceous Research</i> , 2017, 69, 56-61. | 0.6 | 10 |
| 282 | Further evidence of Cretaceous termitophily: Description of new termite hosts of the trichopseniine <i>Cretotrichopsenius</i> (Coleoptera: Staphylinidae), with emendations to the classification of lower termites (Isoptera). <i>Palaeoentomology</i> , 2021, 4, 1-10. | 0.4 | 10 |
| 283 | A Robinson Crusoe story in the fossil record: Plant-insect interactions from a Middle Jurassic ephemeral volcanic island (Eastern Spain). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2021, 583, 110655. | 1.0 | 10 |
| 284 | Nest Architecture, Immature Stages, and Ethnoentomology of a New Species of <i>Trigonisca</i> from Northern Colombia (Hymenoptera: Apidae). <i>American Museum Novitates</i> , 2019, 3319, 1. | 0.2 | 10 |
| 285 | A revision of the augochlorine bee genus <i>Chlerogas</i> Vachal (Hymenoptera: Halictidae). <i>Zoological Journal of the Linnean Society</i> , 1999, 125, 463-486. | 1.0 | 9 |
| 286 | The First Fossil of the Subfamily Trypanaeinae (Coleoptera: Histeridae): A New Species of <i>Trypanaeus</i> in Dominican Amber. <i>The Coleopterists Bulletin</i> , 2006, 60, 333-340. | 0.1 | 9 |
| 287 | An adventitious distal abscissa in the forewing of honey bees (Hymenoptera: Apidae: <i>Apis</i>). <i>Apidologie</i> , 2008, 39, 674-682. | 0.9 | 9 |
| 288 | <I>Laasbium</I> Scudder: A Genus of Tertiary Earwigs, Not Rove Beetles, and the Classification of Florissant Fossil Dermaptera (Insecta). <i>Annales Zoologici</i> , 2010, 60, 101-108. | 0.1 | 9 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 289 | A new species of <i>Microsphecodes</i> from Jamaica (Hymenoptera, Halictidae).. <i>ZooKeys</i> , 2011, 111, 33-40. | 0.5 | 9 |
| 290 | The apid cuckoo bees of the Cape Verde Islands (Hymenoptera, Apidae). <i>ZooKeys</i> , 2012, 218, 77-109. | 0.5 | 9 |
| 291 | A new Cenomanian–Turonian (Late Cretaceous) insect assemblage from southeastern Morocco. <i>Cretaceous Research</i> , 2012, 35, 88-93. | 0.6 | 9 |
| 292 | A new <i>Stelis</i> (<i>Dolichostelis</i>) from northern Colombia (Hymenoptera: Megachilidae): first records for South America and a synopsis of the bee fauna from the Caribbean region of Colombia. <i>Journal of Natural History</i> , 2012, 46, 2919-2934. | 0.2 | 9 |
| 293 | A New Xyelydid Sawfly from the Early Cretaceous of China (Hymenoptera: Xyelydidae). <i>Journal of the Kansas Entomological Society</i> , 2013, 86, 78-83. | 0.1 | 9 |
| 294 | The first male of the extinct bee tribe Melikertini (Hymenoptera: Apidae). <i>Journal of Melittology</i> , 2014, , 1-18. | 0.2 | 9 |
| 295 | New Evanoid Wasps from the Cenomanian of Myanmar (Hymenoptera: Othniellithidae, Aulacidae), with a Summary of Family-Group Names Among Evanioidea. <i>American Museum Novitates</i> , 2017, 3871, 1-28. | 0.2 | 9 |
| 296 | A new genus of protorhyssaline wasps in Raritan amber (Hymenoptera, Braconidae). <i>ZooKeys</i> , 2017, 711, 103-111. | 0.5 | 9 |
| 297 | A new lineage of braconid wasps in Burmese Cenomanian amber (Hymenoptera, Braconidae). <i>ZooKeys</i> , 2018, 730, 75-86. | 0.5 | 9 |
| 298 | Abdominal serial homologues of wings in Paleozoic insects. <i>Current Biology</i> , 2022, 32, 3414-3422.e1. | 1.8 | 9 |
| 299 | Mesozoic insect fossils reveal the early evolution of twig mimicry. <i>Science Bulletin</i> , 2022, 67, 1641-1643. | 4.3 | 9 |
| 300 | A Miocene Halictine Bee from Rubielos de Mora Basin, Spain (Hymenoptera: Halictidae). <i>American Museum Novitates</i> , 2006, 3503, 1-10. | 0.2 | 8 |
| 301 | Oculogryphus, A Remarkable New Genus of Fireflies from Asia (Coleoptera: Lampyridae). <i>American Museum Novitates</i> , 2007, 3600, 1-19. | 0.2 | 8 |
| 302 | A New Species of <i>Dialictus</i> from Sombrero Island, Anguilla (Hymenoptera, Halictidae). <i>ZooKeys</i> , 2011, 86, 61-68. | 0.5 | 8 |
| 303 | <i>Alocanthedon</i> , a new subgenus of <i>Chalicodoma</i> from Southeast Asia (Hymenoptera, Megachilidae). <i>ZooKeys</i> , 2011, 101, 51-80. | 0.5 | 8 |
| 304 | The Oldest Fossil of the Subfamily Osoriinae (Coleoptera: Staphylinidae), from Eocene Cambay Amber (India). <i>The Coleopterists Bulletin</i> , 2013, 67, 304-308. | 0.1 | 8 |
| 305 | Species status and new distribution records for <i>Lithurgus huberi</i> (Hymenoptera, Megachilidae,) Tj ETQql 1 0.784314 rgBT /Oyerlock 10 | 0.8 | 10 |
| 306 | <i> <i>Incasarus garciai</i> </i>, a new genus and species of panurgine bees from the Peruvian Andes (Hymenoptera: Andrenidae). <i>Journal of Melittology</i> , 2013, , 1-9. | 0.2 | 8 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 307 | An orchid bee of the genus <i>Eulaema</i> in Early Miocene Mexican amber (Hymenoptera: Apidae). Novitates Paleoentomologicae, 2014, , 1. | 0.6 | 8 |
| 308 | Allodapine Bees in the Arabian Peninsula (Hymenoptera: Apidae): A New Species of Braunsapis from the Sarawat Mountains, with an Overview of the Arabian Fauna. American Museum Novitates, 2014, 3801, 1-15. | 0.2 | 8 |
| 309 | The wasp family Spathiopterygidae in mid-Cretaceous amber from Myanmar (Hymenoptera: Diaprioidea). Comptes Rendus - Palevol, 2015, 14, 95-100. | 0.1 | 8 |
| 310 | A plesiomorphic gasteruptiid wasp in Cenomanian amber from Myanmar (Hymenoptera: Gasteruptiidae). Cretaceous Research, 2016, 63, 177-182. | 0.6 | 8 |
| 311 | A putative twisted-wing parasitoid planidium (Insecta: Strepsiptera) in Taimyr Upper Cretaceous amber. Cretaceous Research, 2017, 69, 106-112. | 0.6 | 8 |
| 312 | Termite Evolution: A Primal Knock on Wood or a Hearty Mouthful of Dirt. Current Biology, 2019, 29, R1126-R1129. | 1.8 | 8 |
| 313 | Current and future ranges of an elusive North American insect using species distribution models. Journal of Insect Conservation, 2019, 23, 175-186. | 0.8 | 8 |
| 314 | A direct association between amber and dinosaur remains provides paleoecological insights. Scientific Reports, 2019, 9, 17916. | 1.6 | 8 |
| 315 | The last meal of an Eocene pollen-feeding fly. Current Biology, 2021, 31, 2020-2026.e4. | 1.8 | 8 |
| 316 | Stingless bees in Miocene amber of southeastern China (Hymenoptera: Apidae). Journal of Melittology, 2021, , 1-83. | 0.2 | 8 |
| 317 | <p>A holopsenelline wasp in mid-Cretaceous amber from Myanmar (Hymenoptera: Tj ETQq1 1 0.784314 rgBT /Overlock 10 TF</p> | | |
| 318 | A new termite bug in Miocene amber from the Dominican Republic (Hemiptera, Termitaphididae). ZooKeys, 0, 25, 61-68. | 0.5 | 8 |
| 319 | Two New Halictine Bees in Miocene Amber from the Dominican Republic (Hymenoptera, Halictidae). ZooKeys, 0, 29, 1-12. | 0.5 | 8 |
| 320 | Life history and evolution of the enigmatic Cretaceousâ€“Eocene Alienopteridae: A critical review. Earth-Science Reviews, 2022, 225, 103914. | 4.0 | 8 |
| 321 | The first leucospid wasp from the fossil record (Hymenoptera: Leucospidae). Journal of Natural History, 2002, 36, 435-441. | 0.2 | 7 |
| 322 | A New Alderfly in Baltic Amber (Megaloptera: Sialidae). American Museum Novitates, 2006, 3513, 1. | 0.2 | 7 |
| 323 | A new record of Thaumastobombus andreniformis Engel 2001 in Eocene amber (Hymenoptera: Apidae). Annales De La Societe Entomologique De France, 2007, 43, 505-508. | 0.4 | 7 |
| 324 | Anoblepsis, A New, Bizarre Braconid Wasp Genus in Dominican Amber (Hymenoptera: Braconidae). Journal of the Kansas Entomological Society, 2008, 81, 368-372. | 0.1 | 7 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 325 | Antiquity and Evolution of Prosternal Horns in Baridine Weevils (Coleoptera: Curculionidae). <i>Journal of Paleontology</i> , 2010, 84, 918-926. | 0.5 | 7 |
| 326 | A new species of the bee genus <i>Ctenoplectrella</i> in middle Eocene Baltic amber (Hymenoptera) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 702 | 0.5 | 7 |
| 327 | African and Southeast Asian<i>Chalicodoma</i> (Hymenoptera: Megachilidae): New Subgenus, New Species, and Notes on the Composition of<i>Pseudomegachile</i> and<i>Largella</i>. <i>Annales Zoologici</i> , 2012, 62, 599-617. | 0.1 | 7 |
| 328 | Systropha androsthenes in Saudi Arabia (Hymenoptera: Halictidae). <i>Journal of the Kansas Entomological Society</i> , 2012, 85, 62-64. | 0.1 | 7 |
| 329 | A male of the bee genus <i>Agapostemon</i> in Dominican amber (Hymenoptera: Halictidae). <i>Journal of Melittology</i> , 2013, , 1-9. | 0.2 | 7 |
| 330 | The bee genus <i>Ischnomelissa</i> in Peru, with a key to the species (Hymenoptera) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 5 | 0.2 | 7 |
| 331 | Whipspiders (Arachnida: Amblypygi) in amber from the Early Eocene and mid-Cretaceous, including maternal care. <i>Novitates Paleoentomologicae</i> , 2014, , 1. | 0.6 | 7 |
| 332 | The genus <i>Macroteleia</i> Westwood in Middle Miocene amber from Peru (Hymenoptera, Platygastriidae) Tj ETQq0 0 0 rgBT /Overlock 10 Tf | 0.5 | 7 |
| 333 | A New Genus and Two New Extant Species Closely Allied with the Fossil Genus <i>Pauroripidius</i> (Coleoptera: Ripiphoridae). <i>Journal of the Kansas Entomological Society</i> , 2014, 87, 333-344. | 0.1 | 7 |
| 334 | First record of the bee genus <i>Compsomelissa</i> in the Kingdom of Saudi Arabia (Hymenoptera: Apidae). <i>Pan-Pacific Entomologist</i> , 2014, 90, 37-39. | 0.1 | 7 |
| 335 | A new trap-jaw ant species of the genus <i>Odontomachus</i> (Hymenoptera: Formicidae: Ponerinae) from the Early Miocene (Burdigalian) of the Czech Republic. <i>Palaontologische Zeitschrift</i> , 2014, 88, 495-502. | 0.8 | 7 |
| 336 | An apterous scelionid wasp in mid-Cretaceous Burmese amber (Hymenoptera: Scelionidae). <i>Comptes Rendus - Palevol</i> , 2017, 16, 5-11. | 0.1 | 7 |
| 337 | Zorotypus dilaticeps sp. nov., a remarkable zorapteran (Zoraptera) in mid-Cretaceous Burmese amber. <i>Cretaceous Research</i> , 2018, 91, 126-130. | 0.6 | 7 |
| 338 | Mouthpart homologies and life habits of Mesozoic long-proboscid scorpionflies. <i>Science Advances</i> , 2020, 6, eaay1259. | 4.7 | 7 |
| 339 | Mating and aggregative behaviors among basal hexapods in the Early Cretaceous. <i>PLoS ONE</i> , 2018, 13, e0191669. | 1.1 | 7 |
| 340 | A fossil species of the primitive mymarid genus <i>Borneomyrm</i> (Hymenoptera: Mymaridae) in Eocene Baltic amber. <i>Novitates Paleoentomologicae</i> , 2013, , 1. | 0.6 | 7 |
| 341 | New mid-Cretaceous earwigs in amber from Myanmar (Dermaptera). <i>Novitates Paleoentomologicae</i> , 2014, , 1. | 0.6 | 7 |
| 342 | Stingless bees (Hymenoptera: Apidae) in Holocene copal and Defaunation resin from Eastern Africa indicate Recent biodiversity change. <i>Holocene</i> , 2022, 32, 414-432. | 0.9 | 7 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 343 | A New Palearctic Genus of Melectine Bees (Hymenoptera: Apidae). American Museum Novitates, 2003, 3392, 1-22. | 0.2 | 6 |
| 344 | The first Mesozoic stephanid wasp (Hymenoptera: Stephanidae). Journal of Paleontology, 2004, 78, 1192-1197. | 0.5 | 6 |
| 345 | A New Species of Chiasmognathus from Southeastern Pakistan (Hymenoptera: Apidae). Journal of the Kansas Entomological Society, 2007, 80, 169-174. | 0.1 | 6 |
| 346 | The Firefly Genus Vesta in Taiwan (Coleoptera: Lampyridae). Journal of the Kansas Entomological Society, 2007, 80, 265-280. | 0.1 | 6 |
| 347 | An orussid wood wasp in amber from the Dominican Republic (Hymenoptera: Orussidae). Transactions of the Kansas Academy of Science, 2008, 111, 39-44. | 0.0 | 6 |
| 348 | A new genus of sphaeropsocid bark lice from the Early Cretaceous amber of Lebanon (Psocodea:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 50 | 0.4 | 6 |
| 349 | A New Genus of Dustywings Allied to Archiconiocompsa in Baltic Amber (Neuroptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 50 | 0.0 | 6 |
| 350 | The bee genus Chlerogas in Bolivia (Hymenoptera, Halictidae). ZooKeys, 2010, 46, 61-70. | 0.5 | 6 |
| 351 | The melectine bee genera Brachymelecta and Sinomelecta (Hymenoptera, Apidae). ZooKeys, 2012, 244, 1-19. | 0.5 | 6 |
| 352 | A New Species of Roach from the Jurassic of India (Blattaria: Mesoblattinidae). Journal of the Kansas Entomological Society, 2012, 85, 1-4. | 0.1 | 6 |
| 353 | Description of the male of Megalava truncata Perrichot (Hymenoptera: Megalyridae) in Early Cretaceous amber from El Soplao (Spain). Zootaxa, 2012, 3274, 29. | 0.2 | 6 |
| 354 | First record of Chiasmognathus from the Kingdom of Saudi Arabia (Hymenoptera, Apidae). Journal of Hymenoptera Research, 2013, 35, 83-89. | 0.8 | 6 |
| 355 | The first Mesozoic Leptopodidae (Hemiptera: Heteroptera: Leptopodomorpha), from Canadian Late Cretaceous amber. Historical Biology, 2014, 26, 702-709. | 0.7 | 6 |
| 356 | Two new genera of Cretaceous dustywings in amber from northern Myanmar (Neuroptera:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 222 Td | 0.6 | 6 |
| 357 | The first twisted-wing parasitoid in Eocene amber from north-eastern China (Strepsiptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 50 | 0.2 | 6 |
| 358 | A new genus of alderflies (Megaloptera: Sialidae) in Upper Cretaceous Burmese amber. Cretaceous Research, 2016, 64, 7-11. | 0.6 | 6 |
| 359 | Response to "Evidence from amber for the origins of termitophily". Current Biology, 2017, 27, R794-R795. | 1.8 | 6 |
| 360 | Biology of the mason bee Osmia latreillei (Hymenoptera: Megachilidae) under artificial nesting conditions in Egypt. Journal of Asia-Pacific Entomology, 2018, 21, 754-759. | 0.4 | 6 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 361 | A new and diverse paleofauna of the extinct snakefly family Baissopteridae from the mid-Cretaceous of Myanmar (Raphidioptera). <i>Organisms Diversity and Evolution</i> , 2020, 20, 565-595. | 0.7 | 6 |
| 362 | Fossil Social Insects. , 2021, , 384-403. | | 6 |
| 363 | Phylogeny of Chrysopidae (Neuroptera), with emphasis on morphological trait evolution. <i>Zoological Journal of the Linnean Society</i> , 2022, 194, 1374-1395. | 1.0 | 6 |
| 364 | The complete life cycle of a Cretaceous beetle parasitoid. <i>Current Biology</i> , 2021, 31, R118-R119. | 1.8 | 6 |
| 365 | The <i>Sphecodes</i> of Cuba (Hymenoptera: Halictidae). <i>Acta Zoologica Cracoviensia - Series B: Invertebrata</i> , 2006, 49, 73-78. | 0.2 | 6 |
| 366 | Revision of the green lacewing subgenus Ankylopteryx (Sencera) (Neuroptera, Chrysopidae). <i>ZooKeys</i> , 2015, 543, 111-127. | 0.5 | 6 |
| 367 | A new species of the carpenter bee genus Xylocopa from the Sarawat Mountains in southwestern Saudi Arabia (Hymenoptera, Apidae). <i>ZooKeys</i> , 2017, 716, 29-41. | 0.5 | 6 |
| 368 | Bees. , 2020, , 1-17. | | 6 |
| 369 | Current and future distributions of a native Andean bumble bee. <i>Journal of Insect Conservation</i> , 2022, 26, 559-569. | 0.8 | 6 |
| 370 | A new Eocene-Oligocene snakefly from Florissant, Colorado (Raphidioptera: Raphidiidae). <i>Transactions of the Kansas Academy of Science</i> , 2003, 106, 124-128. | 0.0 | 5 |
| 371 | Notes on a megachiline bee (Hymenoptera: Megachilidae) from the Miocene of Idaho. <i>Transactions of the Kansas Academy of Science</i> , 2004, 107, 97-100. | 0.0 | 5 |
| 372 | SPHAEROPSOCUS KUENOWII HAGEN IN ROVNO AMBER FROM THE UKRAINE (PSOCOPTERA: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 302 | | |
| 373 | TWO OVERLOOKED FAMILY-GROUP NAMES FOR FOSSIL TERMITES (ISOPTERA: MASTOTERMITIDAE). <i>Entomological News</i> , 2007, 118, 105-106. | 0.1 | 5 |
| 374 | Case 3461ancylini Michener, 1944 (Insecta, Hymenoptera): proposed emendation of spelling to aencylaini, to remove homonymy with aencylini Rafinesque, 1815 (Mollusca, Gastropoda). <i>Bulletin of Zoological Nomenclature</i> , 2008, 65, 198-201. | 0.2 | 5 |
| 375 | Two New Species of <i>Ammobates</i> from the Arabian Peninsula and Egypt (Hymenoptera: Apidae). <i>Transactions of the Kansas Academy of Science</i> , 2009, 112, 191-197. | 0.0 | 5 |
| 376 | Baltic amber Ibaliiidae (Hymenoptera: Cynipoidea): a new genus with implications for the phylogeny and historical biogeography of the family. <i>Systematic Entomology</i> , 2010, 35, 164-171. | 1.7 | 5 |
| 377 | First fossil Calopterygoidea (Odonata: Zygoptera) from Southeastern Asia: A new genus and species from the Paleogene of China. <i>Geobios</i> , 2010, 43, 349-353. | 0.7 | 5 |
| 378 | The bee genus Caenaugochlora (Hymenoptera, Apoidea) and its constituent subgenera, with new species of Caenaugochlora s.str. from Ecuador. <i>ZooKeys</i> , 0, 37, 69-80. | 0.5 | 5 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 379 | A pupal caddisfly from the Early Cretaceous of China (Trichoptera). <i>Cretaceous Research</i> , 2010, 31, 396-399. | 0.6 | 5 |
| 380 | A New Genus and Species of Aetheogrammatidae from the Jurassic of Inner Mongolia, China (Neuroptera). <i>Journal of the Kansas Entomological Society</i> , 2011, 84, 315-319. | 0.1 | 5 |
| 381 | A new Chlerogelloides from northeastern Brazil and French Guiana, with a key to the species (Hymenoptera, Halictidae). <i>ZooKeys</i> , 2012, 185, 41-53. | 0.5 | 5 |
| 382 | Sambia succinica, a crown group tenthredinid from Eocene Baltic amber (Hymenoptera:) Tj ETQq0 0 0 rgBT /Overlock_10 Tf 50_622 Td (T | 0.2 | |
| 383 | A Checklist of the Orchid Bees of Nicaragua (Hymenoptera: Apidae: Euglossini). <i>Journal of the Kansas Entomological Society</i> , 2012, 85, 135-144. | 0.1 | 5 |
| 384 | New Species of Macrocephalic Halictine Bees (Hymenoptera: Halictidae). <i>Annales Zoologici</i> , 2012, 62, 297-307. | 0.1 | 5 |
| 385 | Revision of the orchid bee subgenus <i>Euglossella</i> (Hymenoptera: Apidae), part II: The <i>viridis</i> and <i>mandibularis</i> species groups. <i>Journal of Melittology</i> , 2014, , 1-108. | 0.2 | 5 |
| 386 | Serendipity at the Smithsonian: The 107-year journey of <i>Rhipidocyrtus muiri</i> Falin & Engel, new genus and species (Ripidiinae, Ripidiini), from jungle beast to valid taxon. <i>ZooKeys</i> , 2014, 424, 101-116. | 0.5 | 5 |
| 387 | A new species of the cleptoparasitic bee genus <i>Thyreus</i> from northern Yemen and southwestern Saudi Arabia (Hymenoptera, Apidae). <i>ZooKeys</i> , 2014, 428, 29-40. | 0.5 | 5 |
| 388 | New Records of Nomiine and Halictine Bees in the Kingdom of Saudi Arabia (Hymenoptera: Halictidae). <i>Journal of the Kansas Entomological Society</i> , 2014, 87, 312-317. | 0.1 | 5 |
| 389 | A new family of primitive serphitoid wasps in Lebanese amber (Hymenoptera: Serphitoidea). <i>Novitates Paleoentomologiae</i> , 2015, , 1. | 0.6 | 5 |
| 390 | The fourth Mesozoic water measurer discovered in mid-Cretaceous Burmese amber (Heteroptera:) Tj ETQq0 0 0 rgBT /Overlock_10 Tf 50_60 | 0.6 | |
| 391 | A new xyelotomid (Hymenoptera) from the Middle Jurassic of China displaying enigmatic venational asymmetry. <i>BMC Evolutionary Biology</i> , 2016, 16, 155. | 3.2 | 5 |
| 392 | First record of the bee genus <i>Melitta</i> from the Arabian Peninsula (Hymenoptera: Apoidea: Melittidae). <i>Zoology in the Middle East</i> , 2016, 62, 352-357. | 0.2 | 5 |
| 393 | Ecological niche modeling of the rare bee <i>Promelitta alboclypeata</i> reveals possible cryptic differentiation across northern Africa and Arabia (Hymenoptera: Melittidae). <i>Apidologie</i> , 2016, 47, 509-514. | 0.9 | 5 |
| 394 | Distributional modeling of Mantophasmatodea (Insecta: Notoptera): a preliminary application and the need for future sampling. <i>Organisms Diversity and Evolution</i> , 2016, 16, 259-268. | 0.7 | 5 |
| 395 | Marsupial brood care in Cretaceous tanaidaceans. <i>Scientific Reports</i> , 2017, 7, 4390. | 1.6 | 5 |
| 396 | The first gynandromorph of a zorapteran and potential thelytokous parthenogenesis in a population of <i>Zorotypus brasiliensis</i> Silvestri (Zoraptera: Zorotypidae). <i>Revista Brasileira De Entomologia</i> , 2017, 61, 318-322. | 0.1 | 5 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 397 | A primitive honey bee from the Middle Miocene deposits of southeastern Yunnan, China (Hymenoptera,) Tj ETQq1 1.0.784314 rgBT /Ov | 0.5 | 5 |
| 398 | Notes on Papuan and Malesian stingless bees, with the descriptions of new taxa (Hymenoptera:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 | 0.2 | 5 |
| 399 | New eodermapteran earwigs (Dermaptera) from the Middle Jurassic Jiulongshan Formation of China. Alcheringa, 2021, 45, 335-343. | 0.5 | 5 |
| 400 | A new species of the extinct family Proceropidae (Hemiptera: Cercopoidea) from the Jurassic of northeastern China. Palaeoentomology, 2018, 1, 51. | 0.4 | 5 |
| 401 | Rediscovered parasitism of <i>Andrena savignyi</i> Spinola (Hymenoptera, Andrenidae) by <i>Stylops</i> (Strepsiptera, Stylopidae) and revised taxonomic status of the parasite. ZooKeys, 2015, 519, 117-139. | 0.5 | 5 |
| 402 | First record of a tear-drinking stingless bee <i>Lisotrigona cacciae</i> (Nurse) (Hymenoptera: Apidae:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 54 Lanka, 2017, 45, 79. | 0.1 | 5 |
| 403 | Augochlorini Beebe, 1925 (Insecta, Hymenoptera): corrected authorship and date (not Moure, 1943). Bulletin of Zoological Nomenclature, 1999, 56, 198-198. | 0.2 | 5 |
| 404 | A new species of spongiphorine earwig in Miocene amber from the Dominican Republic (Dermaptera: Spongiphoridae). Palaeoentomology, 2019, 2, 560-565. | 0.4 | 5 |
| 405 | Conservation Implications of a Newly Discovered Bee Species on Isla Robinson Crusoe, Chile. Conservation Biology, 2001, 15, 803-805. | 2.4 | 4 |
| 406 | The Wasp Genus <i>Clystopsenella</i> in Belize (Hymenoptera: Scolebythidae). Journal of the Kansas Entomological Society, 2005, 78, 186-188. | 0.1 | 4 |
| 407 | The Zorapteran <i>Zorotypus huxleyi</i> in Guyana (Zoraptera: Zorotypidae). Journal of the Kansas Entomological Society, 2008, 81, 394-395. | 0.1 | 4 |
| 408 | Noctoraptor in Bolivia (Hymenoptera: Halictidae). Journal of the Kansas Entomological Society, 2011, 84, 64-70. | 0.1 | 4 |
| 409 | The identity of the Neotropical stingless bee <i>Friesomelitta meadewaldoi</i> (Cockerell, 1915) (Hymenoptera, Apidae). ZooKeys, 2011, 111, 19-31. | 0.5 | 4 |
| 410 | A New Snakefly from the Eocene Green River Formation (Raphidioptera: Raphidiidae). Transactions of the Kansas Academy of Science, 2011, 114, 77-87. | 0.0 | 4 |
| 411 | First Mesozoic Microphysidae (Hemiptera): a new genus and species in Late Cretaceous amber from Canada. Canadian Entomologist, 2011, 143, 349-357. | 0.4 | 4 |
| 412 | Wasp mimicry among Palaeocene reduviid bug from Svalbard. Acta Palaeontologica Polonica, 0, , . | 0.4 | 4 |
| 413 | COMMENT ON MARDEN (2013): âœREANALYSIS AND EXPERIMENTAL EVIDENCE INDICATE THAT THE EARLIEST TRACE FOSSIL OF A WINGED INSECT WAS A SURFACE SKIMMING NEOPTERANâœ Evolution; International Journal of Organic Evolution, 2013, 67, 2142-2149. | 1.1 | 4 |
| 414 | A diminutive pelecinid wasp from the Eocene Kishenehn Formation of northwestern Montana (Hymenoptera: Pelecinidae). Novitates Paleontologicae, 2014, , 1. | 0.6 | 4 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 415 | The bee genus <i>Caenaugeochlora</i> in Venezuela (Hymenoptera: Halictidae). Journal of Melittology, 2014, , 1-10. | 0.2 | 4 |
| 416 | A new paracolletine bee from Colombia (Hymenoptera: Colletidae), with an updated checklist of the tropical Andean bee fauna. Journal of Melittology, 2014, , 1-26. | 0.2 | 4 |
| 417 | A Pentocellar Female of <i>Caenaugeochlora inermis</i> from Southern Mexico (Hymenoptera: Halictidae). Journal of the Kansas Entomological Society, 2014, 87, 392-394. | 0.1 | 4 |
| 418 | Notes on Cretaceous amber Braconidae (Hymenoptera), with descriptions of two new genera. Novitates Paleoentomologicae, 2016, , 1. | 0.6 | 4 |
| 419 | Phylogenetic Relationships of a New Genus of Calliopsis Bees from Peru, with a Review of <i>Spinoliella</i> Ashmead (Hymenoptera: Andrenidae). Bulletin of the American Museum of Natural History, 2017, 412, 1-71. | 1.2 | 4 |
| 420 | A new genus of labidurid earwigs in mid-Cretaceous amber from northern Myanmar (Dermaptera) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 | 0.6 | 4 |
| 421 | Early Cretaceous termites in amber from northern Spain (Isoptera). Cretaceous Research, 2020, 110, 104385. | 0.6 | 4 |
| 422 | Announcing Big-Bee: An initiative to promote understanding of bees through image and trait digitization. Biodiversity Information Science and Standards, 0, 5, . | 0.0 | 4 |
| 423 | <p>A new dustywing (Neuroptera: Coniopterygidae) from the Early Cretaceous amber of Spain</p>. Palaeoentomology, 2019, 2, 279-288. | 0.4 | 4 |
| 424 | A new species of Astreptolabis in mid-Cretaceous amber from northern Myanmar, with the discovery of the first male of Astreptolabidinae (Dermaptera). ZooKeys, 2020, 911, 101-112. | 0.5 | 4 |
| 425 | A new species of the bee genus Chlerogella from Panama (Hymenoptera: Halictidae). Zootaxa, 2003, 286, . | 0.2 | 4 |
| 426 | A new species of Chiasmognathus from Iran, with a note on Chiasmognathus aegyptiacus in Israel (Hymenoptera: Apidae).. Contributions To Entomology, 2008, 58, 223-226. | 0.1 | 4 |
| 427 | A new genus of anaxyelid wood wasps from the mid-Cretaceous and the phylogeny of Anaxyelidae (Hymenoptera). Journal of Hymenoptera Research, 0, 86, 151-169. | 0.8 | 4 |
| 428 | Alitrepaninae, a new subfamily of auger beetles from mid-Cretaceous Kachin amber of northern Myanmar (Coleoptera: Bostrichidae). Cretaceous Research, 2022, 137, 105244. | 0.6 | 4 |
| 429 | Morphological Phylogeny of New Cretaceous Fossils Elucidates the Early History of Soil Dwelling Among Bugs. Frontiers in Ecology and Evolution, 0, 10, . | 1.1 | 4 |
| 430 | The Bee Genus <i>Andinaugochlora</i> in Central America (Hymenoptera: Halictidae). Journal of the Kansas Entomological Society, 2004, 77, 116-120. | 0.1 | 3 |
| 431 | ON THE AVAILABILITY OF FAMILY-GROUP NAMES BASED ON SCRAPTER (HYMENOPTERA: COLLETIDAE). Entomological News, 2006, 117, 117-119. | 0.1 | 3 |
| 432 | Phylogeny of the cleptoparasitic bee genus <i>Exaerete</i> (Hymenoptera: Apidae). Apidologie, 2007, 38, 419-425. | 0.9 | 3 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 433 | A Miocene Snakefly from Stewart Valley, Nevada (Raphidioptera: Raphidiidae). Transactions of the Kansas Academy of Science, 2009, 112, 211-214. | 0.0 | 3 |
| 434 | A new micropterous species of Embolemus Westwood from Baltic amber (Hymenoptera: Embolemidae). Annales De Paleontologie, 2011, 97, 1-7. | 0.1 | 3 |
| 435 | Discovery of the female of Pyrocoelia prolongata in Taiwan (Coleoptera, Lampyridae). ZooKeys, 2011, 116, 49-57. | 0.5 | 3 |
| 436 | Bolivian Neocorynura (Hymenoptera: Halictidae): A new species and preliminary key to the fauna. Tijdschrift Voor Entomologie, 2012, 155, 3-8. | 0.1 | 3 |
| 437 | A New Fossil Crane Fly from the Early Pannonian of the Styrian Basin (Diptera: Tipulidae). Journal of the Kansas Entomological Society, 2012, 85, 160-163. | 0.1 | 3 |
| 438 | <i>Zoropecelinus zigrasi</i>, a pelecinid wasp in mid-Cretaceous amber from Myanmar (Hymenoptera: Pelecinidae). Novitates Paleoentomologicae, 2013, , 1. | 0.6 | 3 |
| 439 | A new species of <i>Chiasmognathus</i> from Kazakhstan (Hymenoptera: Apidae). Journal of Melittology, 2013, , 1-7. | 0.2 | 3 |
| 440 | A new fossil drywood termite species from the Late Eocene of France allied to <i>Cryptotermes</i> and <i>Procryptotermes</i> (Isoptera: Kalotermitidae). Novitates Paleoentomologicae, 2015, , 1. | 0.6 | 3 |
| 441 | Pangusyndicus gen. nov.: a new mid-Cretaceous scydmaenine with reduced antennae and prothoracic gland (Coleoptera, Staphylinidae: Scydmaeninae). Journal of Systematic Palaeontology, 2019, 17, 1129-1141. | 0.6 | 3 |
| 442 | Leaf-mimicking katydids from the Middle Miocene of Yunnan, southwestern China (Orthoptera: Tettigidae). Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 30 | 0.8 | 3 |
| 443 | Mimicry in Cretaceous Bugs. IScience, 2020, 23, 101280. | 1.9 | 3 |
| 444 | New species of webspinners (Insecta: Embioidea) from mid-Cretaceous amber of northern Myanmar. Cretaceous Research, 2020, 113, 104457. | 0.6 | 3 |
| 445 | Beaded lacewings (Neuroptera: Berothidae) in amber from the Lower Cretaceous of Spain. Cretaceous Research, 2021, 119, 104705. | 0.6 | 3 |
| 446 | On the spelling of family-group names based on the genus Trigonalyss Westwood (Hymenoptera: Encyrtidae). Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 20 | 0.1 | 3 |
| 447 | Description of a Cretaceous amber fossil putatively of the tribe Coprophilini (Coleoptera, Tenebrionidae). Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 20 | 0.5 | 3 |
| 448 | A new genus of Pelecotominae from Mexico, with notes on the genera Clinops and Scotoscopus and the description of new species (Coleoptera, Ripiphoridae). ZooKeys, 2019, 857, 59-84. | 0.5 | 3 |
| 449 | A quadriocellar scoliid wasp (Hymenoptera, Scoliidae) from Mallorca, with a brief account of supernumerary ocelli in insects. Zoosystematics and Evolution, 2015, 91, 191-197. | 0.4 | 3 |
| 450 | An anocellar polistine wasp (Hymenoptera, Vespidae, Polistinae) from Texas. Zoosystematics and Evolution, 2016, 92, 251-255. | 0.4 | 3 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 451 | New archidermpteran earwigs (Dermaptera) from the Middle Jurassic of Inner Mongolia, China. ZooKeys, 2021, 1065, 125-139. | 0.5 | 3 |
| 452 | A new genus and species of Baissidae in Late Cretaceous amber from New Jersey (Hymenoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 7 | 0.6 | 3 |
| 453 | New genera of melitturguline bees from Saudi Arabia and Persia, with notes on related genera and a key to the Arabian fauna (Hymenoptera: Andrenidae). Journal of Hymenoptera Research, 0, 69, 1-21. | 0.8 | 3 |
| 454 | How to extract and analyze pollen from internal organs and exoskeletons of fossil insects. STAR Protocols, 2021, 2, 100923. | 0.5 | 3 |
| 455 | Notes on South American stingless bees of the genus <i>Scaptotrigona</i> (Hymenoptera: Apidae), Part I: short-bristle species, the <i>tubiba</i> species group. Entomologist's Monthly Magazine, 2022, 158, 41-59. | 0.1 | 3 |
| 456 | Notes on South American stingless bees of the genus <i>Scaptotrigona</i> (Hymenoptera: Apidae), Part II: Subgroup A of the <i>postica</i> species group. Journal of Melittology, 2022, , 1-51. | 0.2 | 3 |
| 457 | Notes on South American stingless bees of the genus <i>Scaptotrigona</i> (Hymenoptera: Apidae), Part III: A revised infrageneric classification and new species. Journal of Melittology, 2022, , 1-29. | 0.2 | 3 |
| 458 | A Cretaceous balloon lifts the veil on the antiquity and evolution of nuptial gifts. Gondwana Research, 2022, 107, 146-153. | 3.0 | 3 |
| 459 | Mastigocoleidae fam. nov., a New Mesozoic Beetle Family and the Early Evolution of Dryopoidea (Coleoptera). Insect Systematics and Diversity, 2022, 6, . | 0.7 | 3 |
| 460 | Notes on South American stingless bees of the genus <i>Scaptotrigona</i> (Hymenoptera: Apidae), Part IV: Four new species of group B from the Andean region. Journal of Melittology, 2022, , . | 0.2 | 3 |
| 461 | Zoraptera. , 2009, , 1069-1070. | | 2 |
| 462 | A new species of Chilicola from Bahia, Brazil (Hymenoptera, Colletidae), with a key to the species of the megalostigma group. ZooKeys, 2011, 153, 81-90. | 0.5 | 2 |
| 463 | A new species of <i>Microsphecodes</i> from Peru, with notes on the classification of the genus (Hymenoptera: Halictidae). Journal of Melittology, 2013, , 1-9. | 0.2 | 2 |
| 464 | A ceraphronid wasp in Early Miocene amber from the Dominican Republic (Hymenoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 222 Td 0 | 0.6 | 2 |
| 465 | Three new species of the bee genus <i>Ruizantheda</i> sensu lato (Hymenoptera: Halictidae: Caenohalictina). Zootaxa, 2014, 3889, 58-70. | 0.2 | 2 |
| 466 | A new stingless bee species of the genus <i>Nogueirapis</i> from Costa Rica (Hymenoptera: Tj ETQq0 0 0 rgBT /Overlock 10 | 0.2 | 2 |
| 467 | Description of <i>Oculogryphus shuensis</i> sp. n. (Coleoptera, Lampyridae), the first species of the genus in the Sino-Japanese realm, with a modified key to the subfamily Ototretinae. ZooKeys, 2014, 378, 41-47. | 0.5 | 2 |
| 468 | Additional Records of <i>Zorotypus barberi</i> from Puerto Rico (Zoraptera: Zorotypidae). Journal of the Kansas Entomological Society, 2014, 87, 389-391. | 0.1 | 2 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 469 | A review of the genera and subgenera of Oxaeinae (Hymenoptera: Andrenidae). Journal of Melittology, 2015, , 1-18. | 0.2 | 2 |
| 470 | A giant termite of the genus <i>Gyatermes</i> from the late Miocene of Nagano Prefecture, Japan (Isoptera). Novitates Paleoentomologicae, 2015, , 1. | 0.6 | 2 |
| 471 | A new genus and species of maimetshid wasps in Lebanese Early Cretaceous amber (Hymenoptera:) Tj ETQq1 1 0.784314 rgBT /Overlock | 0.6 | 2 |
| 472 | Two New Species of <i>Pterydrias</i> Reitter (Coleoptera: Ripiphoridae), Significantly Expanding the Biogeographic Range of the Genus. The Coleopterists Bulletin, 2016, 70, 203-213. | 0.1 | 2 |
| 473 | Charles D. Michener (1918â€“2015): The Compleat Melittologist ¹ . Journal of the Kansas Entomological Society, 2016, 89, 1-44. | 0.1 | 2 |
| 474 | A new species of <i>Mermiglossa</i> from Kenya, with comments on the arrangement of Old World Panurginae (Hymenoptera: Andrenidae). Journal of Melittology, 2017, , 1-11. | 0.2 | 2 |
| 475 | A new species of the cleptoparasitic orchid bee genus <i>Exaerete</i> from northern Venezuela (Hymenoptera: Apidae). Entomologist's Monthly Magazine, 2018, 154, 161-175. | 0.1 | 2 |
| 476 | South American Leaf-Cutter Bees (Genus Megachile) of the Subgenera Rhyssomegachile and Zonomegachile, with Two New Subgenera (Hymenoptera: Megachilidae). Bulletin of the American Museum of Natural History, 2018, 2018, 1. | 1.2 | 2 |
| 477 | The first predatory dance fly of the subfamily Ocydromiinae with specialized, raptorial legs in midâ€“Cretaceous amber from Myanmar (Diptera: Hybotidae). Cretaceous Research, 2021, 119, 104697. | 0.6 | 2 |
| 478 | New genera of melikertine bees with facial modifications in Baltic amber (Hymenoptera: Apidae). Journal of Melittology, 2021, , 1-52. | 0.2 | 2 |
| 479 | A key to the species of <i>Nanoplebeia</i>, with descriptions of four new species (Hymenoptera:) Tj ETQq1 1 0.784314 rgBT /Overlock | 0.2 | 2 |
| 480 | Corbiculate Bees. , 2021, , 302-310. | | 2 |
| 481 | Synchrotron-radiation computed tomography uncovers ecosystem functions of fly larvae in an Eocene forest. Palaeontologia Electronica, 0, , . | 0.9 | 2 |
| 482 | <p>A new crown wasp in mid-Cretaceous amber from northern Myanmar (Hymenoptera: Stephanidae)</p>. Palaeoentomology, 2019, 2, 229-235. | 0.4 | 2 |
| 483 | The wasp genus <i>Clystopsenella</i> in Early Miocene amber from the Dominican Republic (Hymenoptera: Scolebythidae). Novitates Paleoentomologicae, 2015, , 1. | 0.6 | 2 |
| 484 | The Cleptoparasitic Bee Genus Chiasmognathus (Hymenoptera: Apidae) in Kenya, with the Description of Two New Species. Journal of East African Natural History, 2019, 108, 17. | 0.6 | 2 |
| 485 | A New Species of Tetralonia (Thygatina) from India, with Notes on the Oriental Fauna (Hymenoptera:) Tj ETQq1 1 0.784314 rgBT /Overlock | 0.2 | 2 |
| 486 | A termite from the Late Oligocene of northern Ethiopia. Acta Palaeontologica Polonica, 0, , . | 0.4 | 2 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 487 | A replacement family-group name among fossil Neuroptera (Insecta). <i>Novitates Paleoentomologicae</i> , 2017, , 1. | 0.6 | 2 |
| 488 | Unique Metasomal Musculature in Sweat Bees (Hymenoptera: Apoidea: Halictidae) Revealed by Micro-CT Scanning. <i>American Museum Novitates</i> , 2019, 2019, 1. | 0.2 | 2 |
| 489 | A key to the subgenera of the orchid bee genus <i>Euglossa</i> (Hymenoptera: Apidae). <i>Entomologist's Monthly Magazine</i> , 2021, 157, 225-241. | 0.1 | 2 |
| 490 | A key to the subgenera of the stingless bee genus <i>Melipona</i> (Hymenoptera: Apidae). <i>Entomologist's Monthly Magazine</i> , 2021, 157, 273-281. | 0.1 | 2 |
| 491 | The first xiphydriid wood wasp in Cretaceous amber (Hymenoptera: Xiphydriidae) and a potential association with Cycadales. <i>Fossil Record</i> , 2022, 24, 445-453. | 0.5 | 2 |
| 492 | A new bee genus from the pampas of eastern Argentina, with appended notes on the classification of <i>Æparacolletines</i> (Hymenoptera: Colletidae). <i>Journal of Melittology</i> , 2022, , 1-39. | 0.2 | 2 |
| 493 | »The first adult mantis lacewing from Baltic amber, with an evaluation of the post-Cretaceous loss of morphological diversity of raptorial appendages in Mantispidae. <i>Fossil Record</i> , 2022, 25, 11-24. | 0.5 | 2 |
| 494 | »A new genus of minute stingless bees from Southeast Asia (Hymenoptera, Apidae). <i>ZooKeys</i> , 2022, 1089, 53-72. | 0.5 | 2 |
| 495 | Declining morphological diversity in snakefly larvae during last 100 million years. <i>Palaontologische Zeitschrift</i> , 2022, 96, 749-780. | 0.8 | 2 |
| 496 | Termite Valkyries: Soldier-Like Alate Termites From the Cretaceous and Task Specialization in the Early Evolution of Isoptera. <i>Frontiers in Ecology and Evolution</i> , 2022, 10, . | 1.1 | 2 |
| 497 | Mayflies as resource pulses in Jurassic lacustrine ecosystems. <i>Geology</i> , 2022, 50, 1043-1047. | 2.0 | 2 |
| 498 | Cretaceous lophocoronids with short proboscis and retractable female genitalia provide the earliest evidence for their feeding and oviposition habits. <i>Cladistics</i> , 2022, 38, 684-701. | 1.5 | 2 |
| 499 | Three Replacement Names in the Bee Genus <i>Andrena</i> (Hymenoptera: Andrenidae). <i>Journal of the Kansas Entomological Society</i> , 2005, 78, 179-180. | 0.1 | 1 |
| 500 | A Note on the Taxonomy of Some Fossil Bees from Germany (Hymenoptera: Apidae). <i>Journal of the Kansas Entomological Society</i> , 2005, 78, 82-83. | 0.1 | 1 |
| 501 | A <i>new Amegilla</i> of the <i>zonata</i> group from Malaysia and Thailand (Hymenoptera: Apidae). <i>Transactions of the Kansas Academy of Science</i> , 2007, 110, 16-22. | 0.0 | 1 |
| 502 | Case 3484 nomiidae Gozis, 1875 (Insecta, Coleoptera): proposed emendation of spelling to nomiusidae to remove homonymy with nomiinae Robertson, 1904 (Insecta, Hymenoptera). <i>Bulletin of Zoological Nomenclature</i> , 2009, 66, 30-33. | 0.2 | 1 |
| 503 | Scientific contributions of Alexandre P. Rasnitsyn, 1959 to present. <i>ZooKeys</i> , 2011, 130, 11-40. | 0.5 | 1 |
| 504 | Charles D. Michener - Recipient of 2010 Hamilton Award. <i>Journal of the Kansas Entomological Society</i> , 2011, 84, 82-82. | 0.1 | 1 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 505 | The first record of Protogryllinae from the Jurassic of India (Orthoptera: Protogryllidae). Journal of the Kansas Entomological Society, 2012, 85, 53-58. | 0.1 | 1 |
| 506 | Biographical Notes for Dr. Jules Desneux (1885–1962): A Multitalented Man of Science. Journal of the Kansas Entomological Society, 2012, 85, 265-267. | 0.1 | 1 |
| 507 | Insect Evolution in an Amberiferous and Stone Alphabet. Insect Systematics and Evolution, 2013, 44, 111-115. | 0.2 | 1 |
| 508 | An evolutionary history embedded in amber: reflection of the Mesozoic shift in weevil-dominated (Coleoptera: Curculionoidea) faunas. Zoological Journal of the Linnean Society, 2014, , . | 1.0 | 1 |
| 509 | A new species of <i>Ptiloglossa</i> from Mexico, with new records of <i>Ptiloglossa cyaniventris</i> from Panama and Costa Rica (Hymenoptera: Colletidae). Journal of Melittology, 2014, , 1-13. | 0.2 | 1 |
| 510 | The bibliography of and taxa proposed by Charles D. Michener. Journal of Melittology, 2015, , 1-109. | 0.2 | 1 |
| 511 | Charles D. Michener (1918–2015): a life among the bees. Arthropod-Plant Interactions, 2017, 11, 243-247. | 0.5 | 1 |
| 512 | First fossil occurrence of the jewel damselflies (Odonata: Chlorocyphidae): a new species from the Late Miocene of Styria, Austria. Annales De La Societe Entomologique De France, 2017, 53, 280-285. | 0.4 | 1 |
| 513 | A new group of species within the bee genus <i>Ruizantheda</i> , with a revised key to the males of the genus (Hymenoptera: Halictidae: Caenohalictini). Zootaxa, 2018, 4415, 513. | 0.2 | 1 |
| 514 | A new genus of augochlorine bees from northern Venezuela (Hymenoptera: Halictidae). Journal of Melittology, 2019, , 1-15. | 0.2 | 1 |
| 515 | On the classification of North American <i>Chelostoma</i> (Hymenoptera: Megachilidae). Journal of Melittology, 2019, , 1-6. | 0.2 | 1 |
| 516 | First fossil Eriocottidae discovered in Eocene Baltic amber (Insecta: Lepidoptera). Zootaxa, 2020, 4834, zootaxa.4834.2.7. | 0.2 | 1 |
| 517 | Bees. , 2021, , 93-109. | | 1 |
| 518 | A new termitophilous genus and species of the tribe Amarygmini Cistel, 1848 from China (Coleoptera: Tj ETQq0 0 0 rgBT /Overlock 101 | 0.2 | |
| 519 | The bee genus <i>Psaenythia</i> in northern South America (Hymenoptera: Andrenidae). Entomologist's Monthly Magazine, 2021, 157, 153-167. | 0.1 | 1 |
| 520 | Terrestrial Isopods from Spanish Amber (Crustacea: Oniscidea): Insights into the Cretaceous Soil Biota. American Museum Novitates, 2021, 2021, . | 0.2 | 1 |
| 521 | A new genus of Early Jurassic earwigs from England (Dermaptera). Novitates Paleoentomologicae, 2021, , 1-3. | 0.6 | 1 |
| 522 | Comment on ‘Mother snail labors for posterity in bed of mid-Cretaceous amber’ by A. Jochum, T. Yu and T.A. Neubauer, Gondwana Research, Volume 97, Pages 68–72. Gondwana Research, 2022, 101, 21-23. | 3.0 | 1 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 523 | On the type locality of <i>Borgatomelissa niveopilosa</i> (Hymenoptera: Andrenidae). Entomologist's Monthly Magazine, 2019, 155, 209-211. | 0.1 | 1 |
| 524 | Apis proava Menge, 1856 (currently Electrapis proava; Insecta, Hymenoptera): proposed conservation by designation of a neotype. Bulletin of Zoological Nomenclature, 1999, 56, 134-135. | 0.2 | 1 |
| 525 | Replacement names for two small carpenter bees from India (Hymenoptera: Apidae, Ceratina). Entomologist's Monthly Magazine, 2018, 154, 296-298. | 0.1 | 1 |
| 526 | Replacement names for sweat bees of the genus <i>Lasioglossum</i> (Hymenoptera: Halictidae). Entomologist's Monthly Magazine, 2018, 154, 266-268. | 0.1 | 1 |
| 527 | First North American species of the European genus <i>Archaemegaptillus</i> from the Upper Carboniferous Pottsville Formation of northern Alabama (Palaeodictyoptera: Archaemegaptillidae). Novitates Paleoentomologicae, 2018, , 1-7. | 0.6 | 1 |
| 528 | A new species of Ctenaugochlora from northwestern Ecuador (Hymenoptera: Halictidae). Entomologist's Monthly Magazine, 2019, 155, 187-192. | 0.1 | 1 |
| 529 | On the subgeneric placement of <i>Megalopta atra</i> (Hymenoptera: Halictidae). Entomologist's Monthly Magazine, 2020, 156, 19-24. | 0.1 | 1 |
| 530 | A semi-aquatic fern (Marsileaceae) from the mid-Cretaceous amber of northern Myanmar. Cretaceous Research, 2022, 133, 105119. | 0.6 | 1 |
| 531 | A second species of the stingless bee genus <i>Plectoplebeia</i> (Hymenoptera: Apidae). Entomologist's Monthly Magazine, 2022, 158, 79-86. | 0.1 | 1 |
| 532 | A New Subgenus and Species of Priochirus from Mid-Cretaceous Kachin Amber (Coleoptera:) Tj ETQq0 0 0 rgBT /Overlock 10 1 Tf 50 382 T | | |
| 533 | »A braconid wasp (Hymenoptera, Braconidae) from the Lower Cretaceous amber of San Just, eastern Iberian Peninsula. ZooKeys, 0, 1103, 65-78. | 0.5 | 1 |
| 534 | The first Sharephemeridae (Insecta: Ephemeroptera) from the Jurassic Shiti Formation of South China. Historical Biology, 2023, 35, 1124-1128. | 0.7 | 1 |
| 535 | A new species of Scaptotrigona from Belize (Hymenoptera: Apidae). Journal of Melittology, 2022, , 1-8. | 0.2 | 1 |
| 536 | The Bee Genus <i>Thrinchostoma</i> Saussure in the Southern Asian Region (Hymenoptera: Halictidae). Proceedings of the Entomological Society of Washington, 2010, 112, 129-139. | 0.0 | 0 |
| 537 | Kumar Krishna, in appreciation. ZooKeys, 2011, 148, 1-13. | 0.5 | 0 |
| 538 | A New Species of Ibaliid Wasp in Baltic Amber (Hymenoptera: Ibaliiidae). Journal of the Kansas Entomological Society, 2012, 85, 164-168. | 0.1 | 0 |
| 539 | 75 Years Ago in Bee Systematics: Cockerell on "The Evolution of the Bees". Journal of the Kansas Entomological Society, 2012, 85, 170-172. | 0.1 | 0 |
| 540 | Family-Group Names for Snakeflies (Raphidioptera). Journal of the Kansas Entomological Society, 2012, 85, 48-52. | 0.1 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 541 | The Apoidea (Hymenoptera) Fauna of Little Tobago Island, West Indies. Journal of the Kansas Entomological Society, 2014, 87, 242-244. | 0.1 | 0 |
| 542 | A review of the New Caledonian Arpactophilus (Hymenoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 70z Td (Crabro) | 0.2 | 0 |
| 543 | Michael Engel. Current Biology, 2016, 26, R905-R907. | 1.8 | 0 |
| 544 | Replacement names for bees in the tribe Megachilini (Hymenoptera: Megachilidae). Journal of Melittology, 2017, , 1-5. | 0.2 | 0 |
| 545 | A tribute to Victor H. Gonzalez: Twenty years of melittological contributions.. Journal of Melittology, 2017, , 1-18. | 0.2 | 0 |
| 546 | A new genus of anthophorine bees from Brunei (Hymenoptera: Apidae). Journal of Melittology, 2018, , 1-13. | 0.2 | 0 |
| 547 | Michener centenary: Memories. Journal of Melittology, 2018, , 1-11. | 0.2 | 0 |
| 548 | Jumping bristletails (Insecta, Archaeognatha) from the Lower Cretaceous amber of Lebanon. Papers in Palaeontology, 2019, 5, 679-697. | 0.7 | 0 |
| 549 | <p>On the availability of the family-group name Lychnocolacidae (Strepsiptera)</p>. Zootaxa, 2020, 4743, 441-442. | 0.2 | 0 |
| 550 | Three new species of the genus Hexarhopalus Fairmaire, 1891 (Coleoptera, Tenebrionidae: Cnadaloniini) from China. Zootaxa, 2021, 5004, 587-597. | 0.2 | 0 |
| 551 | A new baissopterid snakefly (Raphidioptera: Baissopteridae) from mid-Cretaceous amber of northern Myanmar. Cretaceous Research, 2021, , 105028. | 0.6 | 0 |
| 552 | Earliest occurrence of Embiidae: A new genus from earliest Eocene Oise amber (Insecta: Embiodea). Comptes Rendus - Palevol, 2021, , . | 0.1 | 0 |
| 553 | Case 3544ApisarmbrusteriZeuner, 1931 (Insecta, Hymenoptera): proposed conservation by designation of a neotype. Bulletin of Zoological Nomenclature, 2011, 68, 117-121. | 0.2 | 0 |
| 554 | Introducing Novitates Paleoentomologicae: An outlet for occasional fossil insect research at the University of Kansas. Novitates Paleoentomologicae, 2013, , 1. | 0.6 | 0 |
| 555 | On the subgeneric placement of Andrena kraussi Michener (Hymenoptera: Andrenidae). Entomologist's Monthly Magazine, 2019, 155, 38-42. | 0.1 | 0 |
| 556 | First record of the bee genus Meliturgula from Niger (Hymenoptera: Andrenidae). Entomologist's Monthly Magazine, 2019, 155, 168-170. | 0.1 | 0 |
| 557 | New species of the augochlorine bee genus <i>Stilbochlora</i>, with a preliminary key (Hymenoptera: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 70z Td (Crabro) | 0.2 | 0 |
| 558 | Notes on the classification of <i>Ctenocolletes</i> (Hymenoptera: Stenotritidae). Journal of Melittology, 2019, , 1-6. | 0.2 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 559 | A second species of <i>Lonchoprella</i> from northern Argentina (Hymenoptera: Colletidae). Entomologist's Monthly Magazine, 2020, 156, 155-162. | 0.1 | 0 |
| 560 | A new vernal species of Hesperapis from the lower Midwestern United States (Hymenoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 702 | 0.2 | 0 |
| 561 | The bee genus Mesoxaea in Guatemala (Hymenoptera: Andrenidae). Entomologist's Monthly Magazine, 2020, 156, 48-50. | 0.1 | 0 |
| 562 | An overlooked family-group name among bees: Availability of Coelioxoidini (Hymenoptera: Apidae). Journal of Melittology, 2020, , 1-3. | 0.2 | 0 |
| 563 | Genera of the bee tribe Reedapini (Hymenoptera: Colletidae). Journal of Melittology, 2020, , 1-16. | 0.2 | 0 |
| 564 | Notes on the classification of certain colletid bees (Hymenoptera: Colletidae). Entomologist's Monthly Magazine, 2020, 156, 93-101. | 0.1 | 0 |
| 565 | Optical discs in zoological nomenclature: problems and proposed solution. Bionomina, 2021, 24, . | 0.2 | 0 |
| 566 | Notes on the stingless bee genera <i>Scaura</i> and <i>Geotrigona</i> (Hymenoptera: Apidae). Journal of Melittology, 2022, , 1-4. | 0.2 | 0 |
| 567 | Two new genera of South American Eulonchopriini (Hymenoptera: Colletidae). Journal of Melittology, 2021, , 1-24. | 0.2 | 0 |
| 568 | First occurrence of the little-known genus Noteriades (Hymenoptera, Megachilidae) from Vietnam: discovery of a new species and a key to the Southeast Asian fauna. ZooKeys, 0, 1102, 191-200. | 0.5 | 0 |