

Shinji Fukuda

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8653702/publications.pdf>

Version: 2024-02-01

53
papers

759
citations

566801

15
h-index

552369

26
g-index

55
all docs

55
docs citations

55
times ranked

820
citing authors

#	ARTICLE	IF	CITATIONS
1	Species-specific debromination of BDE99 in teleost fish: The relationship between debromination ability and bioaccumulation patterns of PBDEs. <i>Science of the Total Environment</i> , 2022, 806, 151265.	3.9	3
2	Artificial lateral line for aquatic habitat modelling: An example for <i>Lefua echigonia</i> . <i>Ecological Informatics</i> , 2021, 65, 101388.	2.3	11
3	Habitat evaluation for the endangered fish species <i>Lefua echigonia</i> in the Yagawa River, Japan. <i>Journal of Ecohydraulics</i> , 2019, 4, 147-157.	1.6	7
4	Evaluation of water dynamics of contour-levee irrigation system in sloped rice fields in Colombia. <i>Agricultural Water Management</i> , 2019, 217, 107-118.	2.4	8
5	Intra- and intercontinental variation in the functional responses of a high impact alien invasive fish. <i>Biological Invasions</i> , 2019, 21, 1751-1762.	1.2	15
6	International contributions in advancing ecohydraulics. <i>Journal of Ecohydraulics</i> , 2019, 4, 86-87.	1.6	0
7	MODELLING HABITAT SUITABILITY OF <i>LEFUA ECHIGONIA</i> USING RANDOM FORESTS. <i>Journal of Japan Society of Civil Engineers Ser B1 (Hydraulic Engineering)</i> , 2019, 75, 1_541-1_546.	0.0	0
8	Estimating Soil Water Contents from Field Water Tables for Potential Rice Irrigation Criteria under Contour-Levee Irrigation Systems. <i>Environmental Control in Biology</i> , 2019, 57, 15-21.	0.3	1
9	Effect of Storage Conditions on the Postharvest Quality Changes of Fresh Mango Fruits for Export during Transportation. <i>Environmental Control in Biology</i> , 2018, 56, 39-44.	0.3	6
10	Revisiting probabilistic neural networks: a comparative study with support vector machines and the microhabitat suitability for the Eastern Iberian chub (<i>Squalius valentinus</i>). <i>Ecological Informatics</i> , 2018, 43, 24-37.	2.3	17
11	Flow regime shapes seasonal patterns of fish species richness and abundance in main and branch channels of a rice irrigation system. <i>Paddy and Water Environment</i> , 2018, 16, 783-793.	1.0	2
12	Quality Changes in Fresh Mango Fruits (<i>Mangifera indica</i> ; L. "Nam Dok Mai"™) Under Actual Distribution Temperature Profile from Thailand to Japan. <i>Environmental Control in Biology</i> , 2018, 56, 45-49.	0.3	12
13	Effects of water temperature and light intensity on the acute toxicity of herbicide thiobencarb to a green alga, <i>Raphidocelis subcapitata</i> . <i>Environmental Science and Pollution Research</i> , 2018, 25, 25363-25370.	2.7	3
14	Random Forests as a Tool for Analyzing Partial Drought Stress Based on CO ₂ Concentrations in the Rootzone of Longan Trees. <i>Environmental Control in Biology</i> , 2018, 56, 25-31.	0.3	1
15	Sensors and Monitoring for Production and Distribution of a Tropical Fruit. <i>Environmental Control in Biology</i> , 2018, 56, 23-24.	0.3	0
16	Comparing four methods for decision-tree induction: A case study on the invasive Iberian gudgeon (<i>Tj ETQq0 0 0 r0BT /Overlock 10 Tf</i>)	2.3	23
17	Data prevalence matters when assessing species' responses using data-driven species distribution models. <i>Ecological Informatics</i> , 2016, 32, 69-78.	2.3	25
18	Assessment of Depth Measurement Using an Acoustic Doppler Current Profiler and a CTD Profiler in a Small River in Japan. <i>Lecture Notes in Computer Science</i> , 2016, , 308-316.	1.0	1

#	ARTICLE	IF	CITATIONS
19	Online Monitoring System on Controlled Irrigation Experiment for Export Quality Mango in Thailand. Lecture Notes in Computer Science, 2016, , 328-334.	1.0	2
20	Sensitivity-Based Calibration of the Soil and Water Assessment Tool for Hydrologic Cycle Simulation in the Cong Watershed, Vietnam. Water Environment Research, 2015, 87, 735-750.	1.3	4
21	Modeling the Relationship between Hormone Dynamics and Off-season Flowering of Litchi by Using Random Forests. Agriculture and Agricultural Science Procedia, 2015, 5, 9-16.	0.6	2
22	Assessment of spatial habitat heterogeneity by coupling data-driven habitat suitability models with a 2D hydrodynamic model in small-scale streams. Ecological Informatics, 2015, 29, 147-155.	2.3	23
23	Comparison of the growth traits of a commercial pioneer tree species, paper mulberry (<i>Broussonetia</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 5 mechanisms underlying shade-intolerance. Agroforestry Systems, 2014, 88, 907-919.	0.9	4
24	Elevated water temperature reduces the acute toxicity of the widely used herbicide diuron to a green alga, <i>Pseudokirchneriella subcapitata</i> . Environmental Science and Pollution Research, 2014, 21, 1064-1070.	2.7	26
25	Application of a simple genetic algorithm for the calibration of aquatic ecosystem model of an agricultural pond. Paddy and Water Environment, 2014, 12, 1-15.	1.0	1
26	Modelling the relationship between peel colour and the quality of fresh mango fruit using Random Forests. Journal of Food Engineering, 2014, 131, 7-17.	2.7	42
27	Dynamics of Water Qualities under the Anaerobic and Reductive State in an Organically Polluted Closed Water Body. Journal of Rainwater Catchment Systems, 2014, 20, 49-55.	0.2	5
28	Habitat prediction and knowledge extraction for spawning European grayling (<i>Thymallus thymallus</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5 47, 1-6.	1.9	111
29	Effects of data prevalence on species distribution modelling using a genetic takagi-sugeno fuzzy system. , 2013, , .		0
30	Modelling the distribution of the panâ€œcontinental invasive fish <i>Pseudorasbora parva</i> based on landscape features in the northern Kyushu Island, Japan. Aquatic Conservation: Marine and Freshwater Ecosystems, 2013, 23, 901-910.	0.9	7
31	Random Forests modelling for the estimation of mango (<i>Mangifera indica</i> L. cv. Chok Anan) fruit yields under different irrigation regimes. Agricultural Water Management, 2013, 116, 142-150.	2.4	76
32	Predicting potential hybridization between native and non-native <i>Rhodeus ocellatus</i> subspecies: the implications for conservation of a pure native population in northern Kyushu, Japan. Aquatic Invasions, 2013, 8, 219-229.	0.6	8
33	Evaluation of Soil Water Management Difference in Mango Orchards between Thailand and Japan. American Journal of Plant Sciences, 2013, 04, 182-187.	0.3	2
34	DO ABSENCE DATA MATTER WHEN MODELLING FISH HABITAT PREFERENCE USING A GENETIC TAKAGI-SUGENO FUZZY MODEL?. International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems, 2012, 20, 233-245.	0.9	3
35	Effect of aggregation functions on the habitat preference modelling using a genetic Takagi-Sugeno fuzzy system. , 2012, , .		1
36	A short review on the application of computational intelligence and machine learning in the bioenvironmental sciences. , 2012, , .		1

#	ARTICLE	IF	CITATIONS
37	Abundance versus presence/absence data for modelling fish habitat preference with a genetic Takagi-Sugeno fuzzy system. <i>Environmental Monitoring and Assessment</i> , 2012, 184, 6159-6171.	1.3	25
38	Predicting distributions of seven bitterling fishes in northern Kyushu, Japan. <i>Ichthyological Research</i> , 2012, 59, 124-133.	0.5	18
39	A discussion on the accuracy-complexity relationship in modelling fish habitat preference using genetic Takagi-Sugeno fuzzy systems. , 2011, , .		1
40	Assessing the applicability of fuzzy neural networks for habitat preference evaluation of Japanese medaka (<i>Oryzias latipes</i>). <i>Ecological Informatics</i> , 2011, 6, 286-295.	2.3	19
41	Effect of model formulation on the optimization of a genetic Takagi-Sugeno fuzzy system for fish habitat suitability evaluation. <i>Ecological Modelling</i> , 2011, 222, 1401-1413.	1.2	52
42	Evaluating the potential for invasion by alien freshwater fishes in northern Kyushu Island, Japan, using the Fish Invasiveness Scoring Kit. <i>Ichthyological Research</i> , 2011, 58, 382-387.	0.5	33
43	Assessing the effects of zero abundance data on habitat preference modelling using a genetic Takagi-Sugeno fuzzy model. , 2011, , .		2
44	Modelling Fish Habitat Preference with a Genetic Algorithm-Optimized Takagi-Sugeno Model Based on Pairwise Comparisons. <i>Advances in Intelligent and Soft Computing</i> , 2011, , 375-387.	0.2	0
45	The application of entropy for detecting behavioral responses in Japanese medaka (<i>Oryzias latipes</i>). <i>Ecological Modelling</i> , 2011, 222, 1401-1413.	1.2	52
46	Effect of data quality on habitat preference evaluation for Japanese medaka (<i>Oryzias latipes</i>) using a simple genetic fuzzy system. , 2010, , .		5
47	A genetic Takagi-Sugeno fuzzy system for fish habitat preference modelling. , 2010, , .		1
48	Consideration of fuzziness: Is it necessary in modelling fish habitat preference of Japanese medaka (<i>Oryzias latipes</i>)?. <i>Ecological Modelling</i> , 2009, 220, 2877-2884.	1.2	48
49	Coelomycete systematics with special reference to <i>Colletotrichum</i> . <i>Mycoscience</i> , 2008, 49, 373-378.	0.3	6
50	Prediction ability and sensitivity of artificial intelligence-based habitat preference models for predicting spatial distribution of Japanese medaka (<i>Oryzias latipes</i>). <i>Ecological Modelling</i> , 2008, 215, 301-313.	1.2	31
51	Assessing Nonlinearity in Fish Habitat Preference of Japanese Medaka (<i>Oryzias latipes</i>) Using Genetic Algorithm-Optimized Habitat Prediction Models. <i>Japan Agricultural Research Quarterly</i> , 2008, 42, 97-107.	0.1	14
52	Fuzzy neural network model for habitat prediction and HEP for habitat quality estimation focusing on Japanese medaka (<i>Oryzias latipes</i>) in agricultural canals. <i>Paddy and Water Environment</i> , 2006, 4, 119-124.	1.0	19
53	Numerical quantification of the significance of aquatic vegetation affecting spatial distribution of Japanese medaka (<i>Oryzias latipes</i>) in an agricultural canal. <i>Landscape and Ecological Engineering</i> , 2006, 2, 65-80.	0.7	13