

Catherine Werts

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8655180/catherine-werts-publications-by-year.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

75
papers

4,622
citations

32
h-index

67
g-index

82
ext. papers

5,212
ext. citations

8
avg, IF

5.04
L-index

#	Paper	IF	Citations
75	Transient Presence of Live <i>Leptospira interrogans</i> in Murine Testes.. <i>Microbiology Spectrum</i> , 2022 , e0277531	7.9	0
74	Anti- <i>Leptospira</i> immunoglobulin profiling in mice reveals strain specific IgG and persistent IgM responses associated with virulence and renal colonization. <i>PLoS Neglected Tropical Diseases</i> , 2021 , 15, e0008970	4.8	5
73	The route of infection with <i>Leptospira interrogans</i> serovar Copenhageni affects the kinetics of bacterial dissemination and kidney colonization. <i>PLoS Neglected Tropical Diseases</i> , 2020 , 14, e0007950	4.8	14
72	Leptospiral LPS escapes mouse TLR4 internalization and TRIF-associated antimicrobial responses through O antigen and associated lipoproteins. <i>PLoS Pathogens</i> , 2020 , 16, e1008639	7.6	12
71	Phagocyte Escape of : The Role of TLRs and NLRs. <i>Frontiers in Immunology</i> , 2020 , 11, 571816	8.4	12
70	Escape of TLR5 Recognition by spp.: A Rationale for Atypical Endoflagella. <i>Frontiers in Immunology</i> , 2020 , 11, 2007	8.4	10
69	Role of TLR4 in Persistent Infection: A Comparative Study in Mice. <i>Frontiers in Immunology</i> , 2020 , 11, 572999	8.4	3
68	In Vivo Imaging of Bioluminescent <i>Leptospira</i> s. <i>Methods in Molecular Biology</i> , 2020 , 2134, 149-160	1.4	3
67	Purification of LPS from <i>Leptospira</i> . <i>Methods in Molecular Biology</i> , 2020 , 2134, 53-65	1.4	2
66	Leptospiral LPS escapes mouse TLR4 internalization and TRIF-associated antimicrobial responses through O antigen and associated lipoproteins 2020 , 16, e1008639		
65	Leptospiral LPS escapes mouse TLR4 internalization and TRIF-associated antimicrobial responses through O antigen and associated lipoproteins 2020 , 16, e1008639		
64	Leptospiral LPS escapes mouse TLR4 internalization and TRIF-associated antimicrobial responses through O antigen and associated lipoproteins 2020 , 16, e1008639		
63	Leptospiral LPS escapes mouse TLR4 internalization and TRIF-associated antimicrobial responses through O antigen and associated lipoproteins 2020 , 16, e1008639		
62	Innate immune memory through TLR2 and NOD2 contributes to the control of <i>Leptospira interrogans</i> infection. <i>PLoS Pathogens</i> , 2019 , 15, e1007811	7.6	32
61	Murine Models for Leptospirosis Kidney Disease. <i>Translational Research in Biomedicine</i> , 2019 , 65-75	0.1	0
60	Interaction of <i>Leptospira</i> with the Innate Immune System. <i>Current Topics in Microbiology and Immunology</i> , 2018 , 415, 163-187	3.3	13
59	Recent findings related to immune responses against leptospirosis and novel strategies to prevent infection. <i>Microbes and Infection</i> , 2018 , 20, 578-588	9.3	12

58	Mycobacterial Phenolic Glycolipids Selectively Disable TRIF-Dependent TLR4 Signaling in Macrophages. <i>Frontiers in Immunology</i> , 2018 , 9, 2	8.4	16
57	LIM-Only Protein FHL2 Is a Negative Regulator of Transforming Growth Factor β Expression. <i>Molecular and Cellular Biology</i> , 2017 , 37,	4.8	12
56	LipL21 lipoprotein binding to peptidoglycan enables <i>Leptospira interrogans</i> to escape NOD1 and NOD2 recognition. <i>PLoS Pathogens</i> , 2017 , 13, e1006725	7.6	26
55	Animal Models of Leptospirosis: Of Mice and Hamsters. <i>Frontiers in Immunology</i> , 2017 , 8, 58	8.4	51
54	Pre-treatment with <i>Lactobacillus plantarum</i> prevents severe pathogenesis in mice infected with <i>Leptospira interrogans</i> and may be associated with recruitment of myeloid cells. <i>PLoS Neglected Tropical Diseases</i> , 2017 , 11, e0005870	4.8	19
53	CCL20 Displays Antimicrobial Activity Against <i>Cryptosporidium parvum</i> , but Its Expression Is Reduced During Infection in the Intestine of Neonatal Mice. <i>Journal of Infectious Diseases</i> , 2015 , 212, 1332-40	7	25
52	Common Cell Shape Evolution of Two Nasopharyngeal Pathogens. <i>PLoS Genetics</i> , 2015 , 11, e1005338	6	21
51	Peptidoglycan and Nod Receptor 2015 , 737-747		5
50	Calcineurin/NFAT signaling and innate host defence: a role for NOD1-mediated phagocytic functions. <i>Cell Communication and Signaling</i> , 2014 , 12, 8	7.5	32
49	Poly(I:C)-induced protection of neonatal mice against intestinal <i>Cryptosporidium parvum</i> infection requires an additional TLR5 signal provided by the gut flora. <i>Journal of Infectious Diseases</i> , 2014 , 209, 457-67	7	33
48	Flagellin/TLR5 signalling activates renal collecting duct cells and facilitates invasion and cellular translocation of uropathogenic <i>Escherichia coli</i> . <i>Cellular Microbiology</i> , 2014 , 16, 1503-17	3.9	21
47	<i>Leptospira Interrogans</i> induces fibrosis in the mouse kidney through Inos-dependent, TLR- and NLR-independent signaling pathways. <i>PLoS Neglected Tropical Diseases</i> , 2014 , 8, e2664	4.8	48
46	Live imaging of bioluminescent <i>leptospira interrogans</i> in mice reveals renal colonization as a stealth escape from the blood defenses and antibiotics. <i>PLoS Neglected Tropical Diseases</i> , 2014 , 8, e3359	4.8	54
45	CCL17 production by dendritic cells is required for NOD1-mediated exacerbation of allergic asthma. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2014 , 189, 899-908	10.2	19
44	Peptidoglycan and Nod Receptor 2014 , 1-10		3
43	Penicillin resistance compromises Nod1-dependent proinflammatory activity and virulence fitness of <i>neisseria meningitidis</i> . <i>Cell Host and Microbe</i> , 2013 , 13, 735-45	23.4	20
42	Intestinal CD103+ dendritic cells are key players in the innate immune control of <i>Cryptosporidium parvum</i> infection in neonatal mice. <i>PLoS Pathogens</i> , 2013 , 9, e1003801	7.6	54
41	Cyclosporine A impairs nucleotide binding oligomerization domain (Nod1)-mediated innate antibacterial renal defenses in mice and human transplant recipients. <i>PLoS Pathogens</i> , 2013 , 9, e1003152	7.6	38

40	LIM-only protein FHL2 activates NF- κ B signaling in the control of liver regeneration and hepatocarcinogenesis. <i>Molecular and Cellular Biology</i> , 2013 , 33, 3299-308	4.8	29
39	The effect of bulgecin A on peptidoglycan metabolism and physiology of <i>Helicobacter pylori</i> . <i>Microbial Drug Resistance</i> , 2012 , 18, 230-9	2.9	17
38	Downregulation of the Na/K-ATPase pump by leptospiral glycolipoprotein activates the NLRP3 inflammasome. <i>Journal of Immunology</i> , 2012 , 188, 2805-14	5.3	57
37	Nod-like receptors in intestinal homeostasis, inflammation, and cancer. <i>Journal of Leukocyte Biology</i> , 2011 , 90, 471-82	6.5	44
36	Heat shock protein gp96 and NAD(P)H oxidase 4 play key roles in Toll-like receptor 4-activated apoptosis during renal ischemia/reperfusion injury. <i>Cell Death and Differentiation</i> , 2010 , 17, 1474-85	12.7	71
35	TLR9-dependent induction of intestinal alpha-defensins by <i>Toxoplasma gondii</i> . <i>Journal of Immunology</i> , 2010 , 184, 7022-9	5.3	57
34	Leptospirosis: a Toll road from B lymphocytes. <i>Chang Gung Medical Journal</i> , 2010 , 33, 591-601		18
33	TLR4- and TLR2-mediated B cell responses control the clearance of the bacterial pathogen, <i>Leptospira interrogans</i> . <i>Journal of Immunology</i> , 2009 , 183, 2669-77	5.3	96
32	Heat shock protein gp96 interacts with protein phosphatase 5 and controls toll-like receptor 2 (TLR2)-mediated activation of extracellular signal-regulated kinase (ERK) 1/2 in post-hypoxic kidney cells. <i>Journal of Biological Chemistry</i> , 2009 , 284, 12541-9	5.4	32
31	Lymphoid tissue genesis induced by commensals through NOD1 regulates intestinal homeostasis. <i>Nature</i> , 2008 , 456, 507-10	50.4	779
30	The ubiquitin-editing enzyme A20 restricts nucleotide-binding oligomerization domain containing 2-triggered signals. <i>Immunity</i> , 2008 , 28, 381-90	32.3	262
29	Nod1 and Nod2 induce CCL5/RANTES through the NF-kappaB pathway. <i>European Journal of Immunology</i> , 2007 , 37, 2499-508	6.1	68
28	Role of Nods in bacterial infection. <i>Microbes and Infection</i> , 2007 , 9, 629-36	9.3	38
27	Simple method for repurification of endotoxins for biological use. <i>Applied and Environmental Microbiology</i> , 2007 , 73, 1803-8	4.8	41
26	<i>Toxoplasma gondii</i> and subversion of the immune system. <i>Trends in Parasitology</i> , 2006 , 22, 448-52	6.4	32
25	Role of AmiA in the morphological transition of <i>Helicobacter pylori</i> and in immune escape. <i>PLoS Pathogens</i> , 2006 , 2, e97	7.6	89
24	Renal collecting duct epithelial cells react to pyelonephritis-associated <i>Escherichia coli</i> by activating distinct TLR4-dependent and -independent inflammatory pathways. <i>Journal of Immunology</i> , 2006 , 177, 4773-84	5.3	104
23	TIR, CARD and PYRIN: three domains for an antimicrobial triad. <i>Cell Death and Differentiation</i> , 2006 , 13, 798-815	12.7	114

22	Differential TLR recognition of leptospiral lipid A and lipopolysaccharide in murine and human cells. <i>Journal of Immunology</i> , 2005 , 175, 6022-31	5.3	140
21	Synergistic stimulation of human monocytes and dendritic cells by Toll-like receptor 4 and NOD1- and NOD2-activating agonists. <i>European Journal of Immunology</i> , 2005 , 35, 2459-70	6.1	276
20	The frameshift mutation in Nod2 results in unresponsiveness not only to Nod2- but also Nod1-activating peptidoglycan agonists. <i>Journal of Biological Chemistry</i> , 2005 , 280, 35859-67	5.4	67
19	Enzymatic synthesis of lipid A molecules with four amide-linked acyl chains. LpxA acyltransferases selective for an analog of UDP-N-acetylglucosamine in which an amine replaces the 3"-hydroxyl group. <i>Journal of Biological Chemistry</i> , 2004 , 279, 25411-9	5.4	31
18	A methylated phosphate group and four amide-linked acyl chains in leptospira interrogans lipid A. The membrane anchor of an unusual lipopolysaccharide that activates TLR2. <i>Journal of Biological Chemistry</i> , 2004 , 279, 25420-9	5.4	110
17	Helicobacter pylori heat shock protein 60 mediates interleukin-6 production by macrophages via a toll-like receptor (TLR)-2-, TLR-4-, and myeloid differentiation factor 88-independent mechanism. <i>Journal of Biological Chemistry</i> , 2004 , 279, 245-50	5.4	127
16	Toll-like receptor 2-dependent bacterial sensing does not occur via peptidoglycan recognition. <i>EMBO Reports</i> , 2004 , 5, 1000-6	6.5	390
15	Toll-like receptor-mediated tumor necrosis factor and interleukin-10 production differ during systemic inflammation. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2003 , 168, 158-64	10.2	91
14	Inhibition by eicosapentaenoic acid of IL-1beta-induced PGHS-2 expression in human microvascular endothelial cells: involvement of lipoxygenase-derived metabolites and p38 MAPK pathway. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2003 , 1631, 77-84	5	20
13	TNF-alpha, inefficient by itself, potentiates IL-1beta-induced PGHS-2 expression in human pulmonary microvascular endothelial cells: requirement of NF-kappaB and p38 MAPK pathways. <i>British Journal of Pharmacology</i> , 2002 , 136, 1005-14	8.6	19
12	First evidence for a restriction-modification system in Leptospira sp. <i>FEMS Microbiology Letters</i> , 2001 , 201, 139-43	2.9	4
11	Leptospiral lipopolysaccharide activates cells through a TLR2-dependent mechanism. <i>Nature Immunology</i> , 2001 , 2, 346-52	19.1	545
10	A role for residue 151 of LamB in bacteriophage lambda adsorption: possible steric effect of amino acid substitutions. <i>Journal of Bacteriology</i> , 1994 , 176, 3204-9	3.5	23
9	Adsorption of bacteriophage lambda on the LamB protein of Escherichia coli K-12: point mutations in gene J of lambda responsible for extended host range. <i>Journal of Bacteriology</i> , 1994 , 176, 941-7	3.5	61
8	Construction and first characterization of two reciprocal hybrids between LamB from Escherichia coli K12 and Klebsiella pneumoniae. <i>Research in Microbiology</i> , 1993 , 144, 259-69	4	2
7	Immunological relatedness of the LamB proteins among members of Enterobacteriaceae. <i>Journal of General Microbiology</i> , 1993 , 139, 881-7		11
6	The Maltose B Region in Salmonella Typhimurium, Escherichia Coli and Other Enterobacteriaceae 1993 , 91-104		0
5	DNA sequence analysis of the lamB gene from Klebsiella pneumoniae: implications for the topology and the pore functions in maltoporin. <i>Molecular Genetics and Genomics</i> , 1992 , 233, 372-8		21

4	A genetic system to elicit and monitor antipeptide antibodies without peptide synthesis. <i>Nature Biotechnology</i> , 1991 , 9, 170-2	44.5	27
3	Permissive sites and topology of an outer membrane protein with a reporter epitope. <i>Journal of Bacteriology</i> , 1991 , 173, 262-75	3.5	86
2	Role of TLR4 in persistent <i>Leptospira interrogans</i> infection: a comparative in vivo study in mice		1
1	Leptospiral LPS escapes mouse TLR4 internalization and TRIF-associated antimicrobial responses through O antigen and associated lipoproteins		3