

Hong-Yu Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8655491/publications.pdf>

Version: 2024-02-01

13
papers

294
citations

1040056

9
h-index

940533

16
g-index

17
all docs

17
docs citations

17
times ranked

413
citing authors

#	ARTICLE	IF	CITATIONS
1	Potential Therapeutic Targets of Rehmannia Formulations on Diabetic Nephropathy: A Comparative Network Pharmacology Analysis. <i>Frontiers in Pharmacology</i> , 2022, 13, 794139.	3.5	8
2	Pseudolaric acid B ameliorates synovial inflammation and vessel formation by stabilizing PPAR β to inhibit NF κ B signalling pathway. <i>Journal of Cellular and Molecular Medicine</i> , 2021, 25, 6664-6678.	3.6	10
3	Tubule-specific deletion of LincRNA-p21 ameliorates lipotoxic kidney injury. <i>Molecular Therapy - Nucleic Acids</i> , 2021, 26, 1280-1290.	5.1	3
4	Patients TM and clinicians TM expectations on integrative medicine Services for Diabetes: a focus group study. <i>BMC Complementary Medicine and Therapies</i> , 2020, 20, 205.	2.7	9
5	The PAR-1 antagonist vorapaxar ameliorates kidney injury and tubulointerstitial fibrosis. <i>Clinical Science</i> , 2020, 134, 2873-2891.	4.3	20
6	Dimethylaminomicheliolide ameliorates peritoneal fibrosis through the activation of autophagy. <i>Journal of Molecular Medicine</i> , 2019, 97, 659-674.	3.9	21
7	Micheliolide ameliorates renal fibrosis by suppressing the Mtdh/BMP/MAPK pathway. <i>Laboratory Investigation</i> , 2019, 99, 1092-1106.	3.7	25
8	Micheliolide alleviates hepatic steatosis in db/db mice by inhibiting inflammation and promoting autophagy via PPAR- β -mediated NF- κ B and AMPK/mTOR signaling. <i>International Immunopharmacology</i> , 2018, 59, 197-208.	3.8	50
9	Irbesartan ameliorates hyperlipidemia and liver steatosis in type 2 diabetic db/db mice via stimulating PPAR- β , AMPK/Akt/mTOR signaling and autophagy. <i>International Immunopharmacology</i> , 2017, 42, 176-184.	3.8	56
10	Irbesartan Ameliorates Diabetic Nephropathy by Suppressing the RANKL-RANK-NF- κ B Pathway in Type 2 Diabetic db/db Mice. <i>Mediators of Inflammation</i> , 2016, 2016, 1-10.	3.0	10
11	Metadherin facilitates podocyte apoptosis in diabetic nephropathy. <i>Cell Death and Disease</i> , 2016, 7, e2477-e2477.	6.3	54
12	Cyclopropanyldehydrocostunolide LJ attenuates high glucose-induced podocyte injury by suppressing RANKL/RANK-mediated NF- κ B and MAPK signaling pathways. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 760-769.	2.3	14
13	Aprikalim reduces the Na $^{+}$ -Ca $^{2+}$ exchange outward current enhanced by hyperkalemia in rat ventricular myocytes. <i>Annals of Thoracic Surgery</i> , 2002, 73, 1253-1259.	1.3	10