

Satu Pekkala

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8668453/satu-pekkala-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

31
papers

946
citations

15
h-index

30
g-index

36
ext. papers

1,201
ext. citations

5.1
avg, IF

3.71
L-index

#	Paper	IF	Citations
31	Xylo-Oligosaccharides in Prevention of Hepatic Steatosis and Adipose Tissue Inflammation: Associating Taxonomic and Metabolomic Patterns in Fecal Microbiomes with Biclustering. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	1
30	Vascular Adhesion Protein 1 Mediates Gut Microbial Flagellin-Induced Inflammation, Leukocyte Infiltration, and Hepatic Steatosis. <i>Sci</i> , 2021 , 3, 13	0.7	1
29	Higher glucose availability augments the metabolic responses of the C2C12 myotubes to exercise-like electrical pulse stimulation. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2021 , 321, E229-E245	6	3
28	Rats bred for low intrinsic aerobic exercise capacity link obesity with brain inflammation and reduced structural plasticity of the hippocampus. <i>Brain, Behavior, and Immunity</i> , 2021 , 97, 250-259	16.6	2
27	Irradiation of the head reduces adult hippocampal neurogenesis and impairs spatial memory, but leaves overall health intact in rats. <i>European Journal of Neuroscience</i> , 2021 , 53, 1885-1904	3.5	1
26	Differentiation of Murine C2C12 Myoblasts Strongly Reduces the Effects of Myostatin on Intracellular Signaling. <i>Biomolecules</i> , 2020 , 10,	5.9	11
25	Prebiotic Xylo-Oligosaccharides Ameliorate High-Fat-Diet-Induced Hepatic Steatosis in Rats. <i>Nutrients</i> , 2020 , 12,	6.7	13
24	Beneficial effects of running and milk protein supplements on Sirtuins and risk factors of metabolic disorders in rats with low aerobic capacity. <i>Metabolism Open</i> , 2019 , 4, 100019	2.8	3
23	Blocking Activin Receptor Ligands Is Not Sufficient to Rescue Cancer-Associated Gut Microbiota-A Role for Gut Microbial Flagellin in Colorectal Cancer and Cachexia?. <i>Cancers</i> , 2019 , 11,	6.6	8
22	Shorter recovery time following high-intensity interval training induced higher body fat loss among overweight women. <i>Sport Sciences for Health</i> , 2019 , 15, 157-165	1.3	2
21	Enterobacter cloacae administration induces hepatic damage and subcutaneous fat accumulation in high-fat diet fed mice. <i>PLoS ONE</i> , 2018 , 13, e0198262	3.7	13
20	Six-Week Endurance Exercise Alters Gut Metagenome That Is not Reflected in Systemic Metabolism in Over-weight Women. <i>Frontiers in Microbiology</i> , 2018 , 9, 2323	5.7	85
19	Faecalibacterium prausnitzii treatment improves hepatic health and reduces adipose tissue inflammation in high-fat fed mice. <i>ISME Journal</i> , 2017 , 11, 1667-1679	11.9	98
18	Intrinsic aerobic capacity governs the associations between gut microbiota composition and fat metabolism age-dependently in rat siblings. <i>Physiological Genomics</i> , 2017 , 49, 733-746	3.6	10
17	Gut Microbiota Analysis Results Are Highly Dependent on the 16S rRNA Gene Target Region, Whereas the Impact of DNA Extraction Is Minor. <i>Journal of Biomolecular Techniques</i> , 2017 , 28, 19-30	1.1	87
16	Insulin resistance is associated with altered amino acid metabolism and adipose tissue dysfunction in normoglycemic women. <i>Scientific Reports</i> , 2016 , 6, 24540	4.9	39
15	Adipocytes as a Link Between Gut Microbiota-Derived Flagellin and Hepatocyte Fat Accumulation. <i>PLoS ONE</i> , 2016 , 11, e0152786	3.7	11

14	Cannabinoid receptor 1 and acute resistance exercise--In vivo and in vitro studies in human skeletal muscle. <i>Peptides</i> , 2015 , 67, 55-63	3.8	6
13	PGC-1 isoforms and their target genes are expressed differently in human skeletal muscle following resistance and endurance exercise. <i>Physiological Reports</i> , 2015 , 3, e12563	2.6	44
12	Toll-like receptor 5 in obesity: the role of gut microbiota and adipose tissue inflammation. <i>Obesity</i> , 2015 , 23, 581-90	8	43
11	Does systemic low-grade inflammation associate with fat accumulation and distribution? A 7-year follow-up study with peripubertal girls. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 1411-5	5.6	13
10	Metabolic response to 6-week aerobic exercise training and dieting in previously sedentary overweight and obese pre-menopausal women: A randomized trial. <i>Journal of Sport and Health Science</i> , 2014 , 3, 217-224	8.2	15
9	Gut-adipose tissue axis in hepatic fat accumulation in humans. <i>Journal of Hepatology</i> , 2014 , 61, 132-8	13.4	36
8	Molecular characterization of carbamoyl-phosphate synthetase (CPS1) deficiency using human recombinant CPS1 as a key tool. <i>Human Mutation</i> , 2013 , 34, 1149-59	4.7	29
7	Exercise, the endocannabinoid system and metabolic health. <i>Journal of Sport and Health Science</i> , 2013 , 2, 60-61	8.2	1
6	Are skeletal muscle FNDC5 gene expression and irisin release regulated by exercise and related to health?. <i>Journal of Physiology</i> , 2013 , 591, 5393-400	3.9	170
5	Women with and without metabolic disorder differ in their gut microbiota composition. <i>Obesity</i> , 2012 , 20, 1082-7	8	65
4	Serum osteocalcin is not associated with glucose but is inversely associated with leptin across generations of nondiabetic women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, 4106-14	5.6	38
3	Genetic, structural and biochemical basis of carbamoyl phosphate synthetase 1 deficiency. <i>Molecular Genetics and Metabolism</i> , 2010 , 101, 311-23	3.7	43
2	Understanding carbamoyl-phosphate synthetase I (CPS1) deficiency by using expression studies and structure-based analysis. <i>Human Mutation</i> , 2010 , 31, 801-8	4.7	29
1	Structural insight on the control of urea synthesis: identification of the binding site for N-acetyl-L-glutamate, the essential allosteric activator of mitochondrial carbamoyl phosphate synthetase. <i>Biochemical Journal</i> , 2009 , 424, 211-20	3.8	22