

Nibaldo Inestrosa Cantin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8674145/publications.pdf>

Version: 2024-02-01

330
papers

19,413
citations

11235

73
h-index

20023

121
g-index

332
all docs

332
docs citations

332
times ranked

19864
citing authors

#	ARTICLE	IF	CITATIONS
1	Age- and Sex-Associated Glucose Metabolism Decline in a Mouse Model of Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2022, , 1-17.	1.2	3
2	Differential Role of Sex and Age in the Synaptic Transmission of Degus (<i>Octodon degus</i>). <i>Frontiers in Integrative Neuroscience</i> , 2022, 16, 799147.	1.0	1
3	“Live together, die alone”: The effect of re-socialization on behavioural performance and social-affective brain-related proteins after a long-term chronic social isolation stress. <i>Neurobiology of Stress</i> , 2021, 14, 100289.	1.9	12
4	Andrographolide restores glucose uptake in rat hippocampal neurons. <i>Journal of Neurochemistry</i> , 2021, 157, 1222-1233.	2.1	11
5	Wnt5a promotes hippocampal postsynaptic development and GluN2B-induced expression via the eIF2 β HRI kinase. <i>Scientific Reports</i> , 2021, 11, 7395.	1.6	8
6	Wnt5a modulates dendritic spine dynamics through the regulation of Cofilin via small Rho GTPase activity in hippocampal neurons. <i>Journal of Neurochemistry</i> , 2021, 158, 673-693.	2.1	5
7	A Multivariate Assessment of Age-Related Cognitive Impairment in <i>Octodon degus</i> . <i>Frontiers in Integrative Neuroscience</i> , 2021, 15, 719076.	1.0	6
8	Morphological neurite changes induced by porcupine inhibition are rescued by Wnt ligands. <i>Cell Communication and Signaling</i> , 2021, 19, 87.	2.7	4
9	The transcriptional landscape of Alzheimer's disease and its association with Wnt signaling pathway. <i>Neuroscience and Biobehavioral Reviews</i> , 2021, 128, 454-466.	2.9	8
10	Neurodevelopmental impact of the offspring by thyroid hormone system-disrupting environmental chemicals during pregnancy. <i>Environmental Research</i> , 2021, 200, 111345.	3.7	27
11	Discovery of a Potent Dual Inhibitor of Acetylcholinesterase and Butyrylcholinesterase with Antioxidant Activity that Alleviates Alzheimer-like Pathology in Old APP/PS1 Mice. <i>Journal of Medicinal Chemistry</i> , 2021, 64, 812-839.	2.9	45
12	Selective Surface and Intraluminal Localization of Wnt Ligands on Small Extracellular Vesicles Released by HT-22 Hippocampal Neurons. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 735888.	1.8	14
13	Disruption of Glucose Metabolism in Aged <i>Octodon degus</i> : A Sporadic Model of Alzheimer's Disease. <i>Frontiers in Integrative Neuroscience</i> , 2021, 15, 733007.	1.0	2
14	Synaptic Mitochondria: An Early Target of Amyloid- β and Tau in Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2021, 84, 1391-1414.	1.2	26
15	Huperzine A and Its Neuroprotective Molecular Signaling in Alzheimer's Disease. <i>Molecules</i> , 2021, 26, 6531.	1.7	33
16	WNT Signaling Is a Key Player in Alzheimer's Disease. <i>Handbook of Experimental Pharmacology</i> , 2021, 269, 357-382.	0.9	6
17	Andrographolide promotes hippocampal neurogenesis and spatial memory in the APP ^{swe} /PS1 ^{E9} mouse model of Alzheimer's disease. <i>Scientific Reports</i> , 2021, 11, 22904.	1.6	10
18	Wnt5a promotes differentiation and development of adult-born neurons in the hippocampus by noncanonical Wnt signaling. <i>Stem Cells</i> , 2020, 38, 422-436.	1.4	53

#	ARTICLE	IF	CITATIONS
19	Andrographolide Reduces Neuroinflammation and Oxidative Stress in Aged Octodon degus. <i>Molecular Neurobiology</i> , 2020, 57, 1131-1145.	1.9	30
20	Canonical Wnt Signaling Modulates the Expression of Pre- and Postsynaptic Components in Different Temporal Patterns. <i>Molecular Neurobiology</i> , 2020, 57, 1389-1404.	1.9	14
21	Effects of long-lasting social isolation and re-socialization on cognitive performance and brain activity: a longitudinal study in Octodon degus. <i>Scientific Reports</i> , 2020, 10, 18315.	1.6	28
22	Wnt Signaling Pathway Dysregulation in the Aging Brain: Lessons From the Octodon degus. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 734.	1.8	23
23	Glutamatergic Receptor Trafficking and Delivery: Role of the Exocyst Complex. <i>Cells</i> , 2020, 9, 2402.	1.8	5
24	Toll-Like Receptors (TLRs) in Neurodegeneration: Integrative Approach to TLR Cascades in Alzheimer's and Parkinson's Diseases. , 2020, , .		2
25	Hormetic-Like Effects of L-Homocysteine on Synaptic Structure, Function, and A β Aggregation. <i>Pharmaceuticals</i> , 2020, 13, 24.	1.7	11
26	Evidence of Synaptic and Neurochemical Remodeling in the Retina of Aging Degus. <i>Frontiers in Neuroscience</i> , 2020, 14, 161.	1.4	16
27	Toward an integrative understanding of the neuroinflammatory molecular milieu in Alzheimer disease neurodegeneration. , 2020, , 163-176.		0
28	Revisiting the Paraquat-Induced Sporadic Parkinson's Disease-Like Model. <i>Molecular Neurobiology</i> , 2019, 56, 1044-1055.	1.9	65
29	Hypothyroidism and Cognitive Disorders during Development and Adulthood: Implications in the Central Nervous System. <i>Molecular Neurobiology</i> , 2019, 56, 2952-2963.	1.9	48
30	Modulation of Glucose Metabolism in Hippocampal Neurons by Adiponectin and Resistin. <i>Molecular Neurobiology</i> , 2019, 56, 3024-3037.	1.9	34
31	Wnt-7a Stimulates Dendritic Spine Morphogenesis and PSD-95 Expression Through Canonical Signaling. <i>Molecular Neurobiology</i> , 2019, 56, 1870-1882.	1.9	27
32	Fructose and prostate cancer: toward an integrated view of cancer cell metabolism. <i>Prostate Cancer and Prostatic Diseases</i> , 2019, 22, 49-58.	2.0	13
33	Molecular Basis of Neurodegeneration: Lessons from Alzheimer's and Parkinson's Diseases. , 2019, , .		2
34	Serine-Arginine Protein Kinase SRPK2 Modulates the Assembly of the Active Zone Scaffolding Protein CAST1/ERC2. <i>Cells</i> , 2019, 8, 1333.	1.8	6
35	Presymptomatic Treatment With Andrographolide Improves Brain Metabolic Markers and Cognitive Behavior in a Model of Early-Onset Alzheimer's Disease. <i>Frontiers in Cellular Neuroscience</i> , 2019, 13, 295.	1.8	34
36	Wnt Signaling Upregulates Teneurin-3 Expression via Canonical and Non-canonical Wnt Pathway Crosstalk. <i>Frontiers in Neuroscience</i> , 2019, 13, 505.	1.4	6

#	ARTICLE	IF	CITATIONS
37	GALECTIN-8 Is a Neuroprotective Factor in the Brain that Can Be Neutralized by Human Autoantibodies. <i>Molecular Neurobiology</i> , 2019, 56, 7774-7788.	1.9	22
38	Non-canonical function of IRE1 β determines mitochondria-associated endoplasmic reticulum composition to control calcium transfer and bioenergetics. <i>Nature Cell Biology</i> , 2019, 21, 755-767.	4.6	168
39	Modulating Wnt signaling at the root: Porcupine and Wnt acylation. , 2019, 198, 34-45.		65
40	MicroRNAs in Metabolic Syndrome. , 2019, , 709-725.		0
41	The Exocyst Component Exo70 Modulates Dendrite Arbor Formation, Synapse Density, and Spine Maturation in Primary Hippocampal Neurons. <i>Molecular Neurobiology</i> , 2019, 56, 4620-4638.	1.9	19
42	Wnt β -induced activation of glucose metabolism mediates the <i>in vivo</i> neuroprotective roles of Wnt signaling in Alzheimer disease. <i>Journal of Neurochemistry</i> , 2019, 149, 54-72.	2.1	49
43	Local Klotho Enhances Neuronal Progenitor Proliferation in the Adult Hippocampus. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2019, 74, 1043-1051.	1.7	15
44	Cognitive impairment in heart failure is associated with altered Wnt signaling in the hippocampus. <i>Aging</i> , 2019, 11, 5924-5942.	1.4	23
45	Diterpenes and the crosstalk with the arachidonic acid pathways, relevance in neurodegeneration. <i>Neural Regeneration Research</i> , 2019, 14, 1705.	1.6	1
46	Neuroprotective Effects of Ferruginol, Jatrophone, and Junicedric Acid Against Amyloid- β Injury in Hippocampal Neurons. <i>Journal of Alzheimer's Disease</i> , 2018, 63, 705-723.	1.2	8
47	Nicotine Modulates Mitochondrial Dynamics in Hippocampal Neurons. <i>Molecular Neurobiology</i> , 2018, 55, 8965-8977.	1.9	13
48	Long-Term, Fructose-Induced Metabolic Syndrome-Like Condition Is Associated with Higher Metabolism, Reduced Synaptic Plasticity and Cognitive Impairment in <i>Octodon degus</i> . <i>Molecular Neurobiology</i> , 2018, 55, 9169-9187.	1.9	16
49	APP/Go protein G β -complex signaling mediates A β degeneration and cognitive impairment in Alzheimer's disease models. <i>Neurobiology of Aging</i> , 2018, 64, 44-57.	1.5	15
50	Wnt Signaling in the Central Nervous System: New Insights in Health and Disease. <i>Progress in Molecular Biology and Translational Science</i> , 2018, 153, 81-130.	0.9	68
51	Wnt3a ligand facilitates autophagy in hippocampal neurons by modulating a novel GSK-3 β -AMPK axis. <i>Cell Communication and Signaling</i> , 2018, 16, 15.	2.7	36
52	Vertebrate Presynaptic Active Zone Assembly: a Role Accomplished by Diverse Molecular and Cellular Mechanisms. <i>Molecular Neurobiology</i> , 2018, 55, 4513-4528.	1.9	23
53	Wnt signaling loss accelerates the appearance of neuropathological hallmarks of Alzheimer's disease in J20 β APP transgenic and wild-type mice. <i>Journal of Neurochemistry</i> , 2018, 144, 443-465.	2.1	66
54	New Insights into the Spontaneous Human Alzheimer's Disease-Like Model <i>Octodon degus</i> : Unraveling Amyloid- β Peptide Aggregation and Age-Related Amyloid Pathology. <i>Journal of Alzheimer's Disease</i> , 2018, 66, 1145-1163.	1.2	21

#	ARTICLE	IF	CITATIONS
55	Loss of canonical Wnt signaling is involved in the pathogenesis of Alzheimer's disease. <i>Neural Regeneration Research</i> , 2018, 13, 1705.	1.6	100
56	PSD-95 (Postsynaptic Density Protein-95)., 2018, , 4263-4269.		0
57	Posttranslational Modifications Regulate the Postsynaptic Localization of PSD-95. <i>Molecular Neurobiology</i> , 2017, 54, 1759-1776.	1.9	60
58	Induction of hypothyroidism during early postnatal stages triggers a decrease in cognitive performance by decreasing hippocampal synaptic plasticity. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2017, 1863, 870-883.	1.8	28
59	<scp>PPARs</scp> in the central nervous system: roles in neurodegeneration and neuroinflammation. <i>Biological Reviews</i> , 2017, 92, 2046-2069.	4.7	80
60	Brain glucose metabolism: Role of Wnt signaling in the metabolic impairment in Alzheimer's disease. <i>Neuroscience and Biobehavioral Reviews</i> , 2017, 80, 316-328.	2.9	32
61	INT131 increases dendritic arborization and protects against A β toxicity by inducing mitochondrial changes in hippocampal neurons. <i>Biochemical and Biophysical Research Communications</i> , 2017, 490, 955-962.	1.0	6
62	Quercetin Exerts Differential Neuroprotective Effects Against H ₂ O ₂ and A β Aggregates in Hippocampal Neurons: the Role of Mitochondria. <i>Molecular Neurobiology</i> , 2017, 54, 7116-7128.	1.9	56
63	Epigenetic editing of the Dlg4/PSD95 gene improves cognition in aged and Alzheimer's disease mice. <i>Brain</i> , 2017, 140, 3252-3268.	3.7	121
64	Emerging Synaptic Molecules as Candidates in the Etiology of Neurological Disorders. <i>Neural Plasticity</i> , 2017, 2017, 1-25.	1.0	57
65	Wnt/TLR Dialog in Neuroinflammation, Relevance in Alzheimer's Disease. <i>Frontiers in Immunology</i> , 2017, 8, 187.	2.2	39
66	Identification of Cerebral Metal Ion Imbalance in the Brain of Aging Octodon degus. <i>Frontiers in Aging Neuroscience</i> , 2017, 9, 66.	1.7	26
67	Wnt Signaling Prevents the A β Oligomer-Induced Mitochondrial Permeability Transition Pore Opening Preserving Mitochondrial Structure in Hippocampal Neurons. <i>PLoS ONE</i> , 2017, 12, e0168840.	1.1	41
68	MicroRNAs in Metabolic Syndrome. , 2017, , 1-17.		0
69	Wnt5a Increases the Glycolytic Rate and the Activity of the Pentose Phosphate Pathway in Cortical Neurons. <i>Neural Plasticity</i> , 2016, 2016, 1-13.	1.0	10
70	The G β o Activator Mastoparan-7 Promotes Dendritic Spine Formation in Hippocampal Neurons. <i>Neural Plasticity</i> , 2016, 2016, 1-11.	1.0	9
71	Andrographolide recovers cognitive impairment in a natural model of Alzheimer's disease (Octodon) Tj ETQq1 1 0.784314 rgBT /Over 1.5 88		
72	Reduction of Blood Amyloid- β Oligomers in Alzheimer's Disease Transgenic Mice by c-Abl Kinase Inhibition. <i>Journal of Alzheimer's Disease</i> , 2016, 54, 1193-1205.	1.2	23

#	ARTICLE	IF	CITATIONS
73	Wnt-5a/Frizzled9 Receptor Signaling through the $\text{G}\alpha\text{-G}\beta\text{2}\beta\text{3}$ Complex Regulates Dendritic Spine Formation. <i>Journal of Biological Chemistry</i> , 2016, 291, 19092-19107.	1.6	53
74	Activation of Wnt Signaling in Cortical Neurons Enhances Glucose Utilization through Glycolysis. <i>Journal of Biological Chemistry</i> , 2016, 291, 25950-25964.	1.6	46
75	Amyloid- β Peptide Nitrotyrosination Stabilizes Oligomers and Enhances NMDAR-Mediated Toxicity. <i>Journal of Neuroscience</i> , 2016, 36, 11693-11703.	1.7	50
76	Inhibition of Wnt signaling induces amyloidogenic processing of amyloid precursor protein and the production and aggregation of Amyloid- β ($\text{A}\beta$) ₄₂ peptides. <i>Journal of Neurochemistry</i> , 2016, 139, 1175-1191.	2.1	62
77	Frizzled-1 receptor regulates adult hippocampal neurogenesis. <i>Molecular Brain</i> , 2016, 9, 29.	1.3	60
78	On cognitive ecology and the environmental factors that promote Alzheimer disease: lessons from <i>Octodon degus</i> (Rodentia: Octodontidae). <i>Biological Research</i> , 2016, 49, 10.	1.5	25
79	Wnt-5a-regulated miR-101b controls COX2 expression in hippocampal neurons. <i>Biological Research</i> , 2016, 49, 9.	1.5	17
80	Modulation of the NMDA Receptor Through Secreted Soluble Factors. <i>Molecular Neurobiology</i> , 2016, 53, 299-309.	1.9	17
81	Voluntary Running Attenuates Memory Loss, Decreases Neuropathological Changes and Induces Neurogenesis in a Mouse Model of Alzheimer's Disease. <i>Brain Pathology</i> , 2016, 26, 62-74.	2.1	128
82	Are microRNAs the Molecular Link Between Metabolic Syndrome and Alzheimer's Disease?. <i>Molecular Neurobiology</i> , 2016, 53, 2320-2338.	1.9	27
83	Wnt signaling pathway improves central inhibitory synaptic transmission in a mouse model of Duchenne muscular dystrophy. <i>Neurobiology of Disease</i> , 2016, 86, 109-120.	2.1	11
84	Environmental control of microRNAs in the nervous system: Implications in plasticity and behavior. <i>Neuroscience and Biobehavioral Reviews</i> , 2016, 60, 121-138.	2.9	22
85	Role of Wnt Signaling in Central Nervous System Injury. <i>Molecular Neurobiology</i> , 2016, 53, 2297-2311.	1.9	99
86	Recent Advances in Neuroinflammation Therapeutics: PPARs/LXR as Neuroinflammatory Modulators. <i>Current Pharmaceutical Design</i> , 2016, 22, 1312-1323.	0.9	9
87	PSD-95 (Postsynaptic Density Protein-95)., 2016, , 1-7.		0
88	Tetrahydroperforin (IDN5706) targets the endoplasmic reticulum for autophagy activation: potential mechanism for Alzheimer's disease therapy. <i>Neural Regeneration Research</i> , 2016, 11, 242.	1.6	0
89	Is L-methionine a trigger factor for Alzheimer's-like neurodegeneration?: Changes in $\text{A}\beta$ oligomers, tau phosphorylation, synaptic proteins, Wnt signaling and behavioral impairment in wild-type mice. <i>Molecular Neurodegeneration</i> , 2015, 10, 62.	4.4	77
90	How the Wnt signaling pathway protects from neurodegeneration: the mitochondrial scenario. <i>Frontiers in Cellular Neuroscience</i> , 2015, 9, 166.	1.8	61

#	ARTICLE	IF	CITATIONS
91	Wnt-related SynGAP1 is a neuroprotective factor of glutamatergic synapses against A β oligomers. <i>Frontiers in Cellular Neuroscience</i> , 2015, 9, 227.	1.8	10
92	Alzheimer's Disease-Related Protein Expression in the Retina of <i>Octodon degus</i> . <i>PLoS ONE</i> , 2015, 10, e0135499.	1.1	45
93	Andrographolide Stimulates Neurogenesis in the Adult Hippocampus. <i>Neural Plasticity</i> , 2015, 2015, 1-13.	1.0	47
94	β -Catenin-Dependent Signaling Pathway Contributes to Renal Fibrosis in Hypertensive Rats. <i>BioMed Research International</i> , 2015, 2015, 1-13.	0.9	18
95	Andrographolide activates the canonical Wnt signalling pathway by a mechanism that implicates the non-ATP competitive inhibition of GSK-3 β : autoregulation of GSK-3 β <i>in vivo</i> . <i>Biochemical Journal</i> , 2015, 466, 415-430.	1.7	68
96	The ROR2 tyrosine kinase receptor regulates dendritic spine morphogenesis in hippocampal neurons. <i>Molecular and Cellular Neurosciences</i> , 2015, 67, 22-30.	1.0	11
97	Pathogenicity of Lupus Anti-Ribosomal P Antibodies: Role of Cross-Reacting Neuronal Surface P Antigen in Glutamatergic Transmission and Plasticity in a Mouse Model. <i>Arthritis and Rheumatology</i> , 2015, 67, 1598-1610.	2.9	62
98	Teneurinins and Alzheimer's disease: A suggestive role for a unique family of proteins. <i>Medical Hypotheses</i> , 2015, 84, 402-407.	0.8	13
99	A novel function for Wnt signaling modulating neuronal firing activity and the temporal structure of spontaneous oscillation in the entorhinal hippocampal circuit. <i>Experimental Neurology</i> , 2015, 269, 43-55.	2.0	21
100	Accelerating Alzheimer's research through natural animal models. <i>Current Opinion in Psychiatry</i> , 2015, 28, 155-164.	3.1	36
101	The increased potassium intake improves cognitive performance and attenuates histopathological markers in a model of Alzheimer's disease. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015, 1852, 2630-2644.	1.8	26
102	Fructose consumption reduces hippocampal synaptic plasticity underlying cognitive performance. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015, 1852, 2379-2390.	1.8	55
103	Wnt5a inhibits K ⁺ currents in hippocampal synapses through nitric oxide production. <i>Molecular and Cellular Neurosciences</i> , 2015, 68, 314-322.	1.0	15
104	The soluble extracellular fragment of neuroligin-1 targets A β oligomers to the postsynaptic region of excitatory synapses. <i>Biochemical and Biophysical Research Communications</i> , 2015, 466, 66-71.	1.0	23
105	WASP-1, a canonical Wnt signaling potentiator, rescues hippocampal synaptic impairments induced by A β oligomers. <i>Experimental Neurology</i> , 2015, 264, 14-25.	2.0	29
106	Anti-Ribosomal P Protein Autoantibodies From Patients With Neuropsychiatric Lupus Impair Memory in Mice. <i>Arthritis and Rheumatology</i> , 2015, 67, 204-214.	2.9	90
107	Age Progression of Neuropathological Markers in the Brain of the Chilean Rodent <i>Octodon degus</i> , a Natural Model of Alzheimer's Disease. <i>Brain Pathology</i> , 2015, 25, 679-691.	2.1	42
108	The Protein Oxidation Repair Enzyme Methionine Sulfoxide Reductase A Modulates A β Aggregation and Toxicity <i>In Vivo</i> . <i>Antioxidants and Redox Signaling</i> , 2015, 22, 48-62.	2.5	20

#	ARTICLE	IF	CITATIONS
109	Wnt signalling in neuronal differentiation and development. <i>Cell and Tissue Research</i> , 2015, 359, 215-223.	1.5	123
110	Monitoring Mitochondrial Membranes Permeability in Live Neurons and Mitochondrial Swelling Through Electron Microscopy Analysis. <i>Methods in Molecular Biology</i> , 2015, 1254, 87-97.	0.4	13
111	Tetrahydrohyperforin Inhibits the Proteolytic Processing of Amyloid Precursor Protein and Enhances Its Degradation by Atg5-Dependent Autophagy. <i>PLoS ONE</i> , 2015, 10, e0136313.	1.1	34
112	Tetrahydrohyperforin: a neuroprotective modified natural compound against Alzheimer's disease. <i>Neural Regeneration Research</i> , 2015, 10, 552.	1.6	7
113	PSD95 Suppresses Dendritic Arbor Development in Mature Hippocampal Neurons by Occluding the Clustering of NR2B-NMDA Receptors. <i>PLoS ONE</i> , 2014, 9, e94037.	1.1	63
114	Alzheimer's disease: relevant molecular and physiopathological events affecting amyloid- β brain balance and the putative role of PPARs. <i>Frontiers in Aging Neuroscience</i> , 2014, 6, 176.	1.7	46
115	Chronic hypoxia induces the activation of the Wnt/ β -catenin signaling pathway and stimulates hippocampal neurogenesis in wild-type and APP ^{swe-PS1^{E9}} transgenic mice in vivo. <i>Frontiers in Cellular Neuroscience</i> , 2014, 8, 17.	1.8	60
116	Wnt-5a Ligand Modulates Mitochondrial Fission-Fusion in Rat Hippocampal Neurons. <i>Journal of Biological Chemistry</i> , 2014, 289, 36179-36193.	1.6	56
117	Role of Sirt1 During the Ageing Process: Relevance to Protection of Synapses in the Brain. <i>Molecular Neurobiology</i> , 2014, 50, 744-756.	1.9	44
118	Brain metabolite clearance: impact on Alzheimer's disease. <i>Metabolic Brain Disease</i> , 2014, 29, 553-561.	1.4	10
119	Wnt signaling in the nervous system and in Alzheimer's disease. <i>Journal of Molecular Cell Biology</i> , 2014, 6, 64-74.	1.5	260
120	Wnt-5a increases NO and modulates NMDA receptor in rat hippocampal neurons. <i>Biochemical and Biophysical Research Communications</i> , 2014, 444, 189-194.	1.0	39
121	<i>In vivo</i> Activation of Wnt Signaling Pathway Enhances Cognitive Function of Adult Mice and Reverses Cognitive Deficits in an Alzheimer's Disease Model. <i>Journal of Neuroscience</i> , 2014, 34, 2191-2202.	1.7	125
122	Wnt Signaling in Skeletal Muscle Dynamics: Myogenesis, Neuromuscular Synapse and Fibrosis. <i>Molecular Neurobiology</i> , 2014, 49, 574-589.	1.9	107
123	Phosphorylated tau potentiates β -induced mitochondrial damage in mature neurons. <i>Neurobiology of Disease</i> , 2014, 71, 260-269.	2.1	55
124	Is Alzheimer's disease related to metabolic syndrome? A Wnt signaling conundrum. <i>Progress in Neurobiology</i> , 2014, 121, 125-146.	2.8	87
125	Signaling pathway cross talk in Alzheimer's disease. <i>Cell Communication and Signaling</i> , 2014, 12, 23.	2.7	126
126	Synthesis and Multitarget Biological Profiling of a Novel Family of Rhein Derivatives As Disease-Modifying Anti-Alzheimer Agents. <i>Journal of Medicinal Chemistry</i> , 2014, 57, 2549-2567.	2.9	132

#	ARTICLE	IF	CITATIONS
127	Andrographolide reduces cognitive impairment in young and mature A β 2PPswe/PS-1 mice. <i>Molecular Neurodegeneration</i> , 2014, 9, 61.	4.4	95
128	EphA4 Activation of c-Abl Mediates Synaptic Loss and LTP Blockade Caused by Amyloid- β Oligomers. <i>PLoS ONE</i> , 2014, 9, e92309.	1.1	75
129	Metal and complementary molecular bioimaging in Alzheimer's disease. <i>Frontiers in Aging Neuroscience</i> , 2014, 6, 138.	1.7	44
130	Nicotine Prevents Synaptic Impairment Induced by Amyloid- β Oligomers Through α 7-Nicotinic Acetylcholine Receptor Activation. <i>NeuroMolecular Medicine</i> , 2013, 15, 549-569.	1.8	77
131	Tetrahydrohyperforin Induces Mitochondrial Dynamics and Prevents Mitochondrial Ca ²⁺ Overload after A β ² and A β ² -AChE Complex Challenge in Rat Hippocampal Neurons. <i>Journal of Alzheimer's Disease</i> , 2013, 37, 735-746.	1.2	12
132	Tetrahydrohyperforin Decreases Cholinergic Markers associated with Amyloid- β Plaques, 4-Hydroxynonenal Formation, and Caspase-3 Activation in A β ² PP/PS1 Mice. <i>Journal of Alzheimer's Disease</i> , 2013, 36, 99-118.	1.2	26
133	Peroxisome Proliferators Reduce Spatial Memory Impairment, Synaptic Failure, and Neurodegeneration in Brains of a Double Transgenic Mice Model of Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2013, 33, 941-959.	1.2	49
134	Tetrahydrohyperforin Increases Adult Hippocampal Neurogenesis in Wild-Type and APPswe/PS1 Δ E9 Mice. <i>Journal of Alzheimer's Disease</i> , 2013, 34, 873-885.	1.2	34
135	Peroxisome Proliferator-Activated Receptor (PPAR) γ and PPAR α Agonists Modulate Mitochondrial Fusion-Fission Dynamics: Relevance to Reactive Oxygen Species (ROS)-Related Neurodegenerative Disorders?. <i>PLoS ONE</i> , 2013, 8, e64019.	1.1	84
136	Wnt signaling: Role in LTP, neural networks and memory. <i>Ageing Research Reviews</i> , 2013, 12, 786-800.	5.0	76
137	Peroxisome Proliferator-activated Receptors and Alzheimer's Disease: Hitting the Blood-Brain Barrier. <i>Molecular Neurobiology</i> , 2013, 48, 438-451.	1.9	36
138	Wnt signaling in the regulation of adult hippocampal neurogenesis. <i>Frontiers in Cellular Neuroscience</i> , 2013, 7, 100.	1.8	151
139	Frizzled-5 Receptor Is Involved in Neuronal Polarity and Morphogenesis of Hippocampal Neurons. <i>PLoS ONE</i> , 2013, 8, e78892.	1.1	32
140	Canonical Wnt signaling protects hippocampal neurons from A β ² oligomers: role of non-canonical Wnt-5a/Ca ²⁺ in mitochondrial dynamics. <i>Frontiers in Cellular Neuroscience</i> , 2013, 7, 97.	1.8	77
141	WNT signaling in neuronal maturation and synaptogenesis. <i>Frontiers in Cellular Neuroscience</i> , 2013, 7, 103.	1.8	204
142	Wnts in adult brain: from synaptic plasticity to cognitive deficiencies. <i>Frontiers in Cellular Neuroscience</i> , 2013, 7, 224.	1.8	128
143	ATP Induces NO Production in Hippocampal Neurons by P2X7 Receptor Activation Independent of Glutamate Signaling. <i>PLoS ONE</i> , 2013, 8, e57626.	1.1	27
144	Thiazolidinediones Promote Axonal Growth through the Activation of the JNK Pathway. <i>PLoS ONE</i> , 2013, 8, e65140.	1.1	24

#	ARTICLE	IF	CITATIONS
145	Postsynaptic Receptors for Amyloid- β^2 Oligomers as Mediators of Neuronal Damage in Alzheimer's Disease. <i>Frontiers in Physiology</i> , 2012, 3, 464.	1.3	84
146	Wnt Signaling: Role in Alzheimer Disease and Schizophrenia. <i>Journal of NeuroImmune Pharmacology</i> , 2012, 7, 788-807.	2.1	165
147	Postsynaptic dysfunction is associated with spatial and object recognition memory loss in a natural model of Alzheimer's disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 13835-13840.	3.3	113
148	Wnt-5a Is a Synaptogenic Factor with Neuroprotective Properties against $A\beta^2$ Toxicity. <i>Neurodegenerative Diseases</i> , 2012, 10, 23-26.	0.8	30
149	Frizzled receptors in neurons: From growth cones to the synapse. <i>Cytoskeleton</i> , 2012, 69, 528-534.	1.0	25
150	Reusing and composing models of cell fate regulation of human bone precursor cells. <i>BioSystems</i> , 2012, 108, 63-72.	0.9	5
151	Recent rodent models for Alzheimer's disease: clinical implications and basic research. <i>Journal of Neural Transmission</i> , 2012, 119, 173-195.	1.4	97
152	SIRT1 Regulates Dendritic Development in Hippocampal Neurons. <i>PLoS ONE</i> , 2012, 7, e47073.	1.1	68
153	Regulation of NMDA-Receptor Synaptic Transmission by Wnt Signaling. <i>Journal of Neuroscience</i> , 2011, 31, 9466-9471.	1.7	136
154	The GABA(A) β -receptors in hippocampal spontaneous activity and their distribution in hippocampus, amygdala and visual cortex. <i>Neuroscience Letters</i> , 2011, 500, 20-25.	1.0	18
155	Interactions of AChE with A β Aggregates in Alzheimer's Brain: Therapeutic Relevance of IDN 5706. <i>Frontiers in Molecular Neuroscience</i> , 2011, 4, 19.	1.4	132
156	The Cellular Prion Protein Prevents Copper-Induced Inhibition of P2 _X ₄ International Journal of Alzheimer's Disease, 2011, 2011, 1-6.	1.1	4
157	The Synaptic Protein Neuroligin-1 Interacts with the Amyloid β^2 -Peptide. Is There a Role in Alzheimer's Disease?. <i>Biochemistry</i> , 2011, 50, 8127-8137.	1.2	49
158	Copper Reduces A β Oligomeric Species and Ameliorates Neuromuscular Synaptic Defects in a <i>C. elegans</i> Model of Inclusion Body Myositis. <i>Journal of Neuroscience</i> , 2011, 31, 10149-10158.	1.7	39
159	Wnt signaling modulates pre- and postsynaptic maturation: Therapeutic considerations. <i>Developmental Dynamics</i> , 2010, 239, 94-101.	0.8	30
160	Genome-wide identification of new Wnt/ β -catenin target genes in the human genome using CART method. <i>BMC Genomics</i> , 2010, 11, 348.	1.2	50
161	Adult hippocampal neurogenesis in aging and Alzheimer's disease. <i>Birth Defects Research Part C: Embryo Today Reviews</i> , 2010, 90, 284-296.	3.6	49
162	Wnt-5a occludes $A\beta^2$ oligomer-induced depression of glutamatergic transmission in hippocampal neurons. <i>Molecular Neurodegeneration</i> , 2010, 5, 3.	4.4	107

#	ARTICLE	IF	CITATIONS
163	Amyloid- β -Acetylcholinesterase complexes potentiate neurodegenerative changes induced by the A β ² peptide. Implications for the pathogenesis of Alzheimer's disease. <i>Molecular Neurodegeneration</i> , 2010, 5, 4.	4.4	96
164	Emerging roles of Wnts in the adult nervous system. <i>Nature Reviews Neuroscience</i> , 2010, 11, 77-86.	4.9	558
165	β -Amyloid Causes Depletion of Synaptic Vesicles Leading to Neurotransmission Failure. <i>Journal of Biological Chemistry</i> , 2010, 285, 2506-2514.	1.6	153
166	Wingless-type family member 5A (Wnt-5a) stimulates synaptic differentiation and function of glutamatergic synapses. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 21164-21169.	3.3	185
167	The Hyperforin Derivative IDN5706 Occludes Spatial Memory Impairments and Neuropathological Changes in a Double Transgenic Alzheimers Mouse Model. <i>Current Alzheimer Research</i> , 2010, 7, 126-133.	0.7	38
168	Synaptic Clustering of PSD-95 Is Regulated by c-Abl through Tyrosine Phosphorylation. <i>Journal of Neuroscience</i> , 2010, 30, 3728-3738.	1.7	50
169	Wnt-5a Modulates Recycling of Functional GABAA Receptors on Hippocampal Neurons. <i>Journal of Neuroscience</i> , 2010, 30, 8411-8420.	1.7	112
170	Wnt-5a/JNK Signaling Promotes the Clustering of PSD-95 in Hippocampal Neurons. <i>Journal of Biological Chemistry</i> , 2009, 284, 15857-15866.	1.6	187
171	c-Abl modulates AICD dependent cellular responses: Transcriptional induction and apoptosis. <i>Journal of Cellular Physiology</i> , 2009, 220, 136-143.	2.0	38
172	Calcium/calmodulin-dependent protein kinase type IV is a target gene of the Wnt/ β -catenin signaling pathway. <i>Journal of Cellular Physiology</i> , 2009, 221, 658-667.	2.0	71
173	Role of the Wnt receptor Frizzled-1 in presynaptic differentiation and function. <i>Neural Development</i> , 2009, 4, 41.	1.1	95
174	Intracellular amyloid formation in muscle cells of A β ² -transgenic <i>Caenorhabditis elegans</i> : determinants and physiological role in copper detoxification. <i>Molecular Neurodegeneration</i> , 2009, 4, 2.	4.4	39
175	Methionine sulfoxide reductase A expression is regulated by the DAF-16/FOXO pathway in <i>Caenorhabditis elegans</i> . <i>Aging Cell</i> , 2009, 8, 690-705.	3.0	70
176	Overexpression of amyloid precursor protein increases copper content in HEK293 cells. <i>Biochemical and Biophysical Research Communications</i> , 2009, 382, 740-744.	1.0	15
177	The role of Wnt signaling in neuroprotection. <i>Drug News and Perspectives</i> , 2009, 22, 579.	1.9	30
178	Inclusion Body Myositis: A View from the <i>Caenorhabditis elegans</i> Muscle. <i>Molecular Neurobiology</i> , 2008, 38, 178-198.	1.9	15
179	The role of Wnt signaling in neuronal dysfunction in Alzheimer's Disease. <i>Molecular Neurodegeneration</i> , 2008, 3, 9.	4.4	164
180	Heparin activates Wnt signaling for neuronal morphogenesis. <i>Journal of Cellular Physiology</i> , 2008, 216, 805-815.	2.0	34

#	ARTICLE	IF	CITATIONS
181	Frizzled β 1 is involved in the neuroprotective effect of Wnt3a against A β 2 oligomers. <i>Journal of Cellular Physiology</i> , 2008, 217, 215-227.	2.0	80
182	Amyloid β cholinesterase interactions. <i>FEBS Journal</i> , 2008, 275, 625-632.	2.2	215
183	Release of acetylcholinesterase (AChE) from β 2-amyloid plaques assemblies improves the spatial memory impairments in APP-transgenic mice. <i>Chemico-Biological Interactions</i> , 2008, 175, 142-149.	1.7	37
184	β 2-Amyloid Oligomers Affect the Structure and Function of the Postsynaptic Region: Role of the β 2-Wnt Signaling Pathway. <i>Neurodegenerative Diseases</i> , 2008, 5, 149-152.	0.8	31
185	Wnt signaling in neuroprotection and stem cell differentiation. <i>Progress in Neurobiology</i> , 2008, 86, 281-296.	2.8	182
186	STI571 prevents apoptosis, tau phosphorylation and behavioural impairments induced by Alzheimer's β 2-amyloid deposits. <i>Brain</i> , 2008, 131, 2425-2442.	3.7	136
187	Wnt-7a Modulates the Synaptic Vesicle Cycle and Synaptic Transmission in Hippocampal Neurons. <i>Journal of Biological Chemistry</i> , 2008, 283, 5918-5927.	1.6	205
188	Structure-Function Implications in Alzheimers Disease: Effect of A β 42 Oligomers at Central Synapses. <i>Current Alzheimer Research</i> , 2008, 5, 233-243.	0.7	91
189	Expression of Wnt Receptors, Frizzled, in Rat Neuronal Cells. , 2008, , 317-324.		0
190	Wnt-7a Induces Presynaptic Colocalization of α 7-Nicotinic Acetylcholine Receptors and Adenomatous Polyposis Coli in Hippocampal Neurons. <i>Journal of Neuroscience</i> , 2007, 27, 5313-5325.	1.7	101
191	Peroxisome Proliferator-activated Receptor β 3 Up-regulates the Bcl-2 Anti-apoptotic Protein in Neurons and Induces Mitochondrial Stabilization and Protection against Oxidative Stress and Apoptosis. <i>Journal of Biological Chemistry</i> , 2007, 282, 37006-37015.	1.6	223
192	ApoER2 expression increases A β 2 production while decreasing Amyloid Precursor Protein (APP) endocytosis: Possible role in the partitioning of APP into lipid rafts and in the regulation of β 3-secretase activity. <i>Molecular Neurodegeneration</i> , 2007, 2, 14.	4.4	66
193	Synaptotoxicity in Alzheimer's Disease: The Wnt Signaling Pathway as a Molecular Target. <i>IUBMB Life</i> , 2007, 59, 316-321.	1.5	58
194	The functional links between prion protein and copper. <i>Biological Research</i> , 2006, 39, 39-44.	1.5	20
195	Induction of cellular prion protein gene expression by copper in neurons. <i>American Journal of Physiology - Cell Physiology</i> , 2006, 290, C271-C281.	2.1	58
196	Role of Copper in Prion Diseases: Deleterious or Beneficial?. <i>Current Pharmaceutical Design</i> , 2006, 12, 2587-2595.	0.9	18
197	Copper brain homeostasis: Role of amyloid precursor protein and prion protein. <i>IUBMB Life</i> , 2005, 57, 645-650.	1.5	23
198	ApoER2 is Endocytosed by a Clathrin-Mediated Process Involving the Adaptor Protein Dab2 Independent of its Rafts' Association. <i>Traffic</i> , 2005, 6, 820-838.	1.3	64

#	ARTICLE	IF	CITATIONS
199	Acetylcholinesterase-Amyloid- β -peptide Interaction: Effect of Congo Red and the Role of the Wnt Pathway. <i>Current Alzheimer Research</i> , 2005, 2, 301-306.	0.7	39
200	Peroxisomal Proliferation Protects from β -Amyloid Neurodegeneration. <i>Journal of Biological Chemistry</i> , 2005, 280, 41057-41068.	1.6	137
201	Blood Cells Cholinesterase Activity in Early Stage Alzheimer's Disease and Vascular Dementia. <i>Dementia and Geriatric Cognitive Disorders</i> , 2005, 19, 204-212.	0.7	29
202	Peroxisome proliferator-activated receptor δ is expressed in hippocampal neurons and its activation prevents β -amyloid neurodegeneration: role of Wnt signaling. <i>Experimental Cell Research</i> , 2005, 304, 91-104.	1.2	181
203	Trolox and 17 β -Estradiol Protect against Amyloid β -Peptide Neurotoxicity by a Mechanism That Involves Modulation of the Wnt Signaling Pathway. <i>Journal of Biological Chemistry</i> , 2005, 280, 11615-11625.	1.6	109
204	The anti-inflammatory and cholinesterase inhibitor bifunctional compound IBU-PO protects from β -amyloid neurotoxicity by acting on Wnt signaling components. <i>Neurobiology of Disease</i> , 2005, 18, 176-183.	2.1	35
205	Is there a role for copper in neurodegenerative diseases?. <i>Molecular Aspects of Medicine</i> , 2005, 26, 405-420.	2.7	65
206	Human-like rodent amyloid- β -peptide determines Alzheimer pathology in aged wild-type Octodon degu. <i>Neurobiology of Aging</i> , 2005, 26, 1023-1028.	1.5	106
207	Acetylcholinesterase Interaction with Alzheimer Amyloid β . , 2005, 38, 299-317.		86
208	An Overview of the Current and Novel Drugs for Alzheimers Disease with Particular Reference to Anti-Cholinesterase Compounds. <i>Current Pharmaceutical Design</i> , 2004, 10, 3121-3130.	0.9	75
209	The N-terminal copper-binding domain of the amyloid precursor protein protects against Cu ²⁺ neurotoxicity in vivo. <i>FASEB Journal</i> , 2004, 18, 1701-1703.	0.2	40
210	Stepwise construction of triple-helical heparin binding sites using peptide models. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2004, 1698, 187-195.	1.1	10
211	Heparan sulfate proteoglycans are increased during skeletal muscle regeneration: requirement of syndecan-3 for successful fiber formation. <i>Journal of Cell Science</i> , 2004, 117, 73-84.	1.2	112
212	Signal transduction during amyloid- β -peptide neurotoxicity: role in Alzheimer disease. <i>Brain Research Reviews</i> , 2004, 47, 275-289.	9.1	121
213	Structural and functional organization of synaptic acetylcholinesterase. <i>Brain Research Reviews</i> , 2004, 47, 96-104.	9.1	41
214	Wnt-3a overcomes β -amyloid toxicity in rat hippocampal neurons. <i>Experimental Cell Research</i> , 2004, 297, 186-196.	1.2	203
215	Structure and function of amyloid in Alzheimer's disease. <i>Progress in Neurobiology</i> , 2004, 74, 323-349.	2.8	126
216	Acetylcholinesterase- $A\beta$ Complexes Are More Toxic than $A\beta$ Fibrils in Rat Hippocampus. <i>American Journal of Pathology</i> , 2004, 164, 2163-2174.	1.9	128

#	ARTICLE	IF	CITATIONS
217	M1 muscarinic receptor activation protects neurons from β^2 -amyloid toxicity. A role for Wnt signaling pathway. <i>Neurobiology of Disease</i> , 2004, 17, 337-348.	2.1	71
218	Acetylcholinesterase (AChE) - Amyloid- β ; Peptide Complexes in Alzheimers Disease. The Wnt Signaling Pathway. <i>Current Alzheimer Research</i> , 2004, 1, 249-254.	0.7	51
219	Copper reduction by copper binding proteins and its relation to neurodegenerative diseases. <i>BioMetals</i> , 2003, 16, 91-98.	1.8	73
220	PPAR γ activators induce growth arrest and process extension in B12 oligodendrocyte-like cells and terminal differentiation of cultured oligodendrocytes. <i>Journal of Neuroscience Research</i> , 2003, 72, 425-435.	1.3	69
221	Ethanol specifically decreases peroxisome proliferator activated receptor β^2 in B12 oligodendrocyte-like cells. <i>Journal of Neurochemistry</i> , 2003, 85, 135-141.	2.1	8
222	The human prion octarepeat fragment prevents and reverses the inhibitory action of copper in the P2X ₄ receptor without modifying the zinc action. <i>Journal of Neurochemistry</i> , 2003, 85, 709-716.	2.1	5
223	Acetylcholinesterase induces neuronal cell loss, astrocyte hypertrophy and behavioral deficits in mammalian hippocampus. <i>Journal of Neurochemistry</i> , 2003, 87, 195-204.	2.1	49
224	Acetylcholinesterase induces the expression of the β^2 -amyloid precursor protein in glia and activates glial cells in culture. <i>Neurobiology of Disease</i> , 2003, 14, 447-457.	2.1	26
225	Two Different Heparin-binding Domains in the Triple-helical Domain of ColQ, the Collagen Tail Subunit of Synaptic Acetylcholinesterase. <i>Journal of Biological Chemistry</i> , 2003, 278, 23233-23242.	1.6	46
226	Metalloenzyme-like Activity of Alzheimer's Disease β^2 -Amyloid. <i>Journal of Biological Chemistry</i> , 2002, 277, 40302-40308.	1.6	536
227	Protein kinase C inhibits amyloid β^2 -peptide neurotoxicity by acting on members of the Wnt pathway. <i>FASEB Journal</i> , 2002, 16, 1982-1984.	0.2	156
228	Laminin affects polymerization, depolymerization and neurotoxicity of β^2 peptide. <i>Peptides</i> , 2002, 23, 1229-1240.	1.2	40
229	Wnt signaling involvement in β^2 -amyloid-dependent neurodegeneration. <i>Neurochemistry International</i> , 2002, 41, 341-344.	1.9	80
230	Vitamin E But Not 17 β -Estradiol Protects against Vascular Toxicity Induced by β^2 -Amyloid Wild Type and the Dutch Amyloid Variant. <i>Journal of Neuroscience</i> , 2002, 22, 3081-3089.	1.7	51
231	A Structural Motif of Acetylcholinesterase That Promotes Amyloid β^2 -Peptide Fibril Formation. <i>Biochemistry</i> , 2001, 40, 10447-10457.	1.2	385
232	Cysteine 144 Is a Key Residue in the Copper Reduction by the β^2 -Amyloid Precursor Protein. <i>Journal of Neurochemistry</i> , 2001, 73, 1288-1292.	2.1	51
233	Interaction of the collagen-like tail of asymmetric acetylcholinesterase with heparin depends on triple-helical conformation, sequence and stability. <i>Biochemical Journal</i> , 2000, 350, 283.	1.7	13
234	Interaction of the collagen-like tail of asymmetric acetylcholinesterase with heparin depends on triple-helical conformation, sequence and stability. <i>Biochemical Journal</i> , 2000, 350, 283-290.	1.7	31

#	ARTICLE	IF	CITATIONS
235	Molecular modeling of the collagen-like tail of asymmetric acetylcholinesterase. <i>Protein Engineering, Design and Selection</i> , 2000, 13, 27-34.	1.0	9
236	Expression of β 2-macroglobulin receptor/low density lipoprotein receptor-related protein (LRP) in rat microglial cells. <i>Journal of Neuroscience Research</i> , 2000, 60, 401-411.	1.3	83
237	The N-Terminal Tandem Repeat Region of Human Prion Protein Reduces Copper: Role of Tryptophan Residues. <i>Biochemical and Biophysical Research Communications</i> , 2000, 269, 491-495.	1.0	80
238	Wnt signaling function in Alzheimer's disease. <i>Brain Research Reviews</i> , 2000, 33, 1-12.	9.1	275
239	The role of oxidative stress in the toxicity induced by amyloid β -peptide in Alzheimer's disease. <i>Progress in Neurobiology</i> , 2000, 62, 633-648.	2.8	347
240	Interaction of Collagen-Like Peptide Models of Asymmetric Acetylcholinesterase with Glycosaminoglycans: Spectroscopic Studies of Conformational Changes and Stability. <i>Biochemistry</i> , 2000, 39, 14884-14892.	1.2	15
241	Amyloid- β -peptide reduces copper(II) to copper(I) independent of its aggregation state. <i>Biological Research</i> , 2000, 33, 125-31.	1.5	33
242	Molecular modeling of the amyloid- β -peptide using the homology to a fragment of triosephosphate isomerase that forms amyloid in vitro. <i>Protein Engineering, Design and Selection</i> , 1999, 12, 959-966.	1.0	12
243	PC12 and neuro 2a cells have different susceptibilities to acetylcholinesterase-amyloid complexes, amyloid β 25-35 fragment, glutamate, and hydrogen peroxide. <i>Journal of Neuroscience Research</i> , 1999, 56, 620-631.	1.3	61
244	Mannose receptor is present in a functional state in rat microglial cells. <i>Journal of Neuroscience Research</i> , 1999, 58, 387-395.	1.3	54
245	Neurotoxicity of acetylcholinesterase amyloid β -peptide aggregates is dependent on the type of β -peptide and the AChE concentration present in the complexes. <i>FEBS Letters</i> , 1999, 450, 205-209.	1.3	80
246	Peripheral binding site is involved in the neurotrophic activity of acetylcholinesterase. <i>NeuroReport</i> , 1999, 10, 3621-3625.	0.6	67
247	Brain acetylcholinesterase promotes amyloid-beta-peptide aggregation but does not hydrolyze amyloid precursor protein peptides. <i>Neurochemical Research</i> , 1998, 23, 135-140.	1.6	38
248	Molecular interactions of acetylcholinesterase with senile plaques. <i>Journal of Physiology (Paris)</i> , 1998, 92, 341-344.	2.1	52
249	Responses induced by tacrine in neuronal and non-neuronal cell lines. , 1998, 52, 435-444.		12
250	Cellular and molecular basis of estrogen's neuroprotection. <i>Molecular Neurobiology</i> , 1998, 17, 73-86.	1.9	109
251	Crosslinking of amyloid- β peptide to brain acetylcholinesterase. <i>Molecular and Chemical Neuropathology</i> , 1998, 33, 39-49.	1.0	10
252	Developmental regulation of mouse brain monomeric acetylcholinesterase. <i>International Journal of Developmental Neuroscience</i> , 1998, 16, 123-134.	0.7	14

#	ARTICLE	IF	CITATIONS
253	Estrogen protects neuronal cells from the cytotoxicity induced by acetylcholinesterase-amyloid complexes. <i>FEBS Letters</i> , 1998, 441, 220-224.	1.3	72
254	Laminin blocks the assembly of wild-type A β and the Dutch variant peptide into Alzheimer's fibrils. <i>Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the International Society of Amyloidosis</i> , 1998, 5, 16-23.	1.4	34
255	At Least Two Receptors of Asymmetric Acetylcholinesterase Are Present at the Synaptic Basal Lamina of Torpedo Electric Organ. <i>Biochemical and Biophysical Research Communications</i> , 1998, 250, 312-317.	1.0	12
256	A Major Portion of Synaptic Basal Lamina Acetylcholinesterase Is Detached by High Salt- and Heparin-containing Buffers from Rat Diaphragm Muscle and Torpedo Electric Organ. <i>Journal of Biological Chemistry</i> , 1998, 273, 4258-4265.	1.6	16
257	Stable Complexes Involving Acetylcholinesterase and Amyloid- β Peptide Change the Biochemical Properties of the Enzyme and Increase the Neurotoxicity of Alzheimer's Fibrils. <i>Journal of Neuroscience</i> , 1998, 18, 3213-3223.	1.7	264
258	Molecular Interactions of Acetylcholinesterase with the Synaptic Basal Lamina and the Senile Plaques. , 1998, , 167-173.		0
259	Identification of an Acetylcholinesterase Fragment that Promotes Alzheimer β -Amyloid Fibril Formation. , 1998, , 185-186.		1
260	The Heparin-Binding Sites in the Collagenic Tail of Acetylcholinesterase. , 1998, , 444-445.		0
261	Acetylcholinesterase Enhances the Neurotoxicity of β -Amyloid Fibrils. , 1998, , 182-182.		0
262	A Monoclonal Antibody against Acetylcholinesterase Inhibits the Formation of Amyloid Fibrils Induced by the Enzyme. <i>Biochemical and Biophysical Research Communications</i> , 1997, 232, 652-655.	1.0	102
263	Acetylcholinesterase promotes the aggregation of amyloid- β -peptide fragments by forming a complex with the growing fibrils 1 Edited by A. R. Fersht. <i>Journal of Molecular Biology</i> , 1997, 272, 348-361.	2.0	274
264	Amyloid Precursor Protein Fragment and Acetylcholinesterase Increase with Cell Confluence and Differentiation in a Neuronal Cell Line. <i>Experimental Cell Research</i> , 1996, 229, 93-99.	1.2	22
265	Laminin inhibits amyloid- β -peptide fibrillation. <i>Neuroscience Letters</i> , 1996, 218, 201-203.	1.0	70
266	Acetylcholinesterase Accelerates Assembly of Amyloid- β -Peptides into Alzheimer's Fibrils: Possible Role of the Peripheral Site of the Enzyme. <i>Neuron</i> , 1996, 16, 881-891.	3.8	1,032
267	Sprouting and abnormal contacts of nonmyelinated axons, and deposition of extracellular material induced by the amyloid precursor protein (APP) and other protease inhibitors. <i>Brain Research</i> , 1996, 718, 13-24.	1.1	25
268	Extracellular matrix regulates the amount of the β -amyloid precursor protein and its amyloidogenic fragments. , 1996, 166, 360-369.		29
269	Abnormal development of the locomotor activity in yellow larvae of <i>Drosophila</i> : a cuticular defect?. <i>Genetica</i> , 1996, 97, 205-210.	0.5	17
270	Tetrameric (G ₄) Acetylcholinesterase: Structure, Localization, and Physiological Regulation. <i>Journal of Neurochemistry</i> , 1996, 66, 1335-1346.	2.1	60

#	ARTICLE	IF	CITATIONS
271	The α -Helical to β -Strand Transition in the Amino-terminal Fragment of the Amyloid β -Peptide Modulates Amyloid Formation. <i>Journal of Biological Chemistry</i> , 1995, 270, 3063-3067.	1.6	298
272	Mitosis of Schwann Cells and Demyelination are Induced by the Amyloid Precursor Protein and Other Protease Inhibitors in the Rat Sciatic Nerve. <i>European Journal of Neuroscience</i> , 1995, 7, 152-159.	1.2	31
273	Effect of protamine on the solubilization of collagen-tailed acetylcholinesterase: potential heparin-binding consensus sequences in the tail of the enzyme. <i>BBA - Proteins and Proteomics</i> , 1995, 1252, 53-58.	2.1	8
274	Two Heparin-binding Domains Are Present on the Collagenic Tail of Asymmetric Acetylcholinesterase. <i>Journal of Biological Chemistry</i> , 1995, 270, 11043-11046.	1.6	73
275	Acetylcholinesterase, a senile plaque component, affects the fibrillogenesis of amyloid- β -peptides. <i>Neuroscience Letters</i> , 1995, 201, 49-52.	1.0	62
276	Sensitivity of acetylcholinesterase molecular forms to inhibition by high MgCl ₂ concentration. <i>BBA - Proteins and Proteomics</i> , 1994, 1208, 286-293.	2.1	8
277	Metamorphosis of laboratory-reared larvae of <i>Concholepas concholepas</i> (Mollusca; Gastropoda). <i>Aquaculture</i> , 1994, 126, 299-303.	1.7	8
278	Monomeric amphiphilic forms of acetylcholinesterase appear early during brain development and may correspond to biosynthetic precursors of the amphiphilic G4 forms. <i>Neuroscience Letters</i> , 1994, 173, 155-158.	1.0	16
279	Structural Determinants of the Alzheimer's Amyloid β -Peptide. <i>Journal of Neurochemistry</i> , 1994, 63, 1191-1198.	2.1	141
280	Acetylcholinesterase changes in hearts with sinus rhythm and atrial fibrillation. <i>General Pharmacology</i> , 1993, 24, 111-114.	0.7	9
281	Extracellular matrix components and amyloid in neuritic plaques of Alzheimer's disease. <i>General Pharmacology</i> , 1993, 24, 1063-1068.	0.7	31
282	Presence of an heparin-binding growth factor in <i>Concholepas concholepas</i> Bruguiere (Mollusca; Tj ETQq0 0 0 rgBT/Overlock, 10 Tf 50 3	0.7	7
283	Molecular changes induced by metamorphosis in larvae of the prosobranch <i>Concholepas concholepas</i> Bruguiere (Mollusca; Gastropoda; Muricidae). <i>Journal of Experimental Marine Biology and Ecology</i> , 1993, 168, 205-215.	0.7	7
284	A neurofilament polypeptide and the glial fibrillary acidic protein share common epitopes in the variable region. <i>Neuroscience Letters</i> , 1993, 161, 137-140.	1.0	3
285	Nerve regeneration is improved by insulin-like growth factor I (IGF-I) and basic fibroblast growth factor (bFGF). <i>Restorative Neurology and Neuroscience</i> , 1993, 5, 181-189.	0.4	15
286	Axonal sprouting induced in the sciatic nerve by the amyloid precursor protein (APP) and other antiproteases. <i>Neuroscience Letters</i> , 1992, 144, 130-134.	1.0	58
287	Amphiphilic behavior of a brain tetrameric acetylcholinesterase form lacking the plasma membrane anchoring domain. <i>Brain Research</i> , 1992, 580, 1-5.	1.1	8
288	Isolation of proteoglycans synthesized by rat heart: Evidence for the presence of several distinct forms. <i>General Pharmacology</i> , 1992, 23, 249-255.	0.7	3

#	ARTICLE	IF	CITATIONS
289	Sulfation is required for mobility of veliger larvae of <i>Concholepas concholepas</i> (Mollusca; Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50	1.4	5
290	A high molecular weight proteoglycan is differentially expressed during development of the mollusc <i>Concholepas concholepas</i> (Mollusca; Gastropoda; Muricidae). <i>The Journal of Experimental Zoology</i> , 1992, 264, 363-371.	1.4	5
291	Binding of Asymmetric (A2) Acetylcholinesterase to C2 Muscle Cells and to Cho Mutants Defective in Glycosaminoglycan Synthesis. , 1992, , 25-32.		3
292	A comparison of the <i>Xenopus laevis</i> oocyte acetylcholinesterase with the muscle and brain enzyme suggests variations at the post-translational level. <i>Comparative Biochemistry and Physiology Part C: Comparative Pharmacology</i> , 1991, 98, 299-305.	0.2	3
293	Increase of macromolecule synthesis after hatching of <i>Concholepas concholepas</i> veliger larvae: Effect of sulfate in the synthesis of proteoglycans. <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1990, 96, 613-619.	0.2	0
294	Proteoglycan production in <i>Drosophila</i> egg development: Effect of Î ² -d-xyloside on proteoglycan synthesis and larvae motility. <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1990, 97, 307-314.	0.2	9
295	Carrageenans solubilize asymmetric acetylcholinesterase from nicotinic cholinergic synapses. <i>Comparative Biochemistry and Physiology Part C: Comparative Pharmacology</i> , 1990, 96, 77-81.	0.2	0
296	Association of Acetylcholinesterase with the cell surface. <i>Journal of Membrane Biology</i> , 1990, 118, 1-9.	1.0	41
297	Neurotransmitter-related enzyme acetylcholinesterase in juveniles of <i>Concholepas concholepas</i> (Mollusca; gastropoda; muricidae). <i>The Journal of Experimental Zoology</i> , 1990, 255, 1-8.	1.4	6
298	Dermatan sulfate and de-sulfated heparin solubilized collagen-tailed acetylcholinesterase from the rat neuromuscular junction. <i>Brain Research</i> , 1990, 529, 91-95.	1.1	8
299	A simple assay to estimate the acetylcholinesterase molecular forms in crude extracts of rat skeletal muscle. <i>Analytical Biochemistry</i> , 1989, 180, 227-230.	1.1	3
300	Distribution and anchoring of molecular forms of acetylcholinesterase. <i>Trends in Pharmacological Sciences</i> , 1989, 10, 325-329.	4.0	56
301	Axons grow in the aging rat but fast transport and acetylcholinesterase content remain unchanged. <i>Brain Research</i> , 1988, 441, 331-338.	1.1	21
302	A membrane-associated dimer of acetylcholinesterase from <i>Xenopus</i> skeletal muscle is solubilized by phosphatidylinositol-specific phospholipase C. <i>Neuroscience Letters</i> , 1988, 90, 186-190.	1.0	29
303	Changes in contralateral synaptic acetylcholinesterase following motor nerve section in rats. <i>Neuroscience Letters</i> , 1988, 90, 229-233.	1.0	5
304	The sensory projections of <i>Drosophila</i> mutants which show abnormal wing formation or flying behavior. <i>Brain Research</i> , 1987, 416, 248-256.	1.1	4
305	Co-solubilization of asymmetric acetylcholinesterase and dermatan sulfate proteoglycan from the extracellular matrix of rat skeletal muscles. <i>FEBS Letters</i> , 1987, 213, 159-163.	1.3	23
306	Biosynthesis of the neurofilament heavy subunit in <i>Xenopus</i> oocytes microinjected with rat brain poly(A)+ RNA. <i>Molecular Biology Reports</i> , 1987, 12, 265-271.	1.0	5

#	ARTICLE	IF	CITATIONS
307	Isolation of the heparan sulfate proteoglycans from the extracellular matrix of rat skeletal muscle. <i>Journal of Neurobiology</i> , 1987, 18, 271-282.	3.7	46
308	Interaction of heparin with multimolecular aggregates of acetylcholinesterase. <i>Cellular and Molecular Neurobiology</i> , 1985, 5, 303-309.	1.7	11
309	Atypical Distribution of Asymmetric Acetylcholinesterase in Mutant PC12 Pheochromocytoma Cells Lacking a Cell Surface Heparan Sulfate Proteoglycan. <i>Journal of Neurochemistry</i> , 1985, 45, 86-94.	2.1	28
310	Membrane-Bound Form of Acetylcholinesterase Activated during Postnatal Development of the Rat Somatosensory Cortex. <i>Developmental Neuroscience</i> , 1985, 7, 120-132.	1.0	15
311	Age-Related Responses of Skeletal Muscle After Ectopic Innervation, with Particular Reference to 16S Acetylcholinesterase, in Adult Rats. <i>Journal of Neurochemistry</i> , 1984, 43, 375-381.	2.1	1
312	The electric organ of <i>Discopyge tschudii</i> : Its innervated face and the biology of acetylcholinesterase. <i>Cellular and Molecular Neurobiology</i> , 1984, 4, 125-142.	1.7	14
313	Subcellular localization of acetylcholinesterase molecular forms in endplate regions of adult mammalian skeletal muscle. <i>Neurochemical Research</i> , 1984, 9, 1211-1230.	1.6	41
314	Acetylcholinesterase is functional in embryonic rat muscle before its accumulation at the sites of nerve-muscle contact. <i>Developmental Biology</i> , 1984, 103, 369-377.	0.9	21
315	Increase of muscle peroxisomal enzymes and myotonia induced by nafenopin, a hypolipidemic drug. <i>Muscle and Nerve</i> , 1983, 6, 154-159.	1.0	12
316	The A12 acetylcholinesterase and polypeptide composition of electric organ basal lamina of electrophorus and some torpedinae fishes. <i>Cell Biochemistry and Function</i> , 1983, 1, 41-48.	1.4	13
317	Heparin solubilizes asymmetric acetylcholinesterase from rat neuromuscular junction. <i>FEBS Letters</i> , 1983, 154, 265-268.	1.3	47
318	Association of the synaptic form of acetylcholinesterase with extracellular matrix in cultured mouse muscle cells. <i>Cell</i> , 1982, 29, 71-79.	13.5	100
319	Differentiation of skeletal muscle cells in culture.. <i>Cell Structure and Function</i> , 1982, 7, 91-109.	0.5	18
320	Gangliosides and sialoglycoproteins in normal and denervated rat diaphragm muscle. <i>Muscle and Nerve</i> , 1982, 5, 33-38.	1.0	9
321	Aneural muscle cell cultures make synaptic basal lamina components. <i>Nature</i> , 1982, 295, 143-145.	13.7	71
322	Acetylcholinesterase aggregates in a newly formed motor nerve-smooth muscle junction. <i>Brain Research Bulletin</i> , 1981, 7, 17-24.	1.4	3
323	Purification of the peroxisomal fatty acyl-CoA oxidase from rat liver. <i>Biochemical and Biophysical Research Communications</i> , 1980, 95, 7-12.	1.0	49
324	Acetylcholinesterase like that of skeletal muscle in smooth muscle reinnervated by a motor nerve. <i>Nature</i> , 1979, 280, 504-506.	13.7	37

#	ARTICLE	IF	CITATIONS
325	Fatty acid oxidation by human liver peroxisomes. Biochemical and Biophysical Research Communications, 1979, 88, 1030-1036.	1.0	201
326	Properties of fatty acyl-CoA oxidase from rat liver, a peroxisomal flavoprotein. Life Sciences, 1979, 25, 1127-1135.	2.0	24
327	Is there a correspondence between half-lives of motor endplate acetylcholinesterase and junctional acetylcholine receptors?. Neuroscience Letters, 1977, 5, 91-93.	1.0	2
328	Role of axoplasmic transport in neurotrophic regulation of muscle end plate acetylcholinesterase. Nature, 1976, 262, 55-56.	13.7	60
329	Glycogen Synthase Kinase 3 β (GSK-3 β) a Key Signaling Enzyme: A Developmental Neurobiological Perspective. , 0, , 25-43.		1
330	Comprehensive Overview of Alzheimer's Disease Neurodegeneration, from Amyloid- β to Neuroinflammatory Modulation. , 0, , .		0