
Takahiro Watakabe

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8676554/publications.pdf

Version: 2024-02-01



#	Article	IF	CITATIONS
1	Spleen Dose–Volume Parameters as a Predictor of Treatment-related Lymphopenia During Definitive Chemoradiotherapy for Esophageal Cancer. In Vivo, 2018, 32, 1519-1525.	1.3	29
2	Hypofractionated palliative volumetric modulated arc radiotherapy with the Radiation Oncology Study Group 8502 "QUAD shot―regimen for incurable head and neck cancer. Radiation Oncology, 2020, 15, 123.	2.7	17
3	Impact of hybrid FDG-PET/CT on gross tumor volume definition of cervical esophageal cancer: reducing interobserver variation. Journal of Radiation Research, 2019, 60, 348-352.	1.6	15
4	Diagnostic Value of FDG-PET/CT for the Identification of Extranodal Extension in Patients With Head and Neck Squamous Cell Carcinoma. Anticancer Research, 2020, 40, 2073-2077.	1.1	15
5	Treatment response after palliative radiotherapy for bleeding gastric cancer: a multicenter prospective observational study (JROSG 17-3). Gastric Cancer, 2022, 25, 411-421.	5.3	11
6	Impact of 99mTc-GSA SPECT Image-Guided Inverse Planning on Dose–Function Histogram Parameters for Stereotactic Body Radiation Therapy Planning for Patients With Hepatocellular Carcinoma: A Dosimetric Comparison Study. Dose-Response, 2019, 17, 155932581983214.	1.6	10
7	Dose–function Histogram Evaluation Using 99mTc-GSA SPECT/CT Images for Stereotactic Body Radiation Therapy Planning for Hepatocellular Carcinoma Patients: A Dosimetric Parameter Comparison. Anticancer Research, 2018, 38, 1511-1516.	1.1	8
8	Predictors of the Predominance of NonIndex Pain After Palliative Radiation Therapy for Painful Tumors. Advances in Radiation Oncology, 2019, 4, 118-126.	1.2	6
9	A neuropathic pain component as a predictor of improvement in pain interference after radiotherapy for painful tumors: A secondary analysis of a prospective observational study. Clinical and Translational Radiation Oncology, 2018, 12, 34-39.	1.7	5
10	High Spatial Resolution Digital Positron Emission Tomography Images With Dedicated Source-to-background Algorithm for Radiotherapy Planning. Anticancer Research, 2020, 40, 2567-2572.	1.1	5
11	Impact of four-dimensional cone-beam computed tomography on target localization for gastric mucosa-associated lymphoid tissue lymphoma radiotherapy: reducing planning target volume. Radiation Oncology, 2021, 16, 14.	2.7	5
12	Four-dimensional cone-beam computed tomography-guided radiotherapy for gastric lymphoma. Japanese Journal of Radiology, 2018, 36, 159-163.	2.4	4
13	Influence of pain duration on pain outcomes following palliative radiotherapy for painful tumors: the sooner the irradiation, the better?. Strahlentherapie Und Onkologie, 2021, 197, 916-925.	2.0	4
14	Palliative radiotherapy for painful lymph node metastases. Radiation Oncology, 2021, 16, 178.	2.7	4
15	A prospective comparison of adaptive and fixed boost plans in radiotherapy for glioblastoma. Radiation Oncology, 2022, 17, 40.	2.7	4
16	Plan Quality Comparisons Between 3D-CRT, IMRT, and VMAT Based on 4D-CT for Gastric MALT Lymphoma. Anticancer Research, 2021, 41, 3941-3947.	1.1	3
17	Concurrent Chemoradiotherapy With Docetaxel, Cisplatin, and 5-Fluorouracil for T3 N0 Glottic Carcinoma Without Vocal Cord Fixation. Anticancer Research, 2022, 42, 205-209.	1.1	3
18	Improvement in pain interference after palliative radiotherapy for solid and hematologic painful tumors: a secondary analysis of a prospective observational study. Japanese Journal of Clinical Oncology, 2018, 48, 982-987.	1.3	0

#	Article	IF	CITATIONS
19	QUAD shot: an effective cyclical hypofractionated palliative radiotherapy for salivary gland carcinoma. BJR case Reports, 2020, 6, 20190132.	0.2	Ο
20	Implementation of ^{99m} Tc-GSA SPECT Image-guided Inverse Planning into Palliative Radiotherapy for Diffuse Liver Metastases: A Novel Approach. In Vivo, 2022, 36, 1523-1526.	1.3	0