

# Lars Birger Engesäter

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8677644/publications.pdf>

Version: 2024-02-01

39  
papers

2,232  
citations

218677

26  
h-index

289244

40  
g-index

40  
all docs

40  
docs citations

40  
times ranked

2085  
citing authors

#	ARTICLE	IF	CITATIONS
1	Registration completeness in the Norwegian Arthroplasty Register. Monthly Notices of the Royal Astronomical Society: Letters, 2006, 77, 49-56.	3.3	167
2	Risk of revision for infection in primary total hip and knee arthroplasty in patients with rheumatoid arthritis compared with osteoarthritis: A prospective, population-based study on 108,786 hip and knee joint arthroplasties from the Norwegian Arthroplasty Register. Arthritis Care and Research, 2010, 62, 473-479.	3.4	158
3	The Norwegian Hip Fracture Register: Experiences after the first 2 years and 15,576 reported operations. Monthly Notices of the Royal Astronomical Society: Letters, 2008, 79, 583-593.	3.3	139
4	Does cement increase the risk of infection in primary total hip arthroplasty? Revision rates in 56,275 cemented and uncemented primary THAs followed for 0-16 years in the Norwegian Arthroplasty Register. Monthly Notices of the Royal Astronomical Society: Letters, 2006, 77, 351-358.	3.3	115
5	Surgical procedures in the treatment of 784 infected THAs reported to the Norwegian Arthroplasty Register. Monthly Notices of the Royal Astronomical Society: Letters, 2011, 82, 530-537.	3.3	108
6	Infection after primary hip arthroplasty. Monthly Notices of the Royal Astronomical Society: Letters, 2011, 82, 646-654.	3.3	105
7	18 years of results with cemented primary hip prostheses in the Norwegian Arthroplasty Register. Monthly Notices of the Royal Astronomical Society: Letters, 2009, 80, 402-412.	3.3	99
8	Dependency issues in survival analyses of 55 782 primary hip replacements from 47 355 patients. Statistics in Medicine, 2004, 23, 3227-3240.	1.6	97
9	Prevention of deep infection in joint replacement surgery. Monthly Notices of the Royal Astronomical Society: Letters, 2010, 81, 660-666.	3.3	87
10	Does time from fracture to surgery affect mortality and intraoperative medical complications for hip fracture patients?. Bone and Joint Journal, 2019, 101-B, 1129-1137.	4.4	85
11	Cemented or Uncemented Hemiarthroplasty for Femoral Neck Fracture? Data from the Norwegian Hip Fracture Register. Clinical Orthopaedics and Related Research, 2020, 478, 90-100.	1.5	74
12	Neonatal hip instability and risk of total hip replacement in young adulthood. Monthly Notices of the Royal Astronomical Society: Letters, 2008, 79, 321-326.	3.3	71
13	Implant survival and radiographic outcome of total hip replacement in patients less than 20 years old. Monthly Notices of the Royal Astronomical Society: Letters, 2016, 87, 479-484.	3.3	69
14	Improved results of primary total hip replacement. Monthly Notices of the Royal Astronomical Society: Letters, 2010, 81, 649-659.	3.3	68
15	Patient satisfaction, pain, and quality of life 4 months after displaced femoral neck fractures: A comparison of 663 fractures treated with internal fixation and 906 with bipolar hemiarthroplasty reported to the Norwegian Hip Fracture Register. Monthly Notices of the Royal Astronomical Society: Letters, 2008, 79, 594-601.	3.3	67
16	Low infection rates after 34,361 intramedullary nail operations in 55 low- and middle-income countries. Monthly Notices of the Royal Astronomical Society: Letters, 2011, 82, 737-743.	3.3	67
17	Quality of life following hip fractures: results from the Norwegian hip fracture register. BMC Musculoskeletal Disorders, 2016, 17, 265.	1.9	64
18	Standardized incidence rates of total hip replacement for primary hip osteoarthritis in the 5 Nordic countries: similarities and differences. Monthly Notices of the Royal Astronomical Society: Letters, 2006, 77, 733-740.	3.3	62

#	ARTICLE	IF	CITATIONS
19	Total hip replacement after femoral neck fractures in elderly patients : Results of 8,577 fractures reported to the Norwegian Arthroplasty Register. Monthly Notices of the Royal Astronomical Society: Letters, 2007, 78, 491-497.	3.3	61
20	Low revision rate after total hip arthroplasty in patients with pediatric hip diseases. Monthly Notices of the Royal Astronomical Society: Letters, 2012, 83, 436-441.	3.3	53
21	Protective Effect of Different Types of Bicycle Helmets. Traffic Injury Prevention, 2003, 4, 285-290.	1.4	51
22	Posterior approach compared to direct lateral approach resulted in better patient-reported outcome after hemiarthroplasty for femoral neck fracture. Monthly Notices of the Royal Astronomical Society: Letters, 2017, 88, 29-34.	3.3	49
23	WAITING TIME AND SOCIOECONOMIC STATUSâ€”AN INDIVIDUALâ€LEVEL ANALYSIS. Health Economics (United) Tj ETQq1 1.0.784314	1.7	41
24	Improved outcome after hip fracture surgery in Norway. Monthly Notices of the Royal Astronomical Society: Letters, 2017, 88, 505-511.	3.3	40
25	Early revision among 12,179 hip prostheses: A comparison of 10 different brands reported to the Norwegian Arthroplasty Register, 1987-1993. Acta Orthopaedica, 1995, 66, 487-493.	1.4	33
26	Outcome of 881 total hip arthroplasties in 747 patients 21 years or younger: data from the Nordic Arthroplasty Register Association (NARA) 1995â€2016. Monthly Notices of the Royal Astronomical Society: Letters, 2019, 90, 331-337.	3.3	30
27	Total hip arthroplasty in young adults, with focus on Perthes' disease and slipped capital femoral epiphysis. Monthly Notices of the Royal Astronomical Society: Letters, 2012, 83, 159-164.	3.3	25
28	Increasing Resistance of Coagulase-Negative Staphylococci in Total Hip Arthroplasty Infections: 278 THA-Revisions due to Infection Reported to the Norwegian Arthroplasty Register from 1993 to 2007. Advances in Orthopedics, 2014, 2014, 1-7.	1.0	22
29	Ponseti method compared to previous treatment of clubfoot in Norway. A multicenter study of 205 children followed for 8â€11 years. Journal of Children's Orthopaedics, 2016, 10, 445-452.	1.1	22
30	Do direct oral anticoagulants (DOACs) cause delayed surgery, longer length of hospital stay, and poorer outcome for hip fracture patients?. European Geriatric Medicine, 2020, 11, 563-569.	2.8	21
31	Antipsychotic Drugs and Risk of Hip Fracture in People Aged 60 and Older in Norway. Journal of the American Geriatrics Society, 2016, 64, 1203-1209.	2.6	20
32	Postoperative start compared to preoperative start of low-molecular-weight heparin increases mortality in patients with femoral neck fractures. Monthly Notices of the Royal Astronomical Society: Letters, 2017, 88, 48-54.	3.3	14
33	Consumption of hospital resources for hip fracture: Discharge rates for fracture in Norway. Acta Orthopaedica, 1985, 56, 17-20.	1.4	10
34	Bacterial and Hematological Findings in Infected Total Hip Arthroplasties in Norway Assessment of 278 Revisions Due to Infection in the Norwegian Arthroplasty Register. The Open Orthopaedics Journal, 2015, 9, 445-449.	0.2	9
35	Effects of intramedullary reaming and nailing of rat femur: A mechanical and chemical study. Acta Orthopaedica, 1991, 62, 582-586.	1.4	8
36	Intra- and inter-observer repeatability of radiographic measurements for previously slipped capital femoral epiphysis at skeletal maturity. Acta Radiologica, 2013, 54, 587-591.	1.1	7

#	ARTICLE	IF	CITATIONS
37	Validation of orthopaedic surgeons' assessment of cognitive function in patients with acute hip fracture. BMC Musculoskeletal Disorders, 2019, 20, 268.	1.9	6
38	Patient-reported outcome measures after hip fracture in patients with chronic cognitive impairment. Bone & Joint Open, 2021, 2, 454-465.	2.6	6
39	Reply to the Letter to the Editor: Cemented or Uncemented Hemiarthroplasty for Femoral Neck Fracture? Data from the Norwegian Hip Fracture Register. Clinical Orthopaedics and Related Research, 2020, 478, 687-689.	1.5	1