

# Ismet Cok

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8678782/publications.pdf>

Version: 2024-02-01

49  
papers

1,015  
citations

394286

19  
h-index

454834

30  
g-index

50  
all docs

50  
docs citations

50  
times ranked

1310  
citing authors

#	ARTICLE	IF	CITATIONS
1	Analysis of human milk to assess exposure to PAHs, PCBs and organochlorine pesticides in the vicinity Mediterranean city Mersin, Turkey. <i>Environment International</i> , 2012, 40, 63-69.	4.8	99
2	Organochlorine Pesticide Residues in Human Breast Milk from Agricultural Regions of Turkey, 1995A–1996. <i>Bulletin of Environmental Contamination and Toxicology</i> , 1997, 59, 577-582.	1.3	62
3	Cloning, Sequencing, and Tissue Expression of CYP3A27, a New Member of the CYP3A Subfamily from Embryonic and Adult Rainbow Trout Livers. <i>Archives of Biochemistry and Biophysics</i> , 1998, 360, 53-61.	1.4	47
4	The Organochlorine Pesticide Residues and Antioxidant Enzyme Activities in Human Breast Tumors: Is There Any Association?. <i>Breast Cancer Research and Treatment</i> , 2002, 72, 173-182.	1.1	45
5	An in vivo assessment of the genotoxic potential of bisphenol A and 4-tert-octylphenol in rats. <i>Archives of Toxicology</i> , 2011, 85, 995-1001.	1.9	43
6	Polychlorinated dibenzo-p-dioxins, dibenzofurans and polychlorinated biphenyls levels in human breast milk from different regions of Turkey. <i>Chemosphere</i> , 2009, 76, 1563-1571.	4.2	41
7	Organochlorine Pesticide Contaminants in Human Milk Samples Collected in Tebriz (Iran). <i>Bulletin of Environmental Contamination and Toxicology</i> , 1999, 63, 444-450.	1.3	37
8	N-Acetyltransferase Phenotype of Patients with Bladder Cancer. <i>Human Toxicology</i> , 1986, 5, 333-335.	0.9	36
9	Immunohistochemical Localization and Differential Expression of Cytochrome P450 3A27 in the Gastrointestinal Tract of Rainbow Trout. <i>Toxicology and Applied Pharmacology</i> , 2001, 177, 94-102.	1.3	34
10	Polychlorinated Biphenyl and Organochlorine Pesticide Levels in Human Breast Milk from the Mediterranean city Antalya, Turkey. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2011, 86, 423-427.	1.3	34
11	The Contamination Levels of Organochlorine Pesticides in Water and Sediment Samples in Uluabat Lake, Turkey. <i>Environmental Monitoring and Assessment</i> , 2006, 118, 383-391.	1.3	33
12	Polychlorinated biphenyl levels in adipose tissue of primiparous women in Turkey. <i>Environment International</i> , 2004, 30, 7-10.	4.8	32
13	Determination of urinary levels of Bisphenol A in a Turkish population. <i>Environmental Monitoring and Assessment</i> , 2014, 186, 8443-8452.	1.3	30
14	Concentration of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans and dioxin-like PCBs in human adipose tissue from Turkish men. <i>Chemosphere</i> , 2007, 66, 1955-1961.	4.2	27
15	Urinary cotinine levels of electronic cigarette (e-cigarette) users. <i>Toxicology Mechanisms and Methods</i> , 2016, 26, 441-445.	1.3	27
16	Urinary cotinine levels of smokeless tobacco (MaraE powder) users1. <i>Human and Experimental Toxicology</i> , 2000, 19, 650-655.	1.1	24
17	Expression of CYP2M1, CYP2K1, and CYP3A27 in Brain, Blood, Small Intestine, and Other Tissues of Rainbow Trout. <i>Biochemical and Biophysical Research Communications</i> , 1998, 244, 790-795.	1.0	21
18	Concentrations of Polychlorinated Dibenzo-p-Dioxins (PCDDs), Polychlorinated Dibenzofurans (PCDFs), and Dioxin-Like PCBs in Adipose Tissue of Infertile Men. <i>Archives of Environmental Contamination and Toxicology</i> , 2008, 55, 143-152.	2.1	20

#	ARTICLE	IF	CITATIONS
19	Evaluation of DNA Damage in Common Carp ( <i>Cyprinus carpio</i> L.) by Comet Assay for Determination of Possible Pollution in Lake Mogan (Ankara). Scientific World Journal, The, 2011, 11, 1455-1461.	0.8	19
20	Development and validation of an LC-MS/MS method for simultaneous quantitative analysis of free and conjugated bisphenol A in human urine. Biomedical Chromatography, 2014, 28, 686-693.	0.8	19
21	New psychoactive substances in Turkey: Narcotics cases assessed by the Council of Forensic Medicine between 2016 and 2017 in Ankara, Turkey. Forensic Science International, 2019, 294, 113-123.	1.3	18
22	Polymorphic N-acetylation capacity in breast cancer patients. International Journal of Cancer, 1990, 46, 1138-1139.	2.3	17
23	Levels and Trends of Chlorinated Pesticides in Human Breast Milk from Ankara Residents: Comparison of Concentrations in 1984 and 2002. Bulletin of Environmental Contamination and Toxicology, 2004, 72, 522-529.	1.3	17
24	Organochlorine Pesticide Residue Levels in Human Adipose Tissue of Residents of Manisa (Turkey), 1995-1996. Bulletin of Environmental Contamination and Toxicology, 1998, 61, 311-316.	1.3	15
25	Determination of urinary hippuric acid and o-cresol levels as biological indicators of toluene exposure in shoe-workers and glue sniffers. Biomarkers, 2003, 8, 119-127.	0.9	15
26	Determination of organochlorine pesticide and polychlorinated biphenyl levels in adipose tissue of infertile men. Environmental Monitoring and Assessment, 2010, 162, 301-309.	1.3	14
27	Blood levels of polychlorinated biphenyls and organochlorinated pesticides in women from Istanbul, Turkey. Environmental Monitoring and Assessment, 2015, 187, 132.	1.3	14
28	Determination of coumarin metabolism in Turkish population. Human and Experimental Toxicology, 2001, 20, 179-184.	1.1	13
29	Effects of Smoking during Pregnancy on DNA Damage and ROS Level Consequences in Maternal and Newborns' Blood. Arhiv Za Higijenu Rada I Toksikologiju, 2013, 64, 35-46.	0.4	13
30	Determination of 7-ethoxyresorufin-o-deethylase (EROD) induction in leaping mullet ( <i>Liza saliens</i> ) from the highly contaminated Aliaga Bay, Turkey. Environmental Monitoring and Assessment, 2010, 165, 87-96.	1.3	12
31	Genotoxicity of anti-tumor necrosis factor therapy in patients with juvenile idiopathic arthritis. Arthritis Care and Research, 2010, 62, 73-77.	1.5	12
32	DISPERSIVE LIQUID-LIQUID MICROEXTRACTION BASED ON SOLIDIFICATION OF FLOATING ORGANIC DROP COMBINED WITH COUNTER-ELECTROSMOTIC FLOW NORMAL STACKING MODE IN CAPILLARY ELECTROPHORESIS FOR THE DETERMINATION OF BISPHENOL A IN WATER AND URINE SAMPLES. Journal of Liquid Chromatography and Related Technologies, 2013, 36, 2855-2870.	0.5	11
33	Assessment of types of synthetic cannabinoids in narcotic cases assessed by the Council of Forensic Medicine between 2011-2015, Ankara, Turkey. Forensic Science International, 2017, 280, 124-129.	1.3	11
34	Psychoactive Bath Salts and Neurotoxicity Risk. Turkish Journal of Pharmaceutical Sciences, 2020, 17, 235-241.	0.6	11
35	Evaluation of organochlorine pesticides in breast milk samples in Turkey applying features of the partial order technique. International Journal of Environmental Health Research, 2013, 23, 226-246.	1.3	10
36	Evaluation of the endocrine-disrupting effects of homosalate (HMS) and 2-ethylhexyl 4-dimethylaminobenzoate (OD-PABA) in rat pups during the prenatal, lactation, and early postnatal periods. Toxicology and Industrial Health, 2017, 33, 775-791.	0.6	10

#	ARTICLE	IF	CITATIONS
37	Determination of the impurities in drug products containing montelukast and in silico/in vitro genotoxicological assessments of sulfoxide impurity. <i>Toxicology Letters</i> , 2015, 238, 90-99.	0.4	9
38	Blood concentrations and risk assessment of persistent organochlorine compounds in newborn boys in Turkey. A pilot study. <i>Environmental Science and Pollution Research</i> , 2015, 22, 19896-19904.	2.7	9
39	Assessment of Bisphenol A Levels in Preschool Children: Results of a Human Biomonitoring Study in Ankara, Turkey. <i>JCRPE Journal of Clinical Research in Pediatric Endocrinology</i> , 2020, 12, 86-94.	0.4	9
40	Assessment of DNA damage in glue sniffers by use of the alkaline comet assay. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2004, 557, 131-136.	0.9	8
41	Biomonitoring of bisphenol A, 4-nonylphenol, and 4-t-octylphenol in Turkish population: exposure and risk assessment. <i>Environmental Science and Pollution Research</i> , 2022, 29, 26250-26262.	2.7	8
42	Establishment of rapid, sensitive, and quantitative liquid chromatography-electrospray ionization-tandem mass spectrometry method coupled with liquid-liquid extraction for measurement of urinary bisphenol A, 4-octylphenol, and 4-nonylphenol. <i>Rapid Communications in Mass Spectrometry</i> , 2021, 35, e9084.	0.7	7
43	Biological monitoring of n-hexane exposure in shoe workers. <i>Biomarkers</i> , 1997, 2, 25-28.	0.9	6
44	The role of DENND1A and CYP19A1 gene variants in individual susceptibility to obesity in Turkish population—a preliminary study. <i>Molecular Biology Reports</i> , 2018, 45, 2193-2199.	1.0	5
45	Determination of the acetylator phenotype in a Turkish population. <i>Clinical Genetics</i> , 1986, 29, 185-186.	1.0	4
46	CombiSimilarity, an innovative method to compare environmental and health data sets with different attribute sizes example: Eighteen Organochlorine Pesticides in soil and human breast milk samples. <i>Ecotoxicology and Environmental Safety</i> , 2014, 105, 29-35.	2.9	4
47	An investigation of the mutagenic activity of salamide—a major impurity of hydrochlorothiazide. <i>Toxicology Mechanisms and Methods</i> , 2016, 26, 644-649.	1.3	2
48	Assessment of DNA damage in glue sniffers by use of the alkaline comet assay. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2004, 557, 131-6.	0.4	1
49	PyHasse Software Features Applied on the Evaluation of Chemicals in Human Breast Milk Samples in Turkey. , 2014, , 343-357.		0