

Kristin Holvik

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8680493/kristin-holvik-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

51
papers

1,254
citations

20
h-index

34
g-index

57
ext. papers

1,442
ext. citations

4.9
avg, IF

4.21
L-index

#	Paper	IF	Citations
51	Prevalence and predictors of vitamin D deficiency in five immigrant groups living in Oslo, Norway: the Oslo Immigrant Health Study. <i>European Journal of Clinical Nutrition</i> , 2005 , 59, 57-63	5.2	148
50	Mortality following the first hip fracture in Norwegian women and men (1999-2008). A NOREPOS study. <i>Bone</i> , 2014 , 63, 81-6	4.7	98
49	Standardizing serum 25-hydroxyvitamin D data from four Nordic population samples using the Vitamin D Standardization Program protocols: Shedding new light on vitamin D status in Nordic individuals. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2015 , 75, 549-61	2	80
48	Hip fractures in Norway 1999-2008: time trends in total incidence and second hip fracture rates: a NOREPOS study. <i>European Journal of Epidemiology</i> , 2012 , 27, 807-14	12.1	75
47	Predictors of mortality in older hip fracture inpatients admitted to an orthogeriatric unit in oslo, norway. <i>Journal of Aging and Health</i> , 2010 , 22, 1114-31	2.6	63
46	Low serum levels of 25-hydroxyvitamin D predict hip fracture in the elderly: a NOREPOS study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 3341-50	5.6	59
45	Abdominal obesity increases the risk of hip fracture. A population-based study of 43,000 women and men aged 60-79 years followed for 8 years. Cohort of Norway. <i>Journal of Internal Medicine</i> , 2015 , 277, 306-317	10.8	52
44	Older hip fracture patients: three groups with different needs. <i>BMC Geriatrics</i> , 2010 , 10, 65	4.1	52
43	Should vitamin D supplements be recommended to prevent chronic diseases?. <i>BMJ, The</i> , 2015 , 350, h3215	5.9	41
42	Continued decline in hip fracture incidence in Norway: a NOREPOS study. <i>Osteoporosis International</i> , 2016 , 27, 2217-2222	5.3	36
41	Cohort Profile Update: The Janus Serum Bank Cohort in Norway. <i>International Journal of Epidemiology</i> , 2017 , 46, 1101-1102f	7.8	29
40	Plasma osteocalcin levels as a predictor of cardiovascular disease in older men and women: a population-based cohort study. <i>European Journal of Endocrinology</i> , 2014 , 171, 161-70	6.5	28
39	Vitamin D status in Sri Lankans living in Sri Lanka and Norway. <i>British Journal of Nutrition</i> , 2008 , 99, 941-4	5.6	28
38	Use of warfarin is associated with delay in surgery for hip fracture in older patients. <i>Hospital Practice (1995)</i> , 2011 , 39, 37-40	2.2	27
37	Vitamin D status and current policies to achieve adequate vitamin D intake in the Nordic countries. <i>Scandinavian Journal of Public Health</i> , 2021 , 49, 616-627	3	25
36	Biochemical markers of bone turnover and their relation to forearm bone mineral density in persons of Pakistani and Norwegian background living in Oslo, Norway: The Oslo Health Study. <i>European Journal of Endocrinology</i> , 2006 , 155, 693-9	6.5	24
35	A randomised comparison of increase in serum 25-hydroxyvitamin D concentration after 4 weeks of daily oral intake of 10 microg cholecalciferol from multivitamin tablets or fish oil capsules in healthy young adults. <i>British Journal of Nutrition</i> , 2007 , 98, 620-5	3.6	24

34	Age and Sex Differences in Body Mass Index as a Predictor of Hip Fracture: A NOREPOS Study. <i>American Journal of Epidemiology</i> , 2016 , 184, 510-519	3.8	23
33	Excess mortality following hip fracture: impact of self-perceived health, smoking, and body mass index. A NOREPOS study. <i>Osteoporosis International</i> , 2017 , 28, 881-887	5.3	22
32	Do cadmium, lead, and aluminum in drinking water increase the risk of hip fractures? A NOREPOS study. <i>Biological Trace Element Research</i> , 2014 , 157, 14-23	4.5	20
31	No increase in risk of hip fracture at high serum retinol concentrations in community-dwelling older Norwegians: the Norwegian Epidemiologic Osteoporosis Studies. <i>American Journal of Clinical Nutrition</i> , 2015 , 102, 1289-96	7	19
30	Low serum concentrations of alpha-tocopherol are associated with increased risk of hip fracture. A NOREPOS study. <i>Osteoporosis International</i> , 2014 , 25, 2545-54	5.3	19
29	Sodium and Potassium Intake Assessed by Spot and 24-h Urine in the Population-Based Tromsø Study 2015-2016. <i>Nutrients</i> , 2019 , 11,	6.7	18
28	Impact of comorbidity, age, and gender on seasonal variation in hip fracture incidence. A NOREPOS study. <i>Archives of Osteoporosis</i> , 2014 , 9, 191	2.9	18
27	Association of High Intakes of Vitamins B6 and B12 From Food and Supplements With Risk of Hip Fracture Among Postmenopausal Women in the Nurses' Health Study. <i>JAMA Network Open</i> , 2019 , 2, e193591	10.4	17
26	Vitamin D status in psychotic disorder patients and healthy controls--The influence of ethnic background. <i>Psychiatry Research</i> , 2015 , 230, 616-21	9.9	17
25	Does the Association of Comorbidity with 1-Year Mortality After Hip Fracture Differ According to Gender? The Norwegian Epidemiologic Osteoporosis Studies (NOREPOS). <i>Journal of the American Geriatrics Society</i> , 2018 , 66, 553-558	5.6	17
24	A combination of low serum concentrations of vitamins K1 and D is associated with increased risk of hip fractures in elderly Norwegians: a NOREPOS study. <i>Osteoporosis International</i> , 2016 , 27, 1645-1652	5.3	17
23	Vitamin A and D intake in pregnancy, infant supplementation, and asthma development: the Norwegian Mother and Child Cohort. <i>American Journal of Clinical Nutrition</i> , 2018 , 107, 789-798	7	15
22	25-Hydroxyvitamin D in pregnancy and genome wide cord blood DNA methylation in two pregnancy cohorts (MoBa and ALSPAC). <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2016 , 159, 102-9	5.1	15
21	Pakistanis living in Oslo have lower serum 1,25-dihydroxyvitamin D levels but higher serum ionized calcium levels compared with ethnic Norwegians. The Oslo Health Study. <i>BMC Endocrine Disorders</i> , 2007 , 7, 9	3.3	14
20	The association between alcohol consumption and risk of hip fracture differs by age and gender in Cohort of Norway: a NOREPOS study. <i>Osteoporosis International</i> , 2018 , 29, 2457-2467	5.3	12
19	Nationwide data on municipal drinking water and hip fracture: could calcium and magnesium be protective? A NOREPOS study. <i>Bone</i> , 2013 , 57, 84-91	4.7	12
18	Procollagen type 1 amino-terminal propeptide (P1NP) and risk of hip fractures in elderly Norwegian men and women. A NOREPOS study. <i>Bone</i> , 2014 , 64, 1-7	4.7	12
17	Population data on calcium in drinking water and hip fracture: An association may depend on other minerals in water. A NOREPOS study. <i>Bone</i> , 2015 , 81, 292-299	4.7	11

16	Osteoporosis and osteopenia in the distal forearm predict all-cause mortality independent of grip strength: 22-year follow-up in the population-based Tromsø Study. <i>Osteoporosis International</i> , 2018 , 29, 2447-2456	5.3	11
15	Changes in the vitamin D endocrine system and bone turnover after oral vitamin D3 supplementation in healthy adults: results of a randomised trial. <i>BMC Endocrine Disorders</i> , 2012 , 12, 7	3.3	11
14	Educational Inequalities in Post-Hip Fracture Mortality: A NOREPOS Study. <i>Journal of Bone and Mineral Research</i> , 2015 , 30, 2221-8	6.3	10
13	A restrictive policy for red blood cell transfusion in older hip fracture patients: experiences from a patient register. <i>BMC Research Notes</i> , 2016 , 9, 75	2.3	8
12	Milk drinking and risk of hip fracture: the Norwegian Epidemiologic Osteoporosis Studies (NOREPOS). <i>British Journal of Nutrition</i> , 2019 , 121, 709-718	3.6	7
11	Geographic variations in hip fracture incidence in a high-risk country stretching into the Arctic: a NOREPOS study. <i>Osteoporosis International</i> , 2020 , 31, 1323-1331	5.3	5
10	Urban-Rural Differences in Hip Fracture Mortality: A Nationwide NOREPOS Study. <i>JBMR Plus</i> , 2019 , 3, e10236	3.9	5
9	Increased Mortality in Hip Fracture Patients Living Alone: A NOREPOS Study. <i>Journal of Bone and Mineral Research</i> , 2021 , 36, 480-488	6.3	4
8	Can bone mineral density loss in the non-weight bearing distal forearm predict mortality?. <i>Bone</i> , 2020 , 136, 115347	4.7	2
7	Incidence of injuries in Norway: linking primary and secondary care data. <i>Scandinavian Journal of Public Health</i> , 2020 , 48, 323-330	3	2
6	Means of increasing response rates in a Norwegian dietary survey among infants - results from a pseudo-randomized pilot study. <i>BMC Medical Research Methodology</i> , 2019 , 19, 144	4.7	0
5	Health care utilisation for treatment of injuries among immigrants in Norway: a nationwide register linkage study. <i>Injury Epidemiology</i> , 2020 , 7, 60	1.7	0
4	Contribution of an extensive medication-based comorbidity index (Rx-Risk) in explaining the excess mortality after hip fracture in older Norwegians: a NOREPOS cohort study.. <i>BMJ Open</i> , 2022 , 12, e057823	3	0
3	THE AUTHORS REPLY. <i>American Journal of Epidemiology</i> , 2017 , 185, 511-513	3.8	
2	Individual Variation in Adaptive Immune Responses and Risk of Hip Fracture-A NOREPOS Population-Based Cohort Study. <i>Journal of Bone and Mineral Research</i> , 2020 , 35, 2327-2334	6.3	
1	Re: "Hip Fracture and Mortality: A Loss of Life Expectancy Interpretation" by Thao T Ho-Le and Tuan V Nguyen. <i>Journal of Bone and Mineral Research</i> , 2021 ,	6.3	