Shang-Ze Li

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8681583/publications.pdf Version: 2024-02-01



SHANC-7ELL

#	Article	IF	CITATIONS
1	NLK is required for Ras/ERK/SRF/ELK signaling to tune skeletal muscle development by phosphorylating SRF and antagonizing the SRF/MKL pathway. Cell Death Discovery, 2022, 8, 4.	4.7	1
2	PTPN18 promotes colorectal cancer progression by regulating the c-MYC-CDK4 axis. Genes and Diseases, 2021, 8, 838-848.	3.4	11
3	BRCC3 Promotes Tumorigenesis of Bladder Cancer by Activating the NF-κB Signaling Pathway Through Targeting TRAF2. Frontiers in Cell and Developmental Biology, 2021, 9, 720349.	3.7	10
4	UCHL3 promotes ovarian cancer progression by stabilizing TRAF2 to activate the NF-κB pathway. Oncogene, 2020, 39, 322-333.	5.9	37
5	Cordycepin induces Bax‑dependent apoptosis in colorectal cancer cells. Molecular Medicine Reports, 2019, 19, 901-908.	2.4	14
6	Phosphorylation of MAVS/VISA by Nemo-like kinase (NLK) for degradation regulates the antiviral innate immune response. Nature Communications, 2019, 10, 3233.	12.8	35
7	Cancer testis antigen 55 deficiency attenuates colitis-associated colorectal cancer by inhibiting NF-κB signaling. Cell Death and Disease, 2019, 10, 304.	6.3	9
8	Activation of MAPK Signaling by CXCR7 Leads to Enzalutamide Resistance in Prostate Cancer. Cancer Research, 2019, 79, 2580-2592.	0.9	85
9	Nemo-like kinase (NLK) primes colorectal cancer progression by releasing the E2F1 complex from HDAC1. Cancer Letters, 2018, 431, 43-53.	7.2	23
10	Polycomb-Mediated Disruption of an Androgen Receptor Feedback Loop Drives Castration-Resistant Prostate Cancer. Cancer Research, 2017, 77, 412-422.	0.9	23
11	FOXA1 potentiates lineage-specific enhancer activation through modulating TET1 expression and function. Nucleic Acids Research, 2016, 44, 8153-8164.	14.5	53
12	Dynamic Phosphorylation of CENP-A at Ser68 Orchestrates Its Cell-Cycle-Dependent Deposition at Centromeres. Developmental Cell, 2015, 32, 68-81.	7.0	92
13	Rhomboid domain containing 1 promotes colorectal cancer growth through activation of the EGFR signalling pathway. Nature Communications, 2015, 6, 8022.	12.8	63
14	Stabilization of ATF5 by TAK1–Nemo-Like Kinase Critically Regulates the Interleukin-1β-Stimulated C/EBP Signaling Pathway. Molecular and Cellular Biology, 2015, 35, 778-788.	2.3	20
15	The Selective Activation of p53 Target Genes Regulated by SMYD2 in BIX-01294 Induced Autophagy-Related Cell Death. PLoS ONE, 2015, 10, e0116782.	2.5	29
16	miR-191 promotes tumorigenesis of human colorectal cancer through targeting C/EBPβ. Oncotarget, 2015, 6, 4144-4158.	1.8	58
17	HSCARG, a novel regulator of H2A ubiquitination by downregulating PRC1 ubiquitin E3 ligase activity, is essential for cell proliferation. Nucleic Acids Research, 2014, 42, 5582-5593.	14.5	22
18	Nemo-like kinase (NLK) negatively regulates NF-kappa B activity through disrupting the interaction of TAK1 with IKKβ. Biochimica Et Biophysica Acta - Molecular Cell Research, 2014, 1843, 1365-1372.	4.1	27

#	Article	IF	CITATIONS
19	UbcH10 overexpression increases carcinogenesis and blocks ALLN susceptibility in colorectal cancer. Scientific Reports, 2014, 4, 6910.	3.3	23
20	ALLN hinders HCT116 tumor growth through Bax-dependent apoptosis. Biochemical and Biophysical Research Communications, 2013, 437, 325-330.	2.1	24