

Niels dum

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8682058/niels-odum-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

141
papers

4,494
citations

38
h-index

63
g-index

146
ext. papers

5,442
ext. citations

5
avg. IF

5.04
L-index

#	Paper	IF	Citations
141	Vitamin D controls T cell antigen receptor signaling and activation of human T cells. <i>Nature Immunology</i> , 2010 , 11, 344-9	19.1	408
140	Multilevel dysregulation of STAT3 activation in anaplastic lymphoma kinase-positive T/null-cell lymphoma. <i>Journal of Immunology</i> , 2002 , 168, 466-74	5.3	223
139	Diagnostic microRNA profiling in cutaneous T-cell lymphoma (CTCL). <i>Blood</i> , 2011 , 118, 5891-900	2.2	203
138	The effect of short-chain fatty acids on human monocyte-derived dendritic cells. <i>Scientific Reports</i> , 2015 , 5, 16148	4.9	180
137	Inhibition of constitutively activated Stat3 correlates with altered Bcl-2/Bax expression and induction of apoptosis in mycosis fungoides tumor cells. <i>Leukemia</i> , 1999 , 13, 735-8	10.7	171
136	In vivo activation of STAT3 in cutaneous T-cell lymphoma. Evidence for an antiapoptotic function of STAT3. <i>Leukemia</i> , 2004 , 18, 1288-95	10.7	136
135	Constitutive STAT3-activation in Sezary syndrome: tyrphostin AG490 inhibits STAT3-activation, interleukin-2 receptor expression and growth of leukemic Sezary cells. <i>Leukemia</i> , 2001 , 15, 787-93	10.7	135
134	STAT3-mediated constitutive expression of SOCS-3 in cutaneous T-cell lymphoma. <i>Blood</i> , 2001 , 97, 1056-62	10.7	110
133	Lack of phosphotyrosine phosphatase SHP-1 expression in malignant T-cell lymphoma cells results from methylation of the SHP-1 promoter. <i>American Journal of Pathology</i> , 2000 , 157, 1137-46	5.8	106
132	STAT5-mediated expression of oncogenic miR-155 in cutaneous T-cell lymphoma. <i>Cell Cycle</i> , 2013 , 12, 1939-47	4.7	103
131	Jak3- and JNK-dependent vascular endothelial growth factor expression in cutaneous T-cell lymphoma. <i>Leukemia</i> , 2006 , 20, 1759-66	10.7	95
130	miR-122 regulates p53/Akt signalling and the chemotherapy-induced apoptosis in cutaneous T-cell lymphoma. <i>PLoS ONE</i> , 2012 , 7, e29541	3.7	86
129	Malignant cutaneous T-cell lymphoma cells express IL-17 utilizing the Jak3/Stat3 signaling pathway. <i>Journal of Investigative Dermatology</i> , 2011 , 131, 1331-8	4.3	81
128	Vitamin D-binding protein controls T cell responses to vitamin D. <i>BMC Immunology</i> , 2014 , 15, 35	3.7	77
127	Malignant inflammation in cutaneous T-cell lymphoma-a hostile takeover. <i>Seminars in Immunopathology</i> , 2017 , 39, 269-282	12	73
126	Constitutive SOCS-3 expression protects T-cell lymphoma against growth inhibition by IFNalpha. <i>Leukemia</i> , 2005 , 19, 209-13	10.7	72
125	Notch1 as a potential therapeutic target in cutaneous T-cell lymphoma. <i>Blood</i> , 2010 , 116, 2504-12	2.2	67

124	Staphylococcal enterotoxin A (SEA) stimulates STAT3 activation and IL-17 expression in cutaneous T-cell lymphoma. <i>Blood</i> , 2016 , 127, 1287-96	2.2	60
123	Elucidating the role of interleukin-17F in cutaneous T-cell lymphoma. <i>Blood</i> , 2013 , 122, 943-50	2.2	59
122	Allergic contact dermatitis induces upregulation of identical microRNAs in humans and mice. <i>Contact Dermatitis</i> , 2012 , 67, 298-305	2.7	58
121	Jak3, STAT3, and STAT5 inhibit expression of miR-22, a novel tumor suppressor microRNA, in cutaneous T-Cell lymphoma. <i>Oncotarget</i> , 2015 , 6, 20555-69	3.3	58
120	Activated human CD4+ T cells express transporters for both cysteine and cystine. <i>Scientific Reports</i> , 2012 , 2, 266	4.9	56
119	Nonmalignant T cells stimulate growth of T-cell lymphoma cells in the presence of bacterial toxins. <i>Blood</i> , 2007 , 109, 3325-32	2.2	55
118	Antibiotics inhibit tumor and disease activity in cutaneous T-cell lymphoma. <i>Blood</i> , 2019 , 134, 1072-1083	2.2	54
117	Spontaneous interleukin-5 production in cutaneous T-cell lymphoma lines is mediated by constitutively activated Stat3. <i>Blood</i> , 2002 , 99, 973-7	2.2	52
116	Bacterial toxins fuel disease progression in cutaneous T-cell lymphoma. <i>Toxins</i> , 2013 , 5, 1402-21	4.9	49
115	Rapid allergen-induced interleukin-17 and interferon- γ secretion by skin-resident memory CD8 T cells. <i>Contact Dermatitis</i> , 2017 , 76, 218-227	2.7	48
114	Vitamin D up-regulates the vitamin D receptor by protecting it from proteasomal degradation in human CD4+ T cells. <i>PLoS ONE</i> , 2014 , 9, e96695	3.7	46
113	Characterization and expression of the human T cell receptor-T3 complex by monoclonal antibody F101.01. <i>Scandinavian Journal of Immunology</i> , 1988 , 27, 685-96	3.4	46
112	Single-cell heterogeneity in Sjögren syndrome. <i>Blood Advances</i> , 2018 , 2, 2115-2126	7.8	45
111	MicroRNA expression in early mycosis fungoides is distinctly different from atopic dermatitis and advanced cutaneous T-cell lymphoma. <i>Anticancer Research</i> , 2014 , 34, 7207-17	2.3	45
110	Analysis of STAT4 expression in cutaneous T-cell lymphoma (CTCL) patients and patient-derived cell lines. <i>Cell Cycle</i> , 2014 , 13, 2975-82	4.7	43
109	Activation of Stat-3 Is Involved in the Induction of Apoptosis After Ligation of Major Histocompatibility Complex Class I Molecules on Human Jurkat T Cells. <i>Blood</i> , 1998 , 91, 3566-3573	2.2	41
108	Regulation of vascular endothelial growth factor in prostate cancer. <i>Endocrine-Related Cancer</i> , 2015 , 22, R107-23	5.7	40
107	Staphylococcal enterotoxins stimulate lymphoma-associated immune dysregulation. <i>Blood</i> , 2014 , 124, 761-70	2.2	40

106	cMyc/miR-125b-5p signalling determines sensitivity to bortezomib in preclinical model of cutaneous T-cell lymphomas. <i>PLoS ONE</i> , 2013 , 8, e59390	3.7	40
105	Loss of SHP-1 tyrosine phosphatase expression correlates with the advanced stages of cutaneous T-cell lymphoma. <i>Human Pathology</i> , 2007 , 38, 462-7	3.7	40
104	EGFR induces expression of IRF-1 via STAT1 and STAT3 activation leading to growth arrest of human cancer cells. <i>International Journal of Cancer</i> , 2008 , 122, 342-9	7.5	39
103	Role of Dysregulated Cytokine Signaling and Bacterial Triggers in the Pathogenesis of Cutaneous T-Cell Lymphoma. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 1116-1125	4.3	38
102	MiR137 is an androgen regulated repressor of an extended network of transcriptional coregulators. <i>Oncotarget</i> , 2015 , 6, 35710-25	3.3	38
101	Butyrate and propionate inhibit antigen-specific CD8 T cell activation by suppressing IL-12 production by antigen-presenting cells. <i>Scientific Reports</i> , 2017 , 7, 14516	4.9	37
100	Circulating Cell-Free miR-375 as Surrogate Marker of Tumor Burden in Merkel Cell Carcinoma. <i>Clinical Cancer Research</i> , 2018 , 24, 5873-5882	12.9	35
99	Prognostic miRNA classifier in early-stage mycosis fungoides: development and validation in a Danish nationwide study. <i>Blood</i> , 2018 , 131, 759-770	2.2	34
98	A novel xenograft model of cutaneous T-cell lymphoma. <i>Experimental Dermatology</i> , 2010 , 19, 1096-102	4	33
97	Ectopic expression of embryonic stem cell and other developmental genes in cutaneous T-cell lymphoma. <i>Oncolmmunology</i> , 2014 , 3, e970025	7.2	31
96	STAT5 induces miR-21 expression in cutaneous T cell lymphoma. <i>Oncotarget</i> , 2016 , 7, 45730-45744	3.3	31
95	STAT3/5-Dependent IL9 Overexpression Contributes to Neoplastic Cell Survival in Mycosis Fungoides. <i>Clinical Cancer Research</i> , 2016 , 22, 3328-39	12.9	28
94	SATB1 in Malignant T Cells. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 1805-1815	4.3	28
93	NKG2D-dependent activation of dendritic epidermal T cells in contact hypersensitivity. <i>Journal of Investigative Dermatology</i> , 2015 , 135, 1311-1319	4.3	26
92	Malignant T cells express lymphotoxin and drive endothelial activation in cutaneous T cell lymphoma. <i>Oncotarget</i> , 2015 , 6, 15235-49	3.3	25
91	Analysis of CTCL cell lines reveals important differences between mycosis fungoides/Sjary syndrome leukemic cell lines. <i>Oncotarget</i> , 2017 , 8, 95981-95998	3.3	24
90	Human CD4+ T cells require exogenous cystine for glutathione and DNA synthesis. <i>Oncotarget</i> , 2015 , 6, 21853-64	3.3	24
89	Validation of a diagnostic microRNA classifier in cutaneous T-cell lymphomas. <i>Leukemia and Lymphoma</i> , 2014 , 55, 957-8	1.9	23

88	Spironolactone induces apoptosis and inhibits NF-kappaB independent of the mineralocorticoid receptor. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2006 , 11, 2159-65	5.4	23
87	ZAP-70 and p72syk are signaling response elements through MHC class II molecules. <i>Tissue Antigens</i> , 1995 , 46, 145-54		23
86	The inhibitory checkpoint, PD-L2, is a target for effector T cells: Novel possibilities for immune therapy. <i>Onc Immunology</i> , 2018 , 7, e1390641	7.2	22
85	Investigating potential exogenous tumor initiating and promoting factors for Cutaneous T-Cell Lymphomas (CTCL), a rare skin malignancy. <i>Onc Immunology</i> , 2016 , 5, e1175799	7.2	22
84	Epicutaneous exposure to nickel induces nickel allergy in mice via a MyD88-dependent and interleukin-1-dependent pathway. <i>Contact Dermatitis</i> , 2014 , 71, 224-32	2.7	22
83	Expression of miR-155 and miR-126 in situ in cutaneous T-cell lymphoma. <i>Apmis</i> , 2013 , 121, 1020-4	3.4	22
82	Malignant T cells secrete galectins and induce epidermal hyperproliferation and disorganized stratification in a skin model of cutaneous T-cell lymphoma. <i>Journal of Investigative Dermatology</i> , 2015 , 135, 238-246	4.3	21
81	Three distinct developmental pathways for adaptive and two IFN- γ -producing $\gamma\delta$ subsets in adult thymus. <i>Nature Communications</i> , 2017 , 8, 1911	17.4	20
80	Vitamin D Counteracts -Induced Cathelicidin Downregulation in Dendritic Cells and Allows Th1 Differentiation and IFN γ Secretion. <i>Frontiers in Immunology</i> , 2017 , 8, 656	8.4	20
79	Staphylococcal alpha-toxin tilts the balance between malignant and non-malignant CD4 T cells in cutaneous T-cell lymphoma. <i>Onc Immunology</i> , 2019 , 8, e1641387	7.2	19
78	Vascular endothelial growth factor receptor-3 expression in mycosis fungoides. <i>Leukemia and Lymphoma</i> , 2013 , 54, 819-26	1.9	19
77	FoxP3 mRNA splice forms in synovial CD4+ T cells in rheumatoid arthritis and psoriatic arthritis. <i>Apmis</i> , 2012 , 120, 387-96	3.4	18
76	IL-15 and IL-17F are differentially regulated and expressed in mycosis fungoides (MF). <i>Cell Cycle</i> , 2014 , 13, 1306-12	4.7	18
75	Pathogenic CD8 Epidermis-Resident Memory T Cells Displace Dendritic Epidermal T Cells in Allergic Dermatitis. <i>Journal of Investigative Dermatology</i> , 2020 , 140, 806-815.e5	4.3	18
74	Clonotypic Diversity of the T-cell Receptor Corroborates the Immature Precursor Origin of Cutaneous T-cell Lymphoma. <i>Clinical Cancer Research</i> , 2019 , 25, 3104-3114	12.9	16
73	Staphylococcal enterotoxin-A directly stimulates signal transduction and interferon-gamma production in psoriatic T-cell lines. <i>Tissue Antigens</i> , 1998 , 52, 530-8		13
72	Inhibition of constitutively activated Stat3 correlates with altered Bcl-2/Bax expression and induction of apoptosis in mycosis fungoides tumor cells. <i>Leukemia</i> , 2013 , 13, 735-738	10.7	13
71	MicroRNAs in the Pathogenesis, Diagnosis, Prognosis and Targeted Treatment of Cutaneous T-Cell Lymphomas. <i>Cancers</i> , 2020 , 12,	6.6	12

70	The role of PIP5K1 β /PAKT and targeted inhibition of growth of subtypes of breast cancer using PIP5K1 β inhibitor. <i>Oncogene</i> , 2019 , 38, 375-389	9.2	12
69	Radically altered T cell receptor signaling in glycopeptide-specific T cell hybridoma induced by antigen with minimal differences in the glycan group. <i>European Journal of Immunology</i> , 2001 , 31, 3197-2006	6.1	12
68	SHP2 regulates IL-2 induced MAPK activation, but not Stat3 or Stat5 tyrosine phosphorylation, in cutaneous T cell lymphoma cells. <i>Cytokine</i> , 2002 , 20, 141-7	4	12
67	Ubiquitin-specific protease 2 decreases p53-dependent apoptosis in cutaneous T-cell lymphoma. <i>Oncotarget</i> , 2016 , 7, 48391-48400	3.3	12
66	Staphylococcus aureus enterotoxins induce FOXP3 in neoplastic T cells in Sjögren syndrome. <i>Blood Cancer Journal</i> , 2020 , 10, 57	7	11
65	Human P2Y Expression Level Affects Human P2X7 Receptor-Mediated Cell Death. <i>Frontiers in Immunology</i> , 2018 , 9, 1159	8.4	11
64	MEK kinase 1 is a negative regulator of virus-specific CD8(+) T cells. <i>European Journal of Immunology</i> , 2006 , 36, 2076-84	6.1	11
63	T-cell receptor downregulation by ceramide-induced caspase activation and cleavage of the zeta chain. <i>Scandinavian Journal of Immunology</i> , 2001 , 53, 176-83	3.4	11
62	Improving oligo-conjugated antibody signal in multimodal single-cell analysis. <i>ELife</i> , 2021 , 10,	8.9	11
61	Merkel cell carcinoma-derived exosome-shuttle miR-375 induces fibroblast polarization by inhibition of RBPJ and p53. <i>Oncogene</i> , 2021 , 40, 980-996	9.2	11
60	Interleukin-26 (IL-26) is a novel anti-microbial peptide produced by T cells in response to staphylococcal enterotoxin. <i>Oncotarget</i> , 2018 , 9, 19481-19489	3.3	11
59	Gene variation in IL-7 receptor (IL-7R) β affects IL-7R response in CD4+ T cells in HIV-infected individuals. <i>Scientific Reports</i> , 2017 , 7, 42036	4.9	10
58	alpha-toxin inhibits CD8 T cell-mediated killing of cancer cells in cutaneous T-cell lymphoma. <i>Oncolimmunology</i> , 2020 , 9, 1751561	7.2	10
57	The metabolic enzyme arginase-2 is a potential target for novel immune modulatory vaccines. <i>Oncolimmunology</i> , 2020 , 9, 1771142	7.2	10
56	STAT3 Dysregulation in Mature T and NK Cell Lymphomas. <i>Cancers</i> , 2019 , 11,	6.6	10
55	Endo- and exocytic rate constants for spontaneous and protein kinase C-activated T cell receptor cycling. <i>European Journal of Immunology</i> , 2002 , 32, 616-626	6.1	10
54	The Escherichia coli protein toxin cytotoxic necrotizing factor 1 induces epithelial mesenchymal transition. <i>Cellular Microbiology</i> , 2020 , 22, e13138	3.9	10
53	STAT3 activation and infiltration of eosinophil granulocytes in mycosis fungoides. <i>Anticancer Research</i> , 2014 , 34, 5277-86	2.3	10

52	Peptidylarginine deiminase-4 gene polymorphisms are associated with systemic lupus erythematosus and lupus nephritis. <i>Scandinavian Journal of Rheumatology</i> , 2019 , 48, 133-140	1.9	9
51	The functional interlink between AR and MMP9/VEGF signaling axis is mediated through PIP5K1 β /pAKT in prostate cancer. <i>International Journal of Cancer</i> , 2020 , 146, 1686-1699	7.5	9
50	Increased Production of IL-17A-Producing $\gamma\delta$ T Cells in the Thymus of Filaggrin-Deficient Mice. <i>Frontiers in Immunology</i> , 2018 , 9, 988	8.4	8
49	Androgen dependent mechanisms of pro-angiogenic networks in placental and tumor development. <i>Placenta</i> , 2017 , 56, 79-85	3.4	7
48	Cellular Interactions and Inflammation in the Pathogenesis of Cutaneous T-Cell Lymphoma. <i>Frontiers in Cell and Developmental Biology</i> , 2020 , 8, 851	5.7	7
47	Midline 1 directs lytic granule exocytosis and cytotoxicity of mouse killer T cells. <i>European Journal of Immunology</i> , 2014 , 44, 3109-18	6.1	6
46	Expression of NAD(P)H quinone dehydrogenase 1 (NQO1) is increased in the endometrium of women with endometrial cancer and women with polycystic ovary syndrome. <i>Clinical Endocrinology</i> , 2017 , 87, 557-565	3.4	6
45	The Thioredoxin-Interacting Protein TXNIP Is a Putative Tumour Suppressor in Cutaneous T-Cell Lymphoma. <i>Dermatology</i> , 2021 , 237, 283-290	4.4	6
44	Discrete immune response signature to SARS-CoV-2 mRNA vaccination versus infection 2021 ,		6
43	Inhibition of succinate dehydrogenase activity impairs human T cell activation and function. <i>Scientific Reports</i> , 2021 , 11, 1458	4.9	6
42	MicroRNA-93 Targets p21 and Promotes Proliferation in Mycosis Fungoides T Cells. <i>Dermatology</i> , 2021 , 237, 277-282	4.4	5
41	Deregulated signalling and inflammation in cutaneous T-cell lymphoma. <i>British Journal of Dermatology</i> , 2020 , 182, 16-17	4	5
40	Hypopigmented Mycosis Fungoides: Loss of Pigmentation Reflects Antitumor Immune Response in Young Patients. <i>Cancers</i> , 2020 , 12,	6.6	5
39	Tumor necrosis factor induces rapid down-regulation of TXNIP in human T cells. <i>Scientific Reports</i> , 2019 , 9, 16725	4.9	5
38	Midline 1 controls polarization and migration of murine cytotoxic T cells. <i>Immunity, Inflammation and Disease</i> , 2014 , 2, 262-71	2.4	4
37	Multimodal single-cell analysis of cutaneous T-cell lymphoma reveals distinct subclonal tissue-dependent signatures. <i>Blood</i> , 2021 , 138, 1456-1464	2.2	4
36	Low SATB1 Expression Promotes IL-5 and IL-9 Expression in Sjögren Syndrome. <i>Journal of Investigative Dermatology</i> , 2020 , 140, 713-716	4.3	4
35	Staphylococcus aureus Induces Signal Transducer and Activator of Transcription 5-Dependent miR-155 Expression in Cutaneous T-Cell Lymphoma. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 2449-2458	4.3	4

34	The Expression of IL-21 Is Promoted by MEKK4 in Malignant T Cells and Associated with Increased Progression Risk in Cutaneous T-Cell Lymphoma. <i>Journal of Investigative Dermatology</i> , 2016 , 136, 866-869	4.3	3
33	Human thymic epithelial cells express functional HLA-DP molecules. <i>Tissue Antigens</i> , 1996 , 47, 300-6		3
32	Expression and function of Kv1.3 channel in malignant T cells in Sjögren syndrome. <i>Oncotarget</i> , 2019 , 10, 4894-4906	3.3	3
31	Ectopic expression of a novel CD22 splice-variant regulates survival and proliferation in malignant T cells from cutaneous T cell lymphoma (CTCL) patients. <i>Oncotarget</i> , 2015 , 6, 14374-84	3.3	3
30	The MicroRNA Expression Profile Differs Between Erythrodermic Mycosis Fungoides and Sjögren Syndrome. <i>Acta Dermato-Venereologica</i> , 2019 , 99, 1148-1153	2.2	3
29	Improving oligo-conjugated antibody signal in multimodal single-cell analysis		3
28	Impaired Vitamin D Signaling in T Cells From a Family With Hereditary Vitamin D Resistant Rickets. <i>Frontiers in Immunology</i> , 2021 , 12, 684015	8.4	3
27	Bacterial genotoxins induce T cell senescence. <i>Cell Reports</i> , 2021 , 35, 109220	10.6	3
26	OMIP-057: Mouse T-Cell Development Characterized by a 14 Color Flow Cytometry Panel. <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2019 , 95, 726-729	4.6	2
25	MID2 can substitute for MID1 and control exocytosis of lytic granules in cytotoxic T cells. <i>Apmis</i> , 2015 , 123, 682-7	3.4	2
24	Suppressed microRNA-195-5p expression in mycosis fungoides promotes tumor cell proliferation. <i>Experimental Dermatology</i> , 2021 , 30, 1141-1149	4	2
23	CDK1 links to RAR α treatment response of cancer cells. <i>Cell Cycle</i> , 2013 , 12, 1659-60	4.7	2
22	Establishment of Prostate Tumor Growth and Metastasis Is Supported by Bone Marrow Cells and Is Mediated by PIP5K1 β Lipid Kinase. <i>Cancers</i> , 2020 , 12,	6.6	2
21	Applicability of Small-Molecule Inhibitors in the Study of Peptidyl Arginine Deiminase 2 (PAD2) and PAD4. <i>Frontiers in Immunology</i> , 2021 , 12, 716250	8.4	2
20	CD8 tissue-resident memory T cells recruit neutrophils that are essential for flare-ups in contact dermatitis. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2021 ,	9.3	2
19	Expression of the Voltage-Gated Potassium Channel Kv1.3 in Lesional Skin from Patients with Cutaneous T-Cell Lymphoma and Benign Dermatitis. <i>Dermatology</i> , 2020 , 236, 123-132	4.4	2
18	Diagnostic Two-Gene Classifier in Early-Stage Mycosis Fungoides: A Retrospective Multicenter Study. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 213-217.e5	4.3	2
17	Vitamin D Inhibits IL-22 Production Through a Repressive Vitamin D Response Element in the Promoter. <i>Frontiers in Immunology</i> , 2021 , 12, 715059	8.4	2

16	Macrophages Control the Bioavailability of Vitamin D and Vitamin D-Regulated T Cell Responses. <i>Frontiers in Immunology</i> , 2021 , 12, 722806	8.4	2
15	Evidence of gene-gene interaction in hidradenitis suppurativa: a nationwide registry study of Danish twins. <i>British Journal of Dermatology</i> , 2021 ,	4	2
14	Low prevalence of antibodies and other plasma factors binding to CC chemokines and IL-2 in HIV-positive patients. <i>Apmis</i> , 2000 , 108, 122-30	3.4	1
13	Peptide vaccination activating Galectin-3-specific T cells offers a novel means to target Galectin-3-expressing cells in the tumor microenvironment.. <i>Oncolmmunology</i> , 2022 , 11, 2026020	7.2	1
12	Anti-regulatory T cells are natural regulatory effector T cells. <i>Cell Stress</i> , 2019 , 3, 310-311	5.5	1
11	Staphylococcus aureus and Antibiotics in Cutaneous T-Cell Lymphoma. <i>Dermatology</i> , 2021 , 1-3	4.4	1
10	Polymorphisms Confer Risk of Anti-CCP-Positive Rheumatoid Arthritis in Synergy With and Smoking. <i>Frontiers in Immunology</i> , 2021 , 12, 707690	8.4	1
9	Skin Associated Staphylococcus Aureus Contributes to Disease Progression in CTCL. <i>Blood</i> , 2019 , 134, 659-659	2.2	1
8	JAK3 Is Expressed in the Nucleus of Malignant T Cells in Cutaneous T Cell Lymphoma (CTCL). <i>Cancers</i> , 2021 , 13,	6.6	1
7	Normal T and B Cell Responses Against SARS-CoV-2 in a Family With a Non-Functional Vitamin D Receptor: A Case Report. <i>Frontiers in Immunology</i> , 2021 , 12, 758154	8.4	1
6	Haematopoietic stem cells and their niches. <i>Cell Cycle</i> , 2015 , 14, 3524-5	4.7	
5	Diagnostic 2-Gene Classifier in Early-Stage Mycosis Fungoides: A Retrospective Multicenter Study. <i>Blood</i> , 2019 , 134, 2772-2772	2.2	
4	Proinflammatory biomarkers are associated with prediabetes in patients with schizophrenia. <i>CNS Spectrums</i> , 2020 , 1-8	1.8	
3	Oncogenic kinase NPM/ALK induces expression of the cell-growth stimulatory receptor ICOS. <i>FASEB Journal</i> , 2011 , 25, 243.7	0.9	
2	Epidermal T cell subsets-Effect of age and antigen exposure in humans and mice. <i>Contact Dermatitis</i> , 2021 , 84, 375-384	2.7	
1	Omalizumab serum levels predict treatment outcomes in patients with chronic spontaneous urticaria: A three months prospective study.. <i>Clinical and Experimental Allergy</i> , 2022 ,	4.1	