

Antonio Molina

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8683120/publications.pdf>

Version: 2024-02-01

83
papers

14,579
citations

34493

54
h-index

68831

81
g-index

87
all docs

87
docs citations

87
times ranked

13391
citing authors

#	ARTICLE	IF	CITATIONS
1	Computational prediction method to decipher receptor-glycoligand interactions in plant immunity. <i>Plant Journal</i> , 2021, 105, 1710-1726.	2.8	14
2	Cell wall-derived mixed-linked β -1,3/1,4-glucans trigger immune responses and disease resistance in plants. <i>Plant Journal</i> , 2021, 106, 601-615.	2.8	69
3	<i>Arabidopsis</i> cell wall composition determines disease resistance specificity and fitness. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	88
4	The role of CYP71A12 monooxygenase in pathogen-triggered tryptophan metabolism and <i>Arabidopsis</i> immunity. <i>New Phytologist</i> , 2020, 225, 400-412.	3.5	51
5	YODA Kinase Controls a Novel Immune Pathway of Tomato Conferring Enhanced Disease Resistance to the Bacterium <i>Pseudomonas syringae</i> . <i>Frontiers in Plant Science</i> , 2020, 11, 584471.	1.7	9
6	Differential Expression of Fungal Genes Determines the Lifestyle of <i>Plectosphaerella</i> Strains During <i>Arabidopsis thaliana</i> Colonization. <i>Molecular Plant-Microbe Interactions</i> , 2020, 33, 1299-1314.	1.4	9
7	Arabinoxylan-Oligosaccharides Act as Damage Associated Molecular Patterns in Plants Regulating Disease Resistance. <i>Frontiers in Plant Science</i> , 2020, 11, 1210.	1.7	49
8	<i>Arabidopsis</i> Response Regulator 6 (ARR6) Modulates Plant Cell-Wall Composition and Disease Resistance. <i>Molecular Plant-Microbe Interactions</i> , 2020, 33, 767-780.	1.4	46
9	Moonlighting Function of Phytochelatin Synthase1 in Extracellular Defense against Fungal Pathogens. <i>Plant Physiology</i> , 2020, 182, 1920-1932.	2.3	26
10	Functional characterization of genes mediating cell wall metabolism and responses to plant cell wall integrity impairment. <i>BMC Plant Biology</i> , 2019, 19, 320.	1.6	20
11	Mitogen-Activated Protein Kinase Phosphatase 1 (MKP1) Negatively Regulates the Production of Reactive Oxygen Species During <i>Arabidopsis</i> Immune Responses. <i>Molecular Plant-Microbe Interactions</i> , 2019, 32, 464-478.	1.4	27
12	Quantitative phosphoproteomic analysis reveals common regulatory mechanisms between effector- and PAMP-triggered immunity in plants. <i>New Phytologist</i> , 2019, 221, 2160-2175.	3.5	102
13	YODA MAP3K kinase regulates plant immune responses conferring broad-spectrum disease resistance. <i>New Phytologist</i> , 2018, 218, 661-680.	3.5	54
14	Plant cell wall-mediated immunity: cell wall changes trigger disease resistance responses. <i>Plant Journal</i> , 2018, 93, 614-636.	2.8	398
15	Non-branched β -1,3-glucan oligosaccharides trigger immune responses in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2018, 93, 34-49.	2.8	112
16	Glutathione Transferase U13 Functions in Pathogen-Triggered Glucosinolate Metabolism. <i>Plant Physiology</i> , 2018, 176, 538-551.	2.3	69
17	Characterization of Plant Cell Wall Damage-Associated Molecular Patterns Regulating Immune Responses. <i>Methods in Molecular Biology</i> , 2017, 1578, 13-23.	0.4	20
18	A computational approach for inferring the cell wall properties that govern guard cell dynamics. <i>Plant Journal</i> , 2017, 92, 5-18.	2.8	62

#	ARTICLE	IF	CITATIONS
19	Alteration of cell wall xylan acetylation triggers defense responses that counterbalance the immune deficiencies of plants impaired in the Î² subunit of the heterotrimeric G protein. <i>Plant Journal</i> , 2017, 92, 386-399.	2.8	68
20	The Arabidopsis leucine-rich repeat receptor kinase MIK2/LRR-KISS connects cell wall integrity sensing, root growth and response to abiotic and biotic stresses. <i>PLoS Genetics</i> , 2017, 13, e1006832.	1.5	187
21	ERECTA and BAK1 Receptor Like Kinases Interact to Regulate Immune Responses in Arabidopsis. <i>Frontiers in Plant Science</i> , 2016, 7, 897.	1.7	99
22	Radiation Damage and Racemic Protein Crystallography Reveal the Unique Structure of the GASA/Snakin Protein Superfamily. <i>Angewandte Chemie</i> , 2016, 128, 8062-8065.	1.6	7
23	Radiation Damage and Racemic Protein Crystallography Reveal the Unique Structure of the GASA/Snakin Protein Superfamily. <i>Angewandte Chemie - International Edition</i> , 2016, 55, 7930-7933.	7.2	45
24	Regulation of Pathogen-Triggered Tryptophan Metabolism in Arabidopsis thaliana by MYB Transcription Factors and Indole Glucosinolate Conversion Products. <i>Molecular Plant</i> , 2016, 9, 682-695.	3.9	149
25	The Arabidopsis NADPH oxidases <i>RbohD</i> and <i>RbohF</i> display differential expression patterns and contributions during plant immunity. <i>Journal of Experimental Botany</i> , 2016, 67, 1663-1676.	2.4	161
26	Expression of fungal acetyl xylan esterase in <i>Arabidopsis thaliana</i> improves saccharification of stem lignocellulose. <i>Plant Biotechnology Journal</i> , 2016, 14, 387-397.	4.1	72
27	Development of a <i>Fusarium oxysporum</i> f. sp. <i>melonis</i> functional GFP fluorescence tool to assist melon resistance breeding programmes. <i>Plant Pathology</i> , 2015, 64, 1349-1357.	1.2	2
28	Mutant Allele-Specific Uncoupling of PENETRATION3 Functions Reveals Engagement of the ATP-Binding Cassette Transporter in Distinct Tryptophan Metabolic Pathways. <i>Plant Physiology</i> , 2015, 168, 814-827.	2.3	71
29	The role of the secondary cell wall in plant resistance to pathogens. <i>Frontiers in Plant Science</i> , 2014, 5, 358.	1.7	455
30	Plant Antimicrobial Peptides Snakin1 and Snakin2: Chemical Synthesis and Insights into the Disulfide Connectivity. <i>Chemistry - A European Journal</i> , 2014, 20, 5102-5110.	1.7	37
31	Functional genomics tools to decipher the pathogenicity mechanisms of the necrotrophic fungus <i>Plectosphaerella cucumerina</i> in <i>Arabidopsis thaliana</i> . <i>Molecular Plant Pathology</i> , 2013, 14, 44-57.	2.0	25
32	Arabidopsis <i>wat1</i> (walls are thin1)-mediated resistance to the bacterial vascular pathogen, <i>Ralstonia solanacearum</i> , is accompanied by cross-regulation of salicylic acid and tryptophan metabolism. <i>Plant Journal</i> , 2013, 73, 225-239.	2.8	154
33	Functional Interplay Between Arabidopsis NADPH Oxidases and Heterotrimeric G Protein. <i>Molecular Plant-Microbe Interactions</i> , 2013, 26, 686-694.	1.4	110
34	Disease resistance or growth: the role of plant hormones in balancing immune responses and fitness costs. <i>Frontiers in Plant Science</i> , 2013, 4, 155.	1.7	505
35	Dissecting Arabidopsis G ¹² Signal Transduction on the Protein Surface. <i>Plant Physiology</i> , 2012, 159, 975-983.	2.3	18
36	Disruption of Abscisic Acid Signaling Constitutively Activates Arabidopsis Resistance to the Necrotrophic Fungus <i>Plectosphaerella cucumerina</i> . <i>Plant Physiology</i> , 2012, 160, 2109-2124.	2.3	132

#	ARTICLE	IF	CITATIONS
37	Arabidopsis Heterotrimeric G-protein Regulates Cell Wall Defense and Resistance to Necrotrophic Fungi. <i>Molecular Plant</i> , 2012, 5, 98-114.	3.9	141
38	Arabidopsis G α protein interactome reveals connections to cell wall carbohydrates and morphogenesis. <i>Molecular Systems Biology</i> , 2011, 7, 532.	3.2	191
39	Autophagy differentially controls plant basal immunity to biotrophic and necrotrophic pathogens. <i>Plant Journal</i> , 2011, 66, 818-830.	2.8	190
40	Plant biotechnology. <i>Current Opinion in Biotechnology</i> , 2010, 21, 182-184.	3.3	1
41	Tryptophan-derived secondary metabolites in <i>Arabidopsis thaliana</i> confer non-host resistance to necrotrophic <i>Plectosphaerella cucumerina</i> fungi. <i>Plant Journal</i> , 2010, 63, no-no.	2.8	191
42	G Proteins and Plant Innate Immunity. <i>Signaling and Communication in Plants</i> , 2010, , 221-250.	0.5	19
43	The ERECTA Receptor-Like Kinase Regulates Cell Wall-Mediated Resistance to Pathogens in <i>Arabidopsis thaliana</i> . <i>Molecular Plant-Microbe Interactions</i> , 2009, 22, 953-963.	1.4	100
44	<i>Leishmania donovani</i> : Thionins, plant antimicrobial peptides with leishmanicidal activity. <i>Experimental Parasitology</i> , 2009, 122, 247-249.	0.5	44
45	Control of the pattern-recognition receptor EFR by an ER protein complex in plant immunity. <i>EMBO Journal</i> , 2009, 28, 3428-3438.	3.5	267
46	A Glucosinolate Metabolism Pathway in Living Plant Cells Mediates Broad-Spectrum Antifungal Defense. <i>Science</i> , 2009, 323, 101-106.	6.0	927
47	Arabidopsis defense response against <i>Fusarium oxysporum</i> . <i>Trends in Plant Science</i> , 2008, 13, 145-150.	4.3	171
48	Repression of the Auxin Response Pathway Increases Arabidopsis Susceptibility to Necrotrophic Fungi. <i>Molecular Plant</i> , 2008, 1, 496-509.	3.9	208
49	Impairment of Cellulose Synthases Required for Arabidopsis Secondary Cell Wall Formation Enhances Disease Resistance. <i>Plant Cell</i> , 2007, 19, 890-903.	3.1	380
50	A Minimalist Design Approach to Antimicrobial Agents Based on a Thionin Template. <i>Journal of Medicinal Chemistry</i> , 2006, 49, 448-451.	2.9	25
51	A Minimalist Approach to Antimicrobial Proteins with Thionin as a Template. , 2006, , 248-251.		0
52	Arabidopsis PEN3/PDR8, an ATP Binding Cassette Transporter, Contributes to Nonhost Resistance to Inappropriate Pathogens That Enter by Direct Penetration. <i>Plant Cell</i> , 2006, 18, 731-746.	3.1	598
53	ERECTA receptor-like kinase and heterotrimeric G protein from Arabidopsis are required for resistance to the necrotrophic fungus <i>Plectosphaerella cucumerina</i> . <i>Plant Journal</i> , 2005, 43, 165-180.	2.8	303
54	A protective role for the embryo surrounding region of the maize endosperm, as evidenced by the characterisation of ZmESR-6, a defensin gene specifically expressed in this region. <i>Plant Molecular Biology</i> , 2005, 58, 269-282.	2.0	79

#	ARTICLE	IF	CITATIONS
55	Structural Dissection of a Highly Knotted Peptide Reveals Minimal Motif with Antimicrobial Activity. <i>Journal of Biological Chemistry</i> , 2005, 280, 1661-1668.	1.6	32
56	Pre- and Postinvasion Defenses Both Contribute to Nonhost Resistance in Arabidopsis. <i>Science</i> , 2005, 310, 1180-1183.	6.0	753
57	Ethylene Response Factor 1 Mediates Arabidopsis Resistance to the Soilborne Fungus <i>Fusarium oxysporum</i> . <i>Molecular Plant-Microbe Interactions</i> , 2004, 17, 763-770.	1.4	268
58	Synthetic and structural studies on <i>Pyricularia puberathionin</i> : a single-residue mutation enhances activity against Gram-negative bacteria. <i>FEBS Letters</i> , 2003, 536, 215-219.	1.3	43
59	Snakin-2, an Antimicrobial Peptide from Potato Whose Gene Is Locally Induced by Wounding and Responds to Pathogen Infection. <i>Plant Physiology</i> , 2002, 128, 951-961.	2.3	289
60	Constitutive expression of ETHYLENE-RESPONSE-FACTOR1 in Arabidopsis confers resistance to several necrotrophic fungi. <i>Plant Journal</i> , 2002, 29, 23-32.	2.8	689
61	Antibiotic activities of peptides, hydrogen peroxide and peroxyxynitrite in plant defence. <i>FEBS Letters</i> , 2001, 498, 219-222.	1.3	90
62	Purification and Characterization of a Mannan-Binding Lectin Specifically Expressed in Corms of Saffron Plant (<i>Crocus sativus</i> L.). <i>Journal of Agricultural and Food Chemistry</i> , 2000, 48, 457-463.	2.4	22
63	Expression of Uroporphyrinogen Decarboxylase or Coproporphyrinogen Oxidase Antisense RNA in Tobacco Induces Pathogen Defense Responses Conferring Increased Resistance to Tobacco Mosaic Virus. <i>Journal of Biological Chemistry</i> , 1999, 274, 4231-4238.	1.6	94
64	Inhibition of protoporphyrinogen oxidase expression in Arabidopsis causes a lesion-mimic phenotype that induces systemic acquired resistance. <i>Plant Journal</i> , 1999, 17, 667-678.	2.8	123
65	Snakin-1, a Peptide from Potato That Is Active Against Plant Pathogens. <i>Molecular Plant-Microbe Interactions</i> , 1999, 12, 16-23.	1.4	281
66	Wheat Genes Encoding Two Types of PR-1 Proteins Are Pathogen Inducible, but Do Not Respond to Activators of Systemic Acquired Resistance. <i>Molecular Plant-Microbe Interactions</i> , 1999, 12, 53-58.	1.4	117
67	Plant defense peptides. , 1998, 47, 479-491.		448
68	Interaction of wheat θ -thionin with large unilamellar vesicles. <i>Protein Science</i> , 1998, 7, 2567-2577.	3.1	23
69	Novel defensin subfamily from spinach (<i>Spinacia oleracea</i>). <i>FEBS Letters</i> , 1998, 435, 159-162.	1.3	141
70	Impaired Fungicide Activity in Plants Blocked in Disease Resistance Signal Transduction. <i>Plant Cell</i> , 1998, 10, 1903-1914.	3.1	88
71	Differential effects of five types of antipathogenic plant peptides on model membranes. <i>FEBS Letters</i> , 1997, 410, 338-342.	1.3	74
72	Differential expression of pathogen-responsive genes encoding two types of glycine-rich proteins in barley. <i>Plant Molecular Biology</i> , 1997, 33, 803-810.	2.0	62

#	ARTICLE	IF	CITATIONS
73	Enhanced tolerance to bacterial pathogens caused by the transgenic expression of barley lipid transfer protein LTP2. <i>Plant Journal</i> , 1997, 12, 669-675.	2.8	196
74	Engineering plants against pathogens: A general strategy. <i>Field Crops Research</i> , 1996, 45, 79-84.	2.3	5
75	Two cold-inducible genes encoding lipid transfer protein LTP4 from barley show differential responses to bacterial pathogens. <i>Molecular Genetics and Genomics</i> , 1996, 252, 162-168.	2.4	73
76	Systemic Acquired Resistance. <i>Plant Cell</i> , 1996, 8, 1809.	3.1	583
77	Systemic Acquired Resistance.. <i>Plant Cell</i> , 1996, 8, 1809-1819.	3.1	1,536
78	The defensive role of nonspecific lipid-transfer proteins in plants. <i>Trends in Microbiology</i> , 1995, 3, 72-74.	3.5	333
79	Developmental and pathogen-induced expression of three barley genes encoding lipid transfer proteins. <i>Plant Journal</i> , 1993, 4, 983-991.	2.8	174
80	Purification, characterization, and cell wall localization of an α -fucosidase that inactivates a xyloglucan oligosaccharin. <i>Plant Journal</i> , 1993, 3, 415-426.	2.8	88
81	Expression of the α -thionin gene from barley in tobacco confers enhanced resistance to bacterial pathogens. <i>Plant Journal</i> , 1993, 3, 457-462.	2.8	156
82	Lipid transfer proteins (nsLTPs) from barley and maize leaves are potent inhibitors of bacterial and fungal plant pathogens. <i>FEBS Letters</i> , 1993, 316, 119-122.	1.3	347
83	Inhibition of bacterial and fungal plant pathogens by thionins of types I and II. <i>Plant Science</i> , 1993, 92, 169-177.	1.7	79