

# GÃtz Hannes Welsch

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8684396/publications.pdf>

Version: 2024-02-01

24  
papers

1,303  
citations

430874

18  
h-index

677142

22  
g-index

24  
all docs

24  
docs citations

24  
times ranked

1160  
citing authors

#	ARTICLE	IF	CITATIONS
1	Changes in T2 Relaxation Time Mapping of Intervertebral Discs Adjacent to Vertebrae after Kyphoplasty Correlate with the Physical Clinical Outcome of Patients. <i>Diagnostics</i> , 2022, 12, 605.	2.6	1
2	Regions at Risk in the Knee Joint of Young Professional Soccer Players: Longitudinal Evaluation of Early Cartilage Degeneration by Quantitative T2 Mapping in 3 T MRI. <i>Cartilage</i> , 2021, 13, 595S-603S.	2.7	10
3	Quantitative T2 Mapping Shows Increased Degeneration in Adjacent Intervertebral Discs Following Kyphoplasty. <i>Cartilage</i> , 2020, 11, 152-159.	2.7	22
4	Using Cartilage MRI T2-Mapping to Analyze Early Cartilage Degeneration in the Knee Joint of Young Professional Soccer Players. <i>Cartilage</i> , 2019, 10, 288-298.	2.7	31
5	Optimized cartilage visualization using 7-T sodium ( <sup>23</sup> Na) imaging after patella dislocation. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2016, 24, 1601-1609.	4.2	10
6	Cartilage Repair Surgery: Outcome Evaluation by Using Noninvasive Cartilage Biomarkers Based on Quantitative MRI Techniques?. <i>BioMed Research International</i> , 2014, 2014, 1-17.	1.9	46
7	dGEMRIC MAPPING OF KNEE JOINT REPAIR. , 2014, , 177-195.		0
8	Magnetic resonance imaging of the knee at 3 and 7 Tesla: a comparison using dedicated multi-channel coils and optimised 2D and 3D protocols. <i>European Radiology</i> , 2012, 22, 1852-1859.	4.5	50
9	Effect of short-term unloading on T2 relaxation time in the lumbar intervertebral disc in vivo magnetic resonance imaging study at 3.0 tesla. <i>Spine Journal</i> , 2012, 12, 257-264.	1.3	24
10	Biochemical (T2, T2* and magnetisation transfer ratio) MRI of knee cartilage: feasibility at ultra-high field (7T) compared with high field (3T) strength. <i>European Radiology</i> , 2011, 21, 1136-1143.	4.5	68
11	Advanced morphological 3D magnetic resonance observation of cartilage repair tissue (MOCART) scoring using a new isotropic 3D proton density, turbo spin echo sequence with variable flip angle distribution (PD-SPACE) compared to an isotropic 3D steady-state free precession sequence (TrueFISP) and standard 2D sequences. <i>Journal of Magnetic Resonance Imaging</i> , 2011, 33, 180-188.	3.4	44
12	Morphological and biochemical T2 evaluation of cartilage repair tissue based on a hybrid double echo at steady state (DESS-T2d) approach. <i>Journal of Magnetic Resonance Imaging</i> , 2011, 34, 895-903.	3.4	17
13	Magnetic Resonance Imaging of Cartilage Repair. <i>Cartilage</i> , 2011, 2, 5-26.	2.7	72
14	T2 and T2* mapping in patients after matrix-associated autologous chondrocyte transplantation: initial results on clinical use with 3.0-Tesla MRI. <i>European Radiology</i> , 2010, 20, 1515-1523.	4.5	59
15	Quantitative T2 Mapping of Knee Cartilage: Differentiation of Healthy Control Cartilage and Cartilage Repair Tissue in the Knee with Unloading Initial Results. <i>Radiology</i> , 2010, 254, 818-826.	7.3	110
16	Tibial Cartilage Hypertrophy Due to Matrix-Associated Autologous Chondrocyte Transplantation of the Medial Femoral Condyle. <i>Journal of Bone and Joint Surgery - Series A</i> , 2009, 91, 1996-2001.	3.0	4
17	Quantitative T2 mapping during follow-up after matrix-associated autologous chondrocyte transplantation (MACT): Full-thickness and zonal evaluation to visualize the maturation of cartilage repair tissue. <i>Journal of Orthopaedic Research</i> , 2009, 27, 957-963.	2.3	69
18	Rapid estimation of cartilage T2 based on double echo at steady state (DESS) with 3 Tesla. <i>Magnetic Resonance in Medicine</i> , 2009, 62, 544-549.	3.0	77

#	ARTICLE	IF	CITATIONS
19	Evaluation and comparison of cartilage repair tissue of the patella and medial femoral condyle by using morphological MRI and biochemical zonal T2 mapping. <i>European Radiology</i> , 2009, 19, 1253-1262.	4.5	56
20	Three-Dimensional Magnetic Resonance Observation of Cartilage Repair Tissue (MOCART) Score Assessed With an Isotropic Three-Dimensional True Fast Imaging With Steady-State Precession Sequence at 3.0 Tesla. <i>Investigative Radiology</i> , 2009, 44, 603-612.	6.2	102
21	High-resolution morphological and biochemical imaging of articular cartilage of the ankle joint at 3.0 T using a new dedicated phased array coil: in vivo reproducibility study. <i>Skeletal Radiology</i> , 2008, 37, 519-526.	2.0	43
22	Magnetization transfer contrast and T2 mapping in the evaluation of cartilage repair tissue with 3T MRI. <i>Journal of Magnetic Resonance Imaging</i> , 2008, 28, 979-986.	3.4	69
23	Cartilage T2 Assessment at 3-T MR Imaging: In Vivo Differentiation of Normal Hyaline Cartilage from Reparative Tissue after Two Cartilage Repair Procedures—Initial Experience. <i>Radiology</i> , 2008, 247, 154-161.	7.3	189
24	In Vivo Biochemical 7.0 Tesla Magnetic Resonance. <i>Investigative Radiology</i> , 2008, 43, 619-626.	6.2	130