

# Bi-Jun Huang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/868883/publications.pdf>

Version: 2024-02-01

34  
papers

1,151  
citations

430874

18  
h-index

395702

33  
g-index

38  
all docs

38  
docs citations

38  
times ranked

2078  
citing authors

#	ARTICLE	IF	CITATIONS
1	Tumor-derived exosomes promote tumor progression and T-cell dysfunction through the regulation of enriched exosomal microRNAs in human nasopharyngeal carcinoma. <i>Oncotarget</i> , 2014, 5, 5439-5452.	1.8	303
2	Epstein-Barr Virus_Encoded LMP1 Upregulates MicroRNA-21 to Promote the Resistance of Nasopharyngeal Carcinoma Cells to Cisplatin-Induced Apoptosis by Suppressing PDCD4 and Fas-L. <i>PLoS ONE</i> , 2013, 8, e78355.	2.5	73
3	Cancer stem-like cell characteristics induced by EB virus-encoded LMP1 contribute to radioresistance in nasopharyngeal carcinoma by suppressing the p53-mediated apoptosis pathway. <i>Cancer Letters</i> , 2014, 344, 260-271.	7.2	70
4	WNT5A promotes stemness characteristics in nasopharyngeal carcinoma cells leading to metastasis and tumorigenesis. <i>Oncotarget</i> , 2015, 6, 10239-10252.	1.8	67
5	Nuclear PGK1 Alleviates ADP-Dependent Inhibition of CDC7 to Promote DNA Replication. <i>Molecular Cell</i> , 2018, 72, 650-660.e8.	9.7	57
6	Promoting tumorigenesis in nasopharyngeal carcinoma, NEDD8 serves as a potential theranostic target. <i>Cell Death and Disease</i> , 2017, 8, e2834-e2834.	6.3	47
7	SPINK6 Promotes Metastasis of Nasopharyngeal Carcinoma via Binding and Activation of Epithelial Growth Factor Receptor. <i>Cancer Research</i> , 2017, 77, 579-589.	0.9	47
8	Urokinase-type plasminogen activator receptor signaling is critical in nasopharyngeal carcinoma cell growth and metastasis. <i>Cell Cycle</i> , 2014, 13, 1958-1969.	2.6	44
9	GTSE1 is involved in breast cancer progression in p53 mutation-dependent manner. <i>Journal of Experimental and Clinical Cancer Research</i> , 2019, 38, 152.	8.6	42
10	Along with its favorable prognostic role, CLCA2 inhibits growth and metastasis of nasopharyngeal carcinoma cells via inhibition of FAK/ERK signaling. <i>Journal of Experimental and Clinical Cancer Research</i> , 2018, 37, 34.	8.6	33
11	The developmental transcription factor IRF6 attenuates ABCG2 gene expression and distinctively reverses stemness phenotype in nasopharyngeal carcinoma. <i>Cancer Letters</i> , 2018, 431, 230-243.	7.2	31
12	The Upregulation of Trophinin-Associated Protein (TROAP) Predicts a Poor Prognosis in Hepatocellular Carcinoma. <i>Journal of Cancer</i> , 2019, 10, 957-967.	2.5	31
13	S100A14 suppresses metastasis of nasopharyngeal carcinoma by inhibition of NF- $\kappa$ B signaling through degradation of IRAK1. <i>Oncogene</i> , 2020, 39, 5307-5322.	5.9	30
14	CDC42-interacting protein 4 promotes metastasis of nasopharyngeal carcinoma by mediating invadopodia formation and activating EGFR signaling. <i>Journal of Experimental and Clinical Cancer Research</i> , 2017, 36, 21.	8.6	26
15	<i>RASSF6</i> promotes p21 <sup>Cip1/Waf1</sup> -dependent cell cycle arrest and apoptosis through activation of the JNK/SAPK pathway in clear cell renal cell carcinoma. <i>Cell Cycle</i> , 2014, 13, 1440-1449.	2.6	24
16	PDZ binding kinase (PBK) is a theranostic target for nasopharyngeal carcinoma: driving tumor growth via ROS signaling and correlating with patient survival. <i>Oncotarget</i> , 2016, 7, 26604-26616.	1.8	23
17	IL-8 suppresses E-cadherin expression in nasopharyngeal carcinoma cells by enhancing E-cadherin promoter DNA methylation. <i>International Journal of Oncology</i> , 2016, 48, 207-214.	3.3	22
18	ETV4 is a theranostic target in clear cell renal cell carcinoma that promotes metastasis by activating the pro-metastatic gene FOSL1 in a PI3K-AKT dependent manner. <i>Cancer Letters</i> , 2020, 482, 74-89.	7.2	19

#	ARTICLE	IF	CITATIONS
19	LACTB promotes metastasis of nasopharyngeal carcinoma via activation of ERBB3/EGFR-ERK signaling resulting in unfavorable patient survival. <i>Cancer Letters</i> , 2021, 498, 165-177.	7.2	19
20	A comparison of weekly versus 3-weekly cisplatin during concurrent chemoradiotherapy for locoregionally advanced nasopharyngeal carcinoma using intensity modulated radiation therapy: a matched study. <i>Journal of Cancer</i> , 2018, 9, 92-99.	2.5	17
21	PTPN3 Inhibits the Growth and Metastasis of Clear Cell Renal Cell Carcinoma via Inhibition of PI3K/AKT Signaling. <i>Molecular Cancer Research</i> , 2020, 18, 903-912.	3.4	17
22	Long-Term Toxicity Studies in Canine of E10A, An Adenoviral Vector for Human Endostatin Gene. <i>Human Gene Therapy</i> , 2007, 18, 207-221.	2.7	16
23	&lt;p&gt;Candidate tumor suppressor gene IRF6 is involved in human breast cancer pathogenesis via modulating PI3K-regulatory subunit PIK3R2 expression&lt;/p&gt;. <i>Cancer Management and Research</i> , 2019, Volume 11, 5557-5572.	1.9	14
24	Antioxidants suppress radiation-induced apoptosis via inhibiting MAPK pathway in nasopharyngeal carcinoma cells. <i>Biochemical and Biophysical Research Communications</i> , 2020, 527, 770-777.	2.1	14
25	Downregulation of Ras Association Domain Family Member 6 (RASSF6) Underlies the Treatment Resistance of Highly Metastatic Nasopharyngeal Carcinoma Cells. <i>PLoS ONE</i> , 2014, 9, e100843.	2.5	13
26	Concurrent Chemoradiotherapy versus Intensity-modulated Radiotherapy Alone for Elderly Nasopharyngeal Carcinoma Patients with Pre-treatment Epstein-Barr Virus DNA: A Cohort Study in an Endemic Area with Long-term Follow-up. <i>Journal of Cancer</i> , 2018, 9, 3023-3031.	2.5	10
27	AKR1C2 acts as a targetable oncogene in esophageal squamous cell carcinoma via activating PI3K/AKT signaling pathway. <i>Journal of Cellular and Molecular Medicine</i> , 2020, 24, 9999-10012.	3.6	10
28	Global expression profiling and pathway analysis of mouse mammary tumor reveals strain and stage specific dysregulated pathways in breast cancer progression. <i>Cell Cycle</i> , 2018, 17, 963-973.	2.6	6
29	Metastasis of nasopharyngeal carcinoma: What we know and do not know. <i>Visualized Cancer Medicine</i> , 2021, 2, 4.	0.9	6
30	Geographical disparities in the prognosis of patients with nasopharyngeal carcinoma treated with intensity-modulated radiation therapy: a large institution-based cohort study from an endemic area. <i>BMJ Open</i> , 2020, 10, e037150.	1.9	4
31	p53, latent membrane protein 1, bcl-2, and prognosis in nasopharyngeal carcinoma: a meta-analysis. <i>Histology and Histopathology</i> , 2019, 34, 103-110.	0.7	4
32	ACE2 in tumor cells and tumor vasculature: Negligible intercellular transfer from cancer cells into endothelial cells. <i>Visualized Cancer Medicine</i> , 2021, 2, 3.	0.9	2
33	The autoregulatory serglycin/CD44 axis drives stemness-like phenotypes in TNBC in a $\beta$ -catenin-dependent manner. <i>Clinical and Translational Medicine</i> , 2021, 11, e311.	4.0	2
34	Observation on effect of treatment of alcoholic fatty liver by traditional medical therapy of liver-clearing, dampness-removing and collaterals-dredging. , 2002, 8, 134-136.		1