Sarah A Moore

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8692976/publications.pdf

Version: 2024-02-01

20 papers

646 citations

623188 14 h-index 752256 20 g-index

24 all docs

24 docs citations

times ranked

24

1173 citing authors

#	Article	IF	CITATIONS
1	Physical Activity, Sedentary Behaviour and Metabolic Control following Stroke: A Cross-Sectional and Longitudinal Study. PLoS ONE, 2013, 8, e55263.	1.1	109
2	Effects of Community Exercise Therapy on Metabolic, Brain, Physical, and Cognitive Function Following Stroke. Neurorehabilitation and Neural Repair, 2015, 29, 623-635.	1.4	102
3	Non-alcoholic fatty liver disease is associated with higher levels of <i>objectively </i> measured sedentary behaviour and lower levels of physical activity than matched healthy controls. Frontline Gastroenterology, 2015, 6, 44-51.	0.9	91
4	Measuring Energy Expenditure After Stroke. Stroke, 2012, 43, 1660-1662.	1.0	41
5	Exercise as a treatment for sarcopenia: an umbrella review of systematic review evidence. Physiotherapy, 2020, 107, 189-201.	0.2	38
6	Comprehensive measurement of stroke gait characteristics with a single accelerometer in the laboratory and community: a feasibility, validity and reliability study. Journal of NeuroEngineering and Rehabilitation, 2017, 14, 130.	2.4	35
7	How should long-term free-living physical activity be targeted after stroke? A systematic review and narrative synthesis. International Journal of Behavioral Nutrition and Physical Activity, 2018, 15, 100.	2.0	34
8	Comparison of cardiac output determined by bioimpedance and bioreactance methods at rest and during exercise. Journal of Clinical Monitoring and Computing, 2012, 26, 63-68.	0.7	31
9	Gait Asymmetry Post-Stroke: Determining Valid and Reliable Methods Using a Single Accelerometer Located on the Trunk. Sensors, 2020, 20, 37.	2.1	29
10	Discrepancy Between Cardiac and Physical Functional Reserves in Stroke. Stroke, 2012, 43, 1422-1425.	1.0	24
11	Wristband Accelerometers to motiVate arm Exercises after Stroke (WAVES): a pilot randomized controlled trial. Clinical Rehabilitation, 2019, 33, 1391-1403.	1.0	24
12	Self-directed therapy programmes for arm rehabilitation after stroke: a systematic review. Clinical Rehabilitation, 2018, 32, 1022-1036.	1.0	23
13	Wristband Accelerometers to motiVate arm Exercise after Stroke (WAVES): study protocol for a pilot randomized controlled trial. Trials, 2016, 17, 508.	0.7	20
14	A study of physical activity comparing people with Charcot-Marie-Tooth disease to normal control subjects. Disability and Rehabilitation, 2017, 39, 1753-1758.	0.9	19
15	Exercise Induces Peripheral Muscle But Not Cardiac Adaptations After Stroke: A Randomized Controlled Pilot Trial. Archives of Physical Medicine and Rehabilitation, 2016, 97, 596-603.	0.5	12
16	A feasibility, acceptability and fidelity study of a multifaceted behaviour change intervention targeting free-living physical activity and sedentary behaviour in community dwelling adult stroke survivors. Pilot and Feasibility Studies, 2020, 6, 58.	0.5	8
17	Auditory rhythmical cueing to improve gait and physical activity in community-dwelling stroke survivors (ACTIVATE): study protocol for a pilot randomised controlled trial. Pilot and Feasibility Studies, 2020, 6, 68.	0.5	2
18	Using intervention mapping to develop and facilitate implementation of a multifaceted behavioural intervention targeting physical activity and sedentary behaviour in stroke survivors: Physical Activity Routines After Stroke (PARAS): intervention development study. Health Psychology and Behavioral Medicine, 2022, 10, 439-466.	0.8	2

#	# Article	IF	CITATIONS
19	Physical activity, sedentary behaviour and energy expenditure post-stroke. Physical The 2015, 20, 264-265.	nerapy Reviews, 0.3	1
20	Acceptability and deliverability of an auditory rhythmical cueing (ARC) training progra home and outdoors to improve gait and physical activity post-stroke. Archives of Phys 12, 1.	amme for use at siotherapy, 2022, 0.7	1