Qingyu Zeng

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8698810/publications.pdf

Version: 2024-02-01



#	Article	IF	CITATIONS
1	Rapamycin inhibits BAFF-stimulated cell proliferation and survival by suppressing mTOR-mediated PP2A-Erk1/2 signaling pathway in normal and neoplastic B-lymphoid cells. Cellular and Molecular Life Sciences, 2015, 72, 4867-4884.	2.4	42
2	BAFF inhibits autophagy promoting cell proliferation and survival by activating Ca2+-CaMKII-dependent Akt/mTOR signaling pathway in normal and neoplastic B-lymphoid cells. Cellular Signalling, 2019, 53, 68-79.	1.7	29
3	BAFF activates Erk1/2 promoting cell proliferation and survival by Ca2+-CaMKII-dependent inhibition of PP2A in normal and neoplastic B-lymphoid cells. Biochemical Pharmacology, 2014, 87, 332-343.	2.0	20
4	Rapamycin attenuates BAFFâ€extended proliferation and survival via disruption of mTORC1/2 signaling in normal and neoplastic B″ymphoid cells. Journal of Cellular Physiology, 2018, 233, 516-529.	2.0	20
5	PD-L1 blockade potentiates the antitumor effects of ALA-PDT and optimizes the tumor microenvironment in cutaneous squamous cell carcinoma. Oncolmmunology, 2022, 11, 2061396.	2.1	20
6	Antibacterial photodynamic therapy mediated by 5-aminolevulinic acid on methicillin-resistant Staphylococcus aureus. Photodiagnosis and Photodynamic Therapy, 2019, 28, 330-337.	1.3	19
7	Single-cell transcriptomic analysis reveals the critical molecular pattern of UV-induced cutaneous squamous cell carcinoma. Cell Death and Disease, 2022, 13, 23.	2.7	19
8	Rapamycin inhibits B-cell activating factor (BAFF)-stimulated cell proliferation and survival by suppressing Ca2+-CaMKII-dependent PTEN/Akt-Erk1/2 signaling pathway in normal and neoplastic B-lymphoid cells. Cell Calcium, 2020, 87, 102171.	1.1	18
9	5â€Aminolaevulinic acid photodynamic therapy amplifies intense inflammatory response in the treatment of acne vulgaris via CXCL8. Experimental Dermatology, 2021, 30, 923-931.	1.4	18
10	IL-2, IL-4, IFN-γ or TNF-α enhances BAFF-stimulated cell viability and survival by activating Erk1/2 and S6K1 pathways in neoplastic B-lymphoid cells. Cytokine, 2016, 84, 37-46.	1.4	14
11	Efficacy of the therapy of 5-aminolevulinic acid photodynamic therapy combined with human umbilical cord mesenchymal stem cells on methicillin-resistant Staphylococcus aureus-infected wound in a diabetic mouse model. Photodiagnosis and Photodynamic Therapy, 2021, 36, 102480.	1.3	14
12	Modified 5-aminolevulinic acid photodynamic therapy (M-PDT) inhibits cutaneous squamous cell carcinoma cell proliferation via targeting PP2A/PP5-mediated MAPK signaling pathway. International Journal of Biochemistry and Cell Biology, 2021, 137, 106036.	1.2	12
13	Celastrol inhibits LL37-induced rosacea by inhibiting Ca2+/CaMKII-mTOR-NF-ήB activation. Biomedicine and Pharmacotherapy, 2022, 153, 113292.	2.5	9
14	Modified 5â€aminolevulinic acid photodynamic therapy reduces pain and improves therapeutic effects in cutaneous squamous cell carcinoma mouse model. Lasers in Surgery and Medicine, 2022, 54, 804-812.	1.1	8
15	5-Aminolaevulinic acid photodynamic therapy suppresses lipid secretion of primary sebocytes through AMPK/SREBP-1 pathway. Photodiagnosis and Photodynamic Therapy, 2021, 36, 102537.	1.3	7
16	Ruyi Jinhuang Powder accelerated diabetic ulcer wound healing by regulating Wnt/β-catenin signaling pathway of fibroblasts In Vivo and In Vitro. Journal of Ethnopharmacology, 2022, 293, 115321.	2.0	7
17	Halofuginone enhances the anti-tumor effect of ALA-PDT by suppressing NRF2 signaling in cSCC. Photodiagnosis and Photodynamic Therapy, 2021, 37, 102572.	1.3	3
18	Molecular characterization of gene expression changes in murine cutaneous squamous cell carcinoma after 5-aminolevulinic acid photodynamic therapy. Photodiagnosis and Photodynamic Therapy, 2022, 102907.	1.3	3